

**Submission of Clinical Data Supporting  
Formulary Consideration of**

**AZOR™ (amlodipine and olmesartan medoxomil)**  
**Daiichi Sankyo, Inc.**

**Prepared by:**

**Meenakshi Arunachalam, Pharm. D.  
Associate Director, Professional Affairs  
Daiichi Sankyo, Inc.**

**John Raia, Pharm.D.  
Director, Professional Affairs  
Daiichi Sankyo, Inc.**

**AMCP Formulary Dossier**

**AZOR™ (amlodipine and olmesartan medoxomil)**

**Table of Contents**

	<b>Page</b>
<b>1. Product Information</b>	
<b>1.1. Product Description</b>	
a. Generic, Brand Name, Therapeutic Class	4
b. Dosage Forms/National Drug Code (NDC)/Cost	4
c. Product Labeling (enclosed in binder)	
d. AHFS Drug Classification	4
e. FDA Approved Indication	4
f. Pharmacology	5
g. Pharmacokinetics	5
h. Contraindications	6
i. Warnings and Precautions	6
j. Adverse Reactions	7
k. Drug Interactions	9
l. Dosing and Administration	10
m. Comparative Pharmacokinetics/Pharmacologic Angiotensin Receptor Blockers (ARB)Profiles	10
n. Comparative Pharmacokinetic/Pharmacologic Profiles of the Dihydropyridine Calcium Channel Blockers (CCBs)	12
o. Comparative Pharmacokinetic/Pharmacologic Profiles of Fixed Dose Amlodipine Combination Products	14
<b>1.2. Place of the Product in Therapy</b>	
<b>1.2.1. Disease Description</b>	
a. Epidemiology and Clinical Aspects of Hypertension	17
b. The Role of Calcium and the Renin-Angiotensin System in Hypertension	17
<b>1.2.2. Approaches to Treatment</b>	
a. JNC-7 Treatment Guidelines	18
b. Angiotensin Receptor Blockers in Antihypertensive Therapy	20
c. Calcium Channel Blockers in Antihypertensive Therapy	20
d. Place of Combination Products in Hypertension Therapy	21
<b>1.3 Evidence for Pharmacogenomic Tests and Drugs</b>	22
<b>2. Supporting Clinical and Economic Information</b>	22
<b>2.1 Summarizing Key Clinical Studies</b>	
a. Review of Clinical Studies	22
b. Amlodipine/Olmesartan Medoxomil Abstracts	47
c. Evidence Table Spreadsheets	60

AZOR™ (amlodipine and olmesartan medoxomil)  
Formulary Dossier

**3. Supporting Information**

**3.1. References**

<b>a.</b>	References Contained in Dossier	68
<b>b.</b>	Amlodipine and Olmesartan Medoxomil Clinical Bibliography	70
<b>c.</b>	Olmesartan Medoxomil Clinical Bibliography	71
<b>d.</b>	Olmesartan Medoxomil Pre-Clinical Bibliography	95

AZOR™ (amlodipine and olmesartan medoxomil)  
Formulary Dossier

**1. PRODUCT INFORMATION**

**a. Generic, Brand Name, Therapeutic Class:**

amlodipine and olmesartan medoxomil

AZOR™

Dihydropyridine Calcium Channel Blocker and Selective AT<sub>1</sub> subtype Angiotensin II Receptor Antagonist.

**b. Dosage Forms/National Drug Code (NDC)/Cost:**

AZOR™ tablets are differentiated by tablet color/size and are debossed with an individual product tablet code on one side. AZOR™ tablets are supplied for oral administration in the following strength and package configurations listed in Table 1.

**Table 1: Tablets are supplied as follows (includes NDC codes and WAC Cost):**

Tablet Strength (amlodipine equivalent/olmesartan medoxomil) mg	Package Configuration	NDC#	WAC Tablet Price	WAC Package Price
5/20 mg	Bottle of 30	65597-110-30	\$2.13	\$63.90
	Bottle of 90	65597-110-90	\$2.13	\$191.70
	10 blisters of 10	65597-110-10	\$2.13	\$213.00
	Bottle of 1000	65597-110-11		
10/20 mg	Bottle of 30	65597-111-30	\$2.41	\$72.30
	Bottle of 90	65597-111-90	\$2.41	\$216.90
	10 blisters of 10	65597-111-10	\$2.41	\$241.00
	Bottle of 1000	65597-111-11		
5/40 mg	Bottle of 30	65597-112-30	\$2.70	\$81.00
	Bottle of 90	65597-112-90	\$2.70	\$243.00
	10 blisters of 10	65597-112-10	\$2.70	\$270.00
	Bottle of 1000	65597-112-11		
10/40 mg	Bottle of 30	65597-113-30	\$3.06	\$91.80
	Bottle of 90	65597-113-90	\$3.06	\$275.40
	10 blisters of 10	65597-113-10	\$3.06	\$306.00
	Bottle of 1000	65597-113-11		

**c. Product Labeling:** Enclosed in binder

**d. AHFS Drug Classification:**

Amlodipine - 24:04

Olmesartan – 24:08

**e. FDA Approved Indication:** AZOR™ is indicated for the treatment of hypertension, alone or with other antihypertensive agents. This fixed combination drug is not indicated for the initial therapy of hypertension  
Date of approval: September 26, 2007.

## AZOR™ (amlodipine and olmesartan medoxomil) Formulary Dossier

### e. Pharmacology:

AZOR™ is a combination of two antihypertensive drugs: a dihydropyridine calcium antagonist (calcium ion antagonist or slow-channel blocker), amlodipine besylate, and an angiotensin II receptor blocker, olmesartan medoxomil. The amlodipine component of AZOR™ inhibits the transmembrane influx of calcium ions into vascular smooth muscle and cardiac muscle, and the olmesartan medoxomil component of AZOR™ blocks the vasoconstrictor effects of angiotensin II.

**Amlodipine.** Experimental data suggests that amlodipine binds to both dihydropyridine and nondihydropyridine binding sites. The contractile processes of cardiac muscle and vascular smooth muscle are dependent upon the movement of extracellular calcium ions into these cells through specific ion channels. Amlodipine inhibits calcium ion influx across cell membranes selectively, with a greater effect on vascular smooth muscle cells than on cardiac muscle cells. Negative inotropic effects can be detected *in vitro* but such effects have not been seen in intact animals at therapeutic doses. Serum calcium concentration is not affected by amlodipine. Within the physiologic pH range, amlodipine is an ionized compound ( $pK_a=8.6$ ), and its kinetic interaction with the calcium channel receptor is characterized by a gradual rate of association and dissociation with the receptor binding site, resulting in a gradual onset of effect.

Amlodipine is a peripheral arterial vasodilator that acts directly on vascular smooth muscle to cause a reduction in peripheral vascular resistance and reduction in blood pressure.

**Olmesartan medoxomil.** Angiotensin II is formed from angiotensin I in a reaction catalyzed by angiotensin converting enzyme (ACE, kininase II). Angiotensin II is the principal pressor agent of the renin-angiotensin system with effects that include vasoconstriction, stimulation of synthesis and release of aldosterone, cardiac stimulation and renal reabsorption of sodium. Olmesartan blocks the vasoconstrictor effects of angiotensin II by selectively blocking the binding of angiotensin II to the AT<sub>1</sub> receptor in vascular smooth muscle. Its action is, therefore, independent of the pathways for angiotensin II synthesis.

An AT<sub>2</sub> receptor is found also in many tissues, but this receptor is not known to be associated with cardiovascular homeostasis. Olmesartan has more than a 12,500-fold greater affinity for the AT<sub>1</sub> receptor than for the AT<sub>2</sub> receptor.

Blockade of the renin-angiotensin system with ACE inhibitors, which inhibit the biosynthesis of angiotensin II from angiotensin I, is a mechanism of many drugs used to treat hypertension. ACE inhibitors also inhibit the degradation of bradykinin, a reaction also catalyzed by ACE. Because olmesartan does not inhibit ACE (kininase II), it does not affect the response to bradykinin. Whether this difference has clinical relevance is not yet known.

Blockade of the angiotensin II receptor inhibits the negative regulatory feedback of angiotensin II on renin secretion, but the resulting increased plasma renin activity and circulating angiotensin II levels do not overcome the effect of olmesartan on blood pressure.

### f. Pharmacokinetics:

The pharmacokinetics of amlodipine and olmesartan medoxomil from AZOR™ are equivalent to the pharmacokinetics of amlodipine and olmesartan medoxomil when administered separately. The bioavailability of both components is well below 100%, but neither component is affected by food. The effective half-lives of amlodipine ( $45\pm11$  hours) and olmesartan ( $7\pm1$  hours) result in a 2- to 3- fold accumulation for amlodipine and negligible accumulation for olmesartan with once-daily dosing.

**Amlodipine.** After oral administration of therapeutic doses of amlodipine, absorption produces peak plasma concentrations between 6 and 12 hours. Absolute bioavailability is estimated as between 64% and 90%.

**Olmesartan medoxomil.** Olmesartan medoxomil is rapidly and completely bioactivated by ester hydrolysis to olmesartan during absorption from the gastrointestinal tract. The absolute bioavailability of olmesartan medoxomil is approximately 26%. After oral administration, the peak plasma concentration ( $C_{max}$ ) of olmesartan is reached after 1 to 2 hours. Food does not affect the bioavailability of olmesartan medoxomil.

#### **Distribution**

**Amlodipine.** Ex vivo studies have shown that approximately 93% of the circulating drug is bound to plasma proteins in hypertensive patients. Steady-state plasma levels of amlodipine are reached after 7 to 8 days of consecutive daily dosing.

**Olmesartan medoxomil.** The volume of distribution of olmesartan is approximately 17 L. Olmesartan is highly bound to plasma proteins (99%) and does not penetrate red blood cells. The protein binding is constant at plasma olmesartan concentrations well above the range achieved with recommended doses.

## AZOR™ (amlodipine and olmesartan medoxomil) Formulary Dossier

In rats, olmesartan crossed the blood-brain barrier poorly, if at all. Olmesartan passed across the placental barrier in rats and was distributed to the fetus. Olmesartan was distributed to milk at low levels in rats.

### **Metabolism and Excretion:**

**Amlodipine.** Amlodipine is extensively (about 90%) converted to inactive metabolites via hepatic metabolism. Elimination from the plasma is biphasic with a terminal elimination half-life of about 30 to 50 hours. Ten percent of the parent compound and 60% of the metabolites are excreted in the urine.

**Olmesartan medoxomil.** Following the rapid and complete conversion of olmesartan medoxomil to olmesartan during absorption, there is virtually no further metabolism of olmesartan. Total plasma clearance of olmesartan is 1.3 L/h with a renal clearance of 0.6 L/h. Approximately 35% to 50% of the absorbed dose is recovered in urine while the remainder is eliminated in feces via the bile.

Olmesartan appears to be eliminated in a biphasic manner with a terminal elimination half-life of approximately 13 hours. Olmesartan shows linear pharmacokinetics following single oral doses of up to 320 mg and multiple oral doses of up to 80 mg. Steady-state levels of olmesartan are achieved within 3 to 5 days and no accumulation in plasma occurs with once-daily dosing.

**g. Contraindications:** None.

**h. Warnings and Precautions:** The adverse reactions of AZOR™ are generally related to those of each of its components.

### **Fetal/Neonatal Morbidity and Mortality**

**Olmesartan medoxomil.** Drugs that act directly on the renin-angiotensin system can cause fetal and neonatal morbidity and death when administered to pregnant women. There have been several dozen cases reported in the world literature of patients who were taking angiotensin converting enzyme inhibitors. When pregnancy is detected, AZOR™ should be discontinued as soon as possible.

During the second and third trimesters of pregnancy, these drugs have been associated with fetal injury that includes hypotension, neonatal skull hypoplasia, anuria, reversible or irreversible renal failure, and death. Oligohydramnios has also been reported, presumably resulting from decreased fetal renal function; oligohydramnios in this setting has been associated with fetal limb contractures, craniofacial deformation, and hypoplastic lung development. Prematurity, intrauterine growth retardation, and patent ductus arteriosus have also been reported, although it is not clear whether these occurrences were due to exposure to the drug.

These adverse effects do not appear to have resulted from intrauterine drug exposure that has been limited to the first trimester. Mothers whose embryos and fetuses are exposed to an angiotensin II receptor antagonist only during the first trimester should be so informed. Nonetheless, when patients become pregnant, physicians should have the patient discontinue the use of AZOR™ as soon as possible.

Rarely (probably less often than once in every thousand pregnancies), no alternative to a drug acting on the renin-angiotensin system will be found. In these rare cases, the mothers should be apprised of the potential hazards to their fetuses and serial ultrasound examinations should be performed to assess the intra-amniotic environment.

If oligohydramnios is observed, AZOR™ should be discontinued unless it is considered life-saving for the mother. Contraction stress testing (CST), a non-stress test (NST), or biophysical profiling (BPP) may be appropriate, depending upon the week of pregnancy. Patients and physicians should be aware however, that oligohydramnios may not appear until after the fetus has sustained irreversible injury.

Infants with histories of *in utero* exposure to an angiotensin II receptor antagonist should be closely observed for hypotension, oliguria, and hyperkalemia. If oliguria occurs, attention should be directed toward support of blood pressure and renal perfusion. Exchange transfusion or dialysis may be required as means of reversing hypotension and/or substituting for disordered renal function.

No teratogenic effects were observed when olmesartan medoxomil was administered to pregnant rats at oral doses up to 1000 mg/kg/day (240 times the maximum recommended human dose (MRHD) on a mg/m<sup>2</sup> basis) or pregnant rabbits at oral doses up to 1 mg/kg/day (half the MRHD on a mg/m<sup>2</sup> basis; higher doses could not be evaluated for effects on fetal development as they were lethal to the dose). In rats, significant decreases in pup birth weight and weight gain were observed at doses ≥1.6 mg/kg/day, and delays in developmental milestones (delayed separation of ear auricular, eruption of lower incisors, appearance of abdominal hair, descent of testes, and separation of eyelids) and dose-dependent increase in the incidence of dilation of the renal pelvis were observed at doses ≥8 mg/kg/day. The no observed effect dose for developmental toxicity in rats is 0.3 mg/kg/day, about one-tenth the MRHD of 40 mg/kg/day.

## AZOR™ (amlodipine and olmesartan medoxomil) Formulary Dossier

### ***Hypotension in Volume- or Salt-Depleted Patients***

**Olmesartan medoxomil.** Symptomatic hypotension may occur after initiation of treatment with olmesartan medoxomil. Patients with an activated renin-angiotensin system, such as volume- and/or salt-depleted patients (e.g., those being treated with high doses of diuretics) may be particularly vulnerable. Treatment with AZOR™ should start under close medical supervision. If hypotension does occur, the patient should be placed in the supine position and, if necessary, given an intravenous infusion of normal saline. A transient hypotensive response is not a contraindication to further treatment, which usually can be continued without difficulty once the blood pressure has stabilized.

### ***Vasodilation***

**Amlodipine.** Since the vasodilation attributable to amlodipine in AZOR™ is gradual in onset, acute hypotension has rarely been reported after oral administration. Nonetheless, caution, as with any other peripheral vasodilator, should be exercised when administering AZOR™, particularly in patients with severe aortic stenosis.

### ***Patients with Severe Obstructive Coronary Artery Disease***

Patients, particularly those with severe obstructive coronary artery disease, may develop increased frequency, duration, or severity of angina or acute myocardial infarction on starting calcium channel blocker therapy or at the time of dosage increase. The mechanism of this effect has not been elucidated.

### ***Patients with Congestive Heart Failure***

**Amlodipine.** In general, calcium channel blockers should be used with caution in patients with heart failure. Amlodipine (5-10 mg per day) has been studied in a placebo-controlled trial of 1153 patients with NYHA (New York Heart Association) Class III or IV heart failure on stable doses of ACE inhibitor, digoxin, and diuretics. Follow-up was at least 6 months, with a mean of about 14 months. There was no overall adverse effect on survival or cardiac morbidity (as defined by life-threatening arrhythmia, acute myocardial infarction, or hospitalization for worsened heart failure). Amlodipine has been compared to placebo in four 8-12 week studies of patients with NYHA class II/III heart failure, involving a total of 697 patients. In these studies, there was no evidence of worsening of heart failure based on measures of exercise tolerance, NYHA classification, symptoms, or left ventricular ejection fraction (LVEF).

### ***Patients with Impaired Renal Function***

**Olmesartan medoxomil.** Changes in renal function may be anticipated in susceptible individuals treated with olmesartan medoxomil as a consequence of inhibiting the renin-angiotensin aldosterone system. In patients whose renal function may depend upon the activity of the renin-angiotensin-aldosterone system (e.g., patients with severe congestive heart failure), treatment with angiotensin converting enzyme inhibitors and angiotensin receptor antagonists has been associated with oliguria or progressive azotemia and (rarely) with acute renal failure and/or death. Similar effects may occur in patients treated with AZOR™ due to the olmesartan medoxomil component [See *Clinical Pharmacology* (12.3)].

In studies of ACE inhibitors in patients with unilateral or bilateral renal artery stenosis, increases in serum creatinine or blood urea nitrogen (BUN) have been reported. There has been no long-term use of olmesartan medoxomil in patients with unilateral or bilateral renal artery stenosis, but similar effects would be expected with AZOR™ because of the olmesartan medoxomil component.

### ***Patients with Hepatic Impairment***

**Amlodipine.** Since amlodipine is extensively metabolized by the liver and the plasma elimination half-life ( $t_{1/2}$ ) is 56 hours in patients with severely impaired hepatic function, caution should be exercised when administering AZOR™ to patients with severe hepatic impairment.

### **i. Adverse Reactions:**

#### **Clinical Trials Experience**

Because clinical studies are conducted under widely varying conditions, adverse reaction rates observed in the clinical studies of a drug cannot be directly compared to rates in the clinical studies of another drug and may not reflect the rates observed in practice.

# AZOR™ (amlodipine and olmesartan medoxomil) Formulary Dossier

## **AZOR™**

The data described below reflect exposure to AZOR™ in more than 1600 patients including more than 1000 exposed for at least 6 months and more than 700 exposed for 1 year. AZOR™ was studied in one placebo-controlled factorial trial (see Section 14.1). The population had a mean age of 54 years and included approximately 55% males. Seventy-one percent were Caucasian and 25% were Black. Patients received doses ranging from 5/20 mg to 10/40 mg orally once daily.

The overall incidence of adverse reactions on therapy with AZOR™ was similar to that seen with corresponding doses of the individual components of AZOR™, and to placebo. The reported adverse reactions were generally mild and seldom led to discontinuation of treatment (2.6% for AZOR™ and 6.8% for placebo).

## **Edema**

Edema is a known, dose-dependent adverse effect of amlodipine but not of olmesartan medoxomil.

The placebo-subtracted incidence of edema during the 8-week, randomized, double-blind treatment period was highest with amlodipine 10 mg monotherapy. The incidence was significantly reduced when 20 mg or 40 mg of olmesartan medoxomil was added to the 10 mg amlodipine dose.

**Table 2: Placebo-Subtracted Incidence of Edema during the Double-Blind Treatment Period**

		Olmesartan Medoxomil		
		Placebo	20 mg	40 mg
	<b>Placebo</b>	0%*	(-2.4%)	6.2%
<b>Amlodipine</b>	<b>5 mg</b>	0.7%	5.7%	6.2%
	<b>10 mg</b>	24.5%	13.3%	11.2%
*12.3%=actual placebo incidence				

Across all treatment groups, the frequency of edema was generally higher in women than men, as has been observed in previous studies of amlodipine.

Adverse reactions seen at lower rates during the double-blind period also occurred in the patients treated with AZOR™ at about the same or greater incidence as in patients receiving placebo. These included hypotension, orthostatic hypotension, rash, pruritus, palpitation, urinary frequency, and nocturia.

The adverse event profile obtained from 44 weeks of open-label combination therapy with amlodipine plus olmesartan medoxomil was similar to that observed during the 8-week, double-blind, placebo-controlled period.

## **Amlodipine**

Amlodipine has been evaluated for safety in more than 11,000 patients in U.S. and foreign clinical trials. Most adverse reactions reported during therapy with amlodipine were of mild or moderate severity. In controlled clinical trials directly comparing amlodipine (N=1730) in doses up to 10 mg to placebo (N=1250), discontinuation of amlodipine due to adverse reactions was required in only about 1.5% of amlodipine-treated patients and about 1% of placebo-treated patients. The most common side effects were headache and edema. The incidence (%) of dose-related side effects was as follows:

**Table 3: Incidence (%) of dose-related side effects for amlodipine**

Adverse Event	Placebo N=520	2.5 mg N=275	5.0 mg N=296	10.0 mg N=268
Edema	0.6	1.8	3.0	10.8
Dizziness	1.5	1.1	3.4	3.4
Flushing	0.0	0.7	1.4	2.5
Palpitation	0.5	0.7	1.4	4.5

For several adverse experiences that appear to be drug- and dose-related, there was a greater incidence in women than men associated with amlodipine treatment as shown in the following table:



AZOR™ (amlodipine and olmesartan medoxomil)  
Formulary Dossier

**Table 4: Incidence of drug- and dose-related side effects in women and men for amlodipine**

Adverse Event	Placebo		Amlodipine	
	Male=% (N=914)	Female=% (N=336)	Male=% (N=1218)	Female=% (N=512)
Edema	1.4	5.1	5.6	14.5
Flushing	0.3	0.9	1.5	4.5
Palpitation	0.9	0.9	1.4	3.3
Somnolence	0.8	0.3	1.3	1.6

**Olmesartan medoxomil**

Olmesartan medoxomil has been evaluated for safety in more than 3825 patients/subjects, including more than 3275 patients treated for hypertension in controlled trials. This experience included about 900 patients treated for at least 6 months and more than 525 for at least 1 year. Treatment with olmesartan medoxomil was well tolerated, with an incidence of adverse events similar to that seen with placebo. Events were generally mild, transient, and without relationship to the dose of olmesartan medoxomil.

The overall frequency of adverse events was not dose-related. Analysis of gender, age, and race groups demonstrated no differences between olmesartan medoxomil- and placebo-treated patients. The rate of withdrawals due to adverse events in all trials of hypertensive patients was 2.4% (i.e., 79/3278) of patients treated with olmesartan medoxomil and 2.7% (i.e., 32/1179) of control patients. In placebo-controlled trials, the only adverse event that occurred in more than 1% of patients treated with olmesartan medoxomil and at a higher incidence in olmesartan medoxomil treated patients vs. placebo was dizziness (3% vs 1%).

**j. Drug Interactions:**

**Drug Interactions with AZOR™**

The pharmacokinetics of amlodipine and olmesartan medoxomil are not altered when the drugs are co-administered.

No drug interaction studies have been conducted with AZOR™ and other drugs, although studies have been conducted with the individual amlodipine and olmesartan medoxomil components of AZOR™, as described below, and no significant drug interactions have been observed.

**Drug Interactions with Amlodipine**

In vitro data indicate that amlodipine has no effect on the human plasma protein binding of digoxin, phenytoin, warfarin, and indomethacin.

**Effect of Other Agents on Amlodipine**

*Cimetidine:* Co-administration of amlodipine with cimetidine did not alter the pharmacokinetics of amlodipine.

*Grapefruit juice:* Co-administration of 240 mL of grapefruit juice with a single oral dose of amlodipine 10 mg in 20 healthy volunteers had no significant effect on the pharmacokinetics of amlodipine.

*Maalox® (antacid):* Co-administration of the antacid Maalox® with a single dose of amlodipine had no significant effect on the pharmacokinetics of amlodipine.

*Sildenafil:* A single 100 mg dose of sildenafil in subjects with essential hypertension had no effect on the pharmacokinetic parameters of amlodipine. When amlodipine and sildenafil were used in combination, each agent independently exerted its own blood pressure lowering effect.

**Effect of Amlodipine on Other Agents**

*Atorvastatin:* Co-administration of multiple 10 mg doses of amlodipine with 80 mg of atorvastatin resulted in no significant change in the steady state pharmacokinetic parameters of atorvastatin.

*Digoxin:* Co-administration of amlodipine with digoxin did not change serum digoxin levels or digoxin renal clearance in normal volunteers.

*Ethanol (alcohol):* Single and multiple 10 mg doses of amlodipine had no significant effect on the pharmacokinetics of ethanol.

*Warfarin:* Co-administration of amlodipine with warfarin did not change the warfarin prothrombin response time.

In clinical trials, amlodipine has been safely administered with thiazide diuretics, beta-blockers, angiotensin-converting enzyme inhibitors, long-acting nitrates, sublingual nitroglycerin, digoxin, warfarin, non-steroidal anti-inflammatory drugs, antibiotics, and oral hypoglycemic drugs.

AZOR™ (amlodipine and olmesartan medoxomil)  
Formulary Dossier

**Drug Interactions with Olmesartan Medoxomil**

No significant drug interactions were reported in studies in which olmesartan medoxomil was co-administered with digoxin or warfarin in healthy volunteers.

The bioavailability of olmesartan medoxomil was not significantly altered by the co-administration of antacids [Al(OH)3/Mg(OH)2].

Olmesartan medoxomil is not metabolized by the cytochrome P450 system and has no effects on P450 enzymes; thus, interactions with drugs that inhibit, induce, or are metabolized by those enzymes are not expected.

**k. Dosage and Administration**

AZOR™ may be substituted for its individually titrated components for patients on amlodipine and olmesartan medoxomil. AZOR™ may also be given with increased amounts of amlodipine, olmesartan medoxomil, or both, as needed.

AZOR™ may be used as add-on therapy for patients not adequately controlled on amlodipine or olmesartan medoxomil. Dosage may be increased after 2 weeks to a maximum dose of 10/40 mg once daily, usually by increasing one component at a time but both components can be raised to achieve more rapid control. Maximum antihypertensive effects are attained within 2 weeks after a change in dose. AZOR™ may be administered with other antihypertensive agents.

The side effects of olmesartan medoxomil are generally rare and apparently independent of dose. Those of amlodipine are generally dose-dependent (mostly edema).

AZOR™ may be taken with or without food.

**l. Comparative Pharmacokinetic/Pharmacologic Profiles of Available ARBs**

**Table 5: Comparative Pharmacokinetic/Pharmacologic Profiles of Available ARBs**

	Olmesartan medoxomil	Candesartan cilexetil <sup>1</sup>	Eprosartan <sup>2</sup>	Irbesartan <sup>3</sup>	Losartan potassium <sup>4</sup>	Telmisartan <sup>5</sup>	Valsartan <sup>6</sup>
<b>Prodrug</b>	Yes	Yes	No	No	Yes	No	No
<b>AT<sub>1</sub> Receptor Affinity</b>	IC <sub>50</sub> 5-6 nmol/L	K <sub>i</sub> 0.6	IC <sub>50</sub> 1-4 nmol/L	IC <sub>50</sub> 1.3 nmol/L	IC <sub>50</sub> 20 nmol/L	K <sub>i</sub> 3.7	IC <sub>50</sub> 2.7 nmol/L
<b>Maximal Onset (wk)</b>	2	2-4	3	2	2-3	3	2
<b>Peak (hr)</b>	1-2	3-4	3	1.5-2	3-4	.5-1	2-4
<b>BA (%)</b>		15	13	60-80	33	42-58	25
<b>Food Effect (↓AUC %)</b>	—	—	25	—	10	6	40-50
<b>T<sub>1/2</sub> (hr)</b>	13	9	5-9	12-20	2	24	6
<b>Protein Binding (%)</b>	99	>99	98	90	99	>99.5	>95
<b>P 450 Metabolism</b>	No	No	No	Yes	Yes	No	Uncertain
<b>Drug Interactions</b>	No	No	No	No	Rifampin, fluconazole	Digoxin	No
<b>Fecal Elimination (%)</b>	50-65	67	90	80	60	98	83
<b>Urinary Elimination (%)</b>	35-50	33	7	20	35	<1	13
<b>Trough to Peak Ratio</b>	52-79	80	67	>60	58-78	>97	69-76
<b>Dosages Available (mg)</b>	5, 20, 40	4,8,16,32	400,600	75, 150, 300	25, 50, 100	20, 40, 80	40, 80, 160, 320
<b>Dosing Frequency</b>	QD	QD/BID	QD/BID	QD	QD/BID	QD	QD

BA=bioavailability

AZOR™ (amlodipine and olmesartan medoxomil)  
Formulary Dossier

$K_i$ =inhibition constant

<sup>1</sup>Atacand® package insert. Wayne, PA: AstraZeneca Pharmaceuticals; 2005.

<sup>2</sup>Teveten® package insert. Bridgewater, NJ: Biovail Pharmaceuticals, Inc.; 2004.

<sup>3</sup>Avapro® package insert. Princeton, NJ: Bristol-Myers Squibb Company; 2004.

<sup>4</sup>Cozaar® package insert. West Point, PA: Merck & Co.; 2004.

<sup>5</sup>Micardis® package insert. Ridgefield, CT: Boehringer Ingelheim Pharmaceuticals; 2003.

<sup>6</sup>Diovan® package insert. East Hanover, NJ: Novartis Pharmaceuticals Corp.; 2005.

AZOR™ (amlodipine and olmesartan medoxomil)  
Formulary Dossier

n. Comparative Pharmacokinetic/Pharmacologic Profiles of Available Dihydropyridine CCBs

Table 6: Comparative Pharmacokinetic/Pharmacologic Profiles of Available Dihydropyridine CCBs

	Amlodipine besylate <sup>1</sup>	Felodipine <sup>2</sup>	Isradipine <sup>3</sup>	Nicardipine hydrochloride <sup>4</sup>	Nifedipine <sup>5</sup>	Nisoldipine <sup>6</sup>
Peak (hr)	6-12	2.5-5	8-10	1-4	6	6-12
BA (%)	64-90	20	15-24	35	86	5
Food Effect (↓AUC%)	_____	_____	25	25	_____	25
T <sub>1/2</sub> (hr)	30-50	11-16	8	8.6	2	7-12
Protein Binding (%)	93	>99	95	>95	92-98	>99
P450 Metabolism	No	Yes - CYP3A4	No	No	Yes	Yes – CYP3A4
Drug Interactions	No significant effects by: cimetidine, grapefruit juice, Maalox (antacid), sildenafil, atorvastatin, digoxin, ethanol (alcohol), warfarin, thiazide diuretics, beta-blockers, ACEIs, long-acting nitrates, sublingual nitroglycerin, NSAIDs, antibiotics, and oral hypoglycemic agents	Increased effect with: CYP3A4 inhibitors (eg, itraconazole, erythromycin, grapefruit juice, cimetidine)  <u>Increases effect of:</u> tacrolimus  <u>Decreased effect:</u> anticonvulsants  <u>No significant effects by:</u> Beta-blockers, digoxin, indomethacin, spironolactone	<u>Noted interaction:</u> Fentanyl anesthesia with concomitant use of a beta-blocker may cause severe hypotension  <u>Increases effect of:</u> propranolol  <u>No significant effects by:</u> nitroglycerin, HCTZ, digoxin	<u>Noted interaction:</u> Fentanyl anesthesia with concomitant use of a beta-blocker may cause severe hypotension  <u>Increased effect with:</u> cimetidine  <u>Increases effect of:</u> cyclosporine  <u>No significant effects by:</u> beta-blockers, digoxin, furosemide, propranolol, dipyridamole, warfarin, quinidine, naproxen	Increased effect <u>with:</u> cimetidine  <u>Increases effect of:</u> digitalis  <u>No significant effects by:</u> beta-blockers, long-acting nitrates, coumarin anticoagulants	Increased <u>effect with:</u> cimetidine  <u>Decreased effect with:</u> CYP3A4 inducers, phenytoin, quinidine  <u>No significant effects by:</u> ranitidine, beta-blockers (eg, atenolol, propranolol), digoxin, warfarin
Incidence of Edema (%)	<u>Dose-Related:</u> * 2.5 mg: 1.8 5.0 mg: 3 10 mg: 10.8 Placebo: 0.6	<u>Dose-Related:</u> *† 2.5 mg: 2.0 5.0 mg: 8.8 10 mg: 17.4 Placebo: 3.3	<u>Dose-Related:</u> 5 mg: 8.9 10 mg: 12.7 15 mg: 15.9 20 mg: 35.9 Placebo: 3.6	<u>Dose-Related</u> *† Cardene® SR(exact doses not listed): 5.9 Placebo: 1.4	<u>Dose-Related</u> *† Low Dose: 10 Highest Dose (180 mg): † 30	<u>Dose-Related:</u> *† 10 mg: 7 20 mg: 15 30 mg: 20 40 mg: 27 60 mg: 29 Placebo: 10
Fecal Elimination (%)	Not listed	10	25-30	35	20-40	Not listed
Urinary Elimination (%)	10% parent compound; 60% metabolites	70	60-65	60	60-80	60-80
Trough:Peak Ratio	Not listed	40-50	76	Not listed	Not listed	70-100
Dosages Available (mg)	2.5, 5, 10	2.5, 5, 10	5, 10	30, 45, 60	30, 60, 90	10, 20, 30, 40
Dosing Frequency	Daily	Daily	Daily	Twice Daily	Daily	Daily

BA = bioavailability

\* Results are from pooled data

AZOR™ (amlodipine and olmesartan medoxomil)  
Formulary Dossier

† Results presented as therapy without regard to causality

<sup>1</sup> Norvasc® package insert. New York, NY: Pfizer Inc.,; 2005

<sup>2</sup> Plendil® package insert. Wilmington, DE: AstraZeneca LP; 2003

<sup>3</sup> DynaCirc CR® package insert. Liberty Corner, NJ: Reliant Pharmaceuticals, Inc.; 2005

<sup>4</sup> Cardene® SR package insert. Nutley, NJ: Roche Pharmaceuticals; 2000

<sup>5</sup> Procardia XL® package insert. New York, NY: Pfizer Inc.; 2003

<sup>6</sup> Sular® package insert. Atlanta, GA: Sciele Pharma, Inc.; 2006

AZOR™ (amlodipine and olmesartan medoxomil)  
Formulary Dossier

- o. Comparative Pharmacokinetic/Pharmacologic Profiles of Fixed Dose Amlodipine Combination Products

**Table 7: Comparative Pharmacokinetic/Pharmacologic Profiles of Fixed Dose Amlodipine Combination Products**

	<b>AZOR™ (amlodipine and olmesartan medoxomil)<sup>1</sup></b>	<b>Exforge® (amlodipine and valsartan)<sup>2</sup></b>	<b>Lotrel® (amlodipine besylate/benazepril HCl)<sup>3</sup></b>
<b>Indications</b>	Hypertension, alone or with other antihypertensive agents  Not indicated for initial therapy	Hypertension  Not indicated for initial therapy	Hypertension  Not indicated for initial therapy
<b>BA (%)</b>	<b>Amlodipine:</b> 64-90 <b>Olmesartan:</b> 26	<b>Amlodipine:</b> 64-90 <b>Valsartan:</b> 25% (range: 10-35)	<b>Amlodipine:</b> 64-90 <b>Benazepril:</b> ≥37
<b>T<sub>1/2</sub> (hr)</b>	<b>Amlodipine:</b> 30-50 <b>Olmesartan:</b> 13 (elimination)	<b>Amlodipine:</b> 30-50 <b>Valsartan:</b> 6 (elimination)	<b>Amlodipine:</b> approximately 2 days (elimination) <b>Benazepril:</b> 10-11 (elimination)
<b>Time to Peak Plasma Concentrations (T<sub>max</sub>) (hr)</b>	<b>Amlodipine:</b> 6-12 <b>Olmesartan:</b> 1-2	<b>Amlodipine:</b> 6-12 <b>Valsartan:</b> 2-4	<b>Amlodipine:</b> 6-12 <b>Benazepril:</b> 1.5-4
<b>Protein Binding (%)</b>	<b>Amlodipine:</b> 93 <b>Olmesartan:</b> 99	<b>Amlodipine:</b> 93 <b>Valsartan:</b> 95	<b>Amlodipine:</b> 93 <b>Benazepril:</b> Not specified
<b>Volume of Distribution (L)</b>	<b>Amlodipine:</b> 21 <b>Olmesartan:</b> 17	<b>Amlodipine:</b> 21 <b>Valsartan:</b> 17	<b>Amlodipine:</b> 21 <b>Benazeprilat:</b> 0.7
<b>Metabolism</b>	<b>Amlodipine:</b> extensively (about 90%) converted to inactive metabolites via hepatic metabolism  <b>Olmesartan:</b> following the rapid and complete conversion of olmesartan medoxomil to olmesartan during absorption, there is virtually no further metabolism of olmesartan	No CYP450 Metabolism  <b>Amlodipine:</b> extensively (about 90%) converted to inactive metabolites via hepatic metabolism <b>Valsartan:</b> primary metabolite is valeryl 4-hydroxy valsartan and accounts for 9% of dose	<b>Amlodipine:</b> extensively metabolized in the liver <b>Benazepril:</b> the metabolism of benazepril to benazeprilat is almost complete
<b>Elimination</b>	<b>Amlodipine:</b> 10% of the parent compound and 60% of the metabolites are excreted in the urine  <b>Olmesartan:</b> Approximately 35% to 50% of the absorbed dose is recovered in urine while the remainder is eliminated in feces via the bile	<b>Amlodipine:</b> 10% of parent compound and 60% of the metabolites excreted in the urine <b>Valsartan:</b> when administered as an oral solution, is primarily recovered in feces (about 83% of dose) and urine (about 13% of dose)	<b>Amlodipine:</b> 10% of the parent compound and 60% of the metabolites excreted in the urine <b>Benazepril:</b> only trace amounts of an administered dose of benazepril can be recovered unchanged in the urine; about 20% of the dose is excreted as benazeprilat, 8% as benazeprilat glucuronide, and 4% as benazepril glucuronide

AZOR™ (amlodipine and olmesartan medoxomil)  
Formulary Dossier

Continued – Table 7 - Comparative Pharmacokinetic/Pharmacologic Profiles of Fixed Dose Amlodipine Combination Products

	AZOR™ (amlodipine and olmesartan medoxomil) <sup>1</sup>	Exforge® (amlodipine and valsartan) <sup>2</sup>	Lotrel® (amlodipine besylate/benazepril HCl) <sup>3</sup>
<b>Drug Interactions</b>	<p>No drug interaction studies have been conducted with AZOR and other drugs, although studies have been conducted with the individual amlodipine and olmesartan components, as described below:</p> <p><b>Amlodipine:</b> <u>No significant effects with/by:</u> cimetidine, grapefruit juice, sildenafil, atorvastatin, digoxin, ethanol (alcohol), warfarin In clinical trials, amlodipine has been safely administered with thiazide diuretics, beta blockers, angiotensin-converting enzyme inhibitors, long-acting nitrates, sublingual nitroglycerin, digoxin, warfarin, non-steroidal anti-inflammatory drugs, antibiotics, and oral hypoglycemic drugs</p> <p><b>Olmesartan:</b> No significant drug interactions were reported in studies in which olmesartan was coadministered with digoxin or warfarin in healthy volunteers. Olmesartan medoxomil is not metabolized by the cytochrome P450 system; thus, interactions with drugs that inhibit, induce, or are metabolized by those enzymes are not expected</p>	<p>No drug interaction studies have been conducted with Exforge and other drugs, although studies have been conducted with the individual amlodipine and valsartan components, as described below:</p> <p><b>Amlodipine:</b> <u>No significant effects with/by:</u> cimetidine, grapefruit juice, Maalox (antacid), sildenafil, atorvastatin, digoxin, ethanol (alcohol), warfarin, thiazide diuretics, beta-blockers, ACEIs, long-acting nitrates, sublingual nitroglycerin, NSAIDs, antibiotics, and oral hypoglycemic agents.</p> <p><b>Valsartan:</b> <u>No significant effects with/by:</u> amlodipine, atenolol, cimetidine, digoxin, furosemide, glyburide, hydrochlorothiazide, indomethacin, or warfarin. Concomitant use of potassium sparing diuretics (e.g., spirinolactone, triamterene, amiloride), potassium supplements, or salt substitutes containing potassium may lead to increases in serum potassium and serum creatinine in heart failure patients, as with other drugs that block angiotensin II or its effects.</p>	<p>Patients receiving concomitant diuretic therapy may occasionally experience an excessive reduction in blood pressure after initiation of therapy with Lotrel.</p> <p><b>Amlodipine:</b> safely administered with thiazide diuretics, beta blockers, ACE inhibitors, long-acting nitrates, sublingual nitroglycerin, digoxin, warfarin, nonsteroidal anti-inflammatory drugs, antibiotics, and oral hypoglycemic drugs.</p> <p><b>Benazepril:</b> safe to administer and use concomitantly with oral anticoagulants, beta-adrenergic blocking agents, calcium-blocking agents, cimetidine, diuretics, digoxin, hydralazine, and naproxen. Increased serum lithium levels and symptoms of lithium toxicity have been reported – coadminister with caution and monitor serum lithium frequently. Benazepril may attenuate potassium loss caused by thiazide diuretics. Potassium-sparing diuretics (spirinolactone, amiloride, triamterene, and others) or potassium supplements can increase the risk of hyperkalemia – coadminister with caution and monitor serum potassium frequently</p>

AZOR™ (amlodipine and olmesartan medoxomil)  
Formulary Dossier

Continued – Table 7: Comparative Pharmacokinetic/Pharmacologic Profiles of Fixed Dose Amlodipine Combination Products

	AZOR™ (amlodipine and olmesartan medoxomil) <sup>1</sup>	Exforge® (amlodipine and valsartan) <sup>2</sup>	Lotrel® (amlodipine besylate/benazepril HCl) <sup>3</sup>																									
Adverse Reactions	<p>Edema is a known, dose-dependent adverse effect of amlodipine but not of olmesartan medoxomil.</p> <p>The placebo-subtracted incidence of edema during the 8-week, randomized, double-blind treatment period was highest with amlodipine 10 mg monotherapy. The incidence was significantly reduced when 20 mg or 40 mg of olmesartan medoxomil was added to the 10 mg amlodipine dose.</p> <table border="1"><tr><td colspan="2"></td><td colspan="3">Olmesartan Medoxomil</td></tr><tr><td colspan="2"></td><td>Placebo</td><td>20 mg</td><td>40 mg</td></tr><tr><td>Amlodipine</td><td>Placebo</td><td>0%*</td><td>(-2.4%)</td><td>6.2%</td></tr><tr><td></td><td>5 mg</td><td>0.7%</td><td>5.7%</td><td>6.2%</td></tr><tr><td></td><td>10 mg</td><td>24.5%</td><td>13.3%</td><td>11.2%</td></tr></table> <p>*12.3% = actual placebo incidence</p> <p>Adverse reactions seen at lower rates during the double-blind period also occurred in the patients treated with AZOR™ at about the same or greater incidence as in patients receiving placebo.</p> <p>These included hypotension, orthostatic hypotension, rash, pruritus, palpitation, urinary frequency, and nocturia.</p>			Olmesartan Medoxomil					Placebo	20 mg	40 mg	Amlodipine	Placebo	0%*	(-2.4%)	6.2%		5 mg	0.7%	5.7%	6.2%		10 mg	24.5%	13.3%	11.2%	<p><b>Incidence of AEs in ≥2% patients and more frequent than in placebo patients:</b></p> <p>Peripheral edema (5.4%) Nasopharyngitis (4.3%) Upper respiratory tract infection (2.9%) Dizziness (2.1%)</p>	<p><b>Incidence of AEs in ≥1% of patients considered possibly or probably related to study drug and more frequent than in placebo patients:</b></p> <p>Cough (3.3%) Headache (2.2%) Edema (2.1%) Dizziness (1.3%)</p>
		Olmesartan Medoxomil																										
		Placebo	20 mg	40 mg																								
Amlodipine	Placebo	0%*	(-2.4%)	6.2%																								
	5 mg	0.7%	5.7%	6.2%																								
	10 mg	24.5%	13.3%	11.2%																								
Dosages Available (mg)	5/20, 5/40, 10/20, 10/40 combination tablets	5/160, 10/160, 5/320, and 10/320 combination tablets	2.5/10, 5/10, 5/20, 5/40, 10/20, 10/40 combination tablets																									
Adult Dosing Frequency	Hypertension: Range: 5/20 mg daily up to a maximum dose of 10/40 mg daily	Hypertension: Range: 5/160 mg daily up to a maximum dose of 10/320 mg daily  For convenience, patients receiving amlodipine and valsartan from separate tablets may instead wish to receive tablets of Exforge® containing the same component doses	Hypertension: Range: 2.5/10 mg daily up to a maximum dose of 10/40 mg daily  For convenience, patients receiving amlodipine and benazepril from separate tablets may instead wish to receive tablets of Lotrel® containing the same component doses																									

BA = bioavailability

<sup>1</sup> AZOR package insert. Parsippany, NJ: Daiichi Sankyo, Inc.; 2007

<sup>2</sup> Exforge package insert. East Hanover, NJ: Novartis Pharmaceuticals; 2007

<sup>3</sup> Lotrel package insert. East Hanover, NJ: Novartis Pharmaceuticals; 2007



## 1.2 PLACE OF THE PRODUCT IN THERAPY

### 1.2.1 DISEASE DESCRIPTION: HYPERTENSION

#### a. *Epidemiology and Clinical Aspects of Hypertension*

Hypertension is a major cardiovascular risk factor that directly contributes to myocardial infarction (MI), cerebrovascular accidents, congestive heart failure (CHF), peripheral arterial insufficiency, and premature mortality. Optimal and cost-effective management of the condition depends on careful diagnosis, treatment minimization, optimized adherence, cost effective selection of tests and treatments, and practice efficiency.

Cardiovascular disease has been the dominant cause of death in the United States for over 50 years. About 12 million Americans currently have coronary heart disease. Another 4 million have had a stroke (Cooper, 2000). Heart disease and stroke are the first- and third-leading causes of death in the US, respectively. They also place a staggering financial burden on the US healthcare system, totaling over \$259 billion in direct and indirect costs (JNC-VI, 1997).

The incidence of morbidity and mortality from cardiovascular disease increases with age. The average annual rate of first major cardiovascular events increases nearly tenfold from 7 per 1000 at ages 35 to 44 years to 68 per 1000 at ages 85 to 94 years in men, with similar rates observed in women 10 years later in life. One quarter of nursing home residents aged 65 years and older have a primary diagnosis of cardiovascular disease at admission. Furthermore, over 80% of the nearly 950,000 annual deaths from cardiovascular disease occur in people aged 65 years old or older.

The overall estimated prevalence of hypertension was 72 million in 2004 (AHA, Heart Disease and Stroke Statistics, 2007). The relationship between BP and risk of CVD events is continuous, consistent, and independent of other risk factors. The higher the BP, the greater is the chance of heart attack, heart failure, stroke, and kidney disease. For individuals 40–70 years of age, each increment of 20 mmHg in systolic BP (SBP) or 10 mmHg in diastolic BP (DBP) doubles the risk of CVD across the entire BP range from 115/75 to 185/115 mmHg (JNC-7, 2003).

The vast majority of patients with hypertension (93% to 95%) display no demonstrable, curable abnormality of anatomy or physiology (Rudd, 1996). Termed *primary hypertension*, the condition carries no consistent hallmark symptoms or signs, except for the elevated BP itself. Most cases are detected incidentally as part of routine examinations and generally in the absence of target organ damage at initial presentation. Reports of headache, dizziness, fatigue, palpitations, and chest discomfort occur commonly among both patients with hypertension and patients without hypertension. Because the symptoms prompt clinician attention, hypertension is more likely to be detected among patients with symptoms. For the majority of patients with hypertension, symptoms and symptom levels do not correlate well with BP level (Pickering, 2000).

In clinical trials, antihypertensive therapy has been associated with reductions in stroke incidence averaging 35–40%; myocardial infarction, 20–25%; and more than 50% for heart failure. It is estimated that in patients with stage 1 hypertension (SBP 140–159 mmHg and/or DBP 90–99 mmHg) and additional cardiovascular risk factors, achieving a sustained 12 mmHg reduction in SBP over 10 years will prevent 1 death for every 11 patients treated. In the presence of CVD or target organ damage, only 9 patients would require such BP reduction to prevent a death (JNC-7, 2003).

#### b. *The Role of Calcium and the Renin-Angiotensin System (RAS) in Hypertension*

Blood pressure is determined by the balance between cardiac output and peripheral resistance, where elevated peripheral resistance is the primary cause of human hypertension. Peripheral resistance is believed to be caused by prolonged smooth muscle contraction of small arterioles as a result of increased intracellular calcium (Beevers 2001). Normally, cells maintain a low resting intracellular concentration of ionized calcium in the face of a large and inwardly directed concentration gradient. As calcium enters the cell, it combines with calcium-binding proteins to stimulate a number of secondary messenger systems and cellular responses, such as nerve excitation, cardiac and vascular smooth muscle contraction, and hormone secretion. Calcium channels can be differentiated into several subtypes, but the L-type channel is the channel that is most directly associated with control of blood pressure. L-type calcium channels are responsible for normal myocardial and smooth vascular muscle contraction (Stern 1992). Prolonged

## AZOR™ (amlodipine and olmesartan medoxomil) Formulary Dossier

activation of the L-type calcium channel results in an increased intracellular calcium concentration and subsequent smooth muscle contraction (Stern 1992). Irreversible increases in peripheral resistance have been reported with prolonged smooth muscle constriction and are most likely associated with structural alterations of arteriolar vessels, including thickening of vessel walls (Beevers 2001).

The renin-angiotensin system (RAS), which plays a pivotal role both in the control of blood pressure and the pathogenesis of hypertension, is a complex cascade initiated by the conversion of angiotensinogen into angiotensin I by renin which is a selective enzyme, in the liver (Dina, 2000; Burnier, 2000). Locally, angiotensin I is then converted into angiotensin II, a peptide hormone, primarily in a reaction catalyzed by angiotensin-converting enzyme (ACE; Dina, 2000).

It has been recognized since the 1970's that elevated levels of angiotensin II adversely affect the heart and kidney, and that elevated plasma renin levels increase the risk of myocardial infarction and stroke (Dina, 2000, Burnier, 2000, Burnier, 2001; Yusuf, 2000). Specifically, angiotensin II causes vasoconstriction, proximal tubular sodium reabsorption, inotropism, chronotropism, and alpha-adrenergic receptor stimulation (Dina, 2000). In early animal studies, intravenous administration of angiotensin II was shown to result in renal failure and tubular necrosis (Gavras, 1971). Likewise, early studies in humans found that patients with elevated plasma renin levels had an 11% frequency of heart attacks and a 14% frequency of strokes compared to 0% over the same time period in patients with low serum renin levels (Brunner, 1972).

Research into intervention with the RAS has led to important developments in antihypertensive therapy. Captopril®, introduced in 1981, was the first oral ACE inhibitor (Burnier, 2000). ACE inhibitors, which block the production of angiotensin II, remain an important therapeutic option for patients with hypertension (Burnier, 2000, 2001).

Numerous clinical studies have demonstrated the efficacy of various ACE inhibitors in patients with hypertension, cardiovascular disorders, and renal disorders (Dina, 2000, Burnier, 2001; Yusuf, 2000; CONSENSUS, 1987; SOLVD Investigators, 1991, 1992; Lewis, 1993; Heart Outcomes Prevention Evaluation Study Investigators, 2000). For example, in 1987, the Cooperative North Scandinavian Enalapril Survival Study (CONSENSUS) concluded that ACE inhibition reduced cardiovascular mortality by 31% after one year and improved symptoms in 127 patients with severe congestive heart failure (CONSENSUS, 1987). In 1991, the Studies of Left Ventricular Dysfunction (SOLVD) reported similar findings (SOLVD Investigators, 1991). The addition of an ACE inhibitor to conventional therapy in patients with reduced left ventricular ejection fractions and congestive heart failure resulted in a 16% reduction in the risk of death compared with placebo. These investigators also found a significant reduction in the rate of heart failure in asymptomatic patients with reduced left ventricular ejection fractions when treated with an ACE inhibitor (SOLVD Investigators, 1992).

In the 1990s, researchers discovered that ACE inhibition also provides significant protection against deterioration of renal function in patients with type 1 diabetes, independent of blood pressure control. ACE-inhibitor therapy was associated with a 50% reduction in the combined endpoints of death, dialysis, and transplantation (Lewis, 1993).

More recently, findings of the Heart Outcomes Prevention Evaluation (HOPE) study led researchers to speculate that the clinical benefit of anti-RAS agents may go beyond reducing blood pressure. The HOPE study demonstrated a significant reduction in the rates of death, myocardial infarction, stroke, other cardiovascular events, and the need for revascularization procedures in high-risk patients following treatment with an ACE inhibitor (ramipril). These benefits, including a 25% reduction in cardiovascular-related mortality, were achieved with only a 2 to 3 mm Hg reduction in blood pressure in patients who were normotensive (139/79 mm Hg) at the time of entry into the study (Yusuf, 2000).

### 1.2.2 APPROACHES TO TREATMENT

The National High Blood Pressure Education Program (NHBPEP), a division of the National Institutes of Health (NIH), issues national recommendations every four years on the prevention, detection, evaluation, and treatment of high blood pressure. The latest version, JNC-7, was published in JAMA in May 2003 by the NIH with important new changes.

Some of the more important revisions from the previous version are:

- A new category designated prehypertension (combined high normal and normal BP from JNC-VI)
- Stages 2 and 3 hypertension from JNC-VI have been combined (now stage 2 in JNC-7)

AZOR™ (amlodipine and olmesartan medoxomil)  
Formulary Dossier

- The risk groups based on risk factors and target organ damage have been eliminated
- Thiazide diuretics are recommended initially alone or in combination with an ACE inhibitor, angiotensin receptor blocker, beta blocker, or calcium channel blocker
- The antihypertensive goal for patients with diabetes has been reduced from 130/85 mm Hg to 130/80 mmHg (it is now the same as the ADA Guidelines)
- The risk of CVD begins at a BP of 115/75 mm Hg (even lower than the JNC VI optimal BP of 120/80 mm Hg)
- Most patients will need two or more antihypertensive medications to achieve their BP goal

Americans who are at high risk of cardiovascular disease and associated morbidity and mortality are often not identified, and as a result, do not receive sufficient treatment. A recent report from the National Health and Nutrition Examination Survey (NHANES) conducted in 1999-2000 concluded that 70% were aware of their hypertension, 59% were treated, and 34% had their hypertension controlled. When clinicians fail to prescribe lifestyle modifications, adequate antihypertensive drug doses, or appropriate drug combinations, inadequate BP control may result.

**a. JNC-7 Treatment Guidelines**

The JNC-7 treatment guidelines classify patients based on blood pressure levels only (Table 8). The 1997 JNC-VI guidelines included risk factor groups in the treatment guidelines. Since then, the new JNC 7 treatment guidelines have been simplified. The new JNC-7 guidelines eliminated the risk factor groups, condensed the 2<sup>nd</sup> and 3<sup>rd</sup> stages of hypertension into one stage, and also added a normal classification for hypertension. The NIH recognized that more stringent BP control is required to achieve optimal benefit in high-risk conditions. Treatment of hypertension to these lower levels may provide benefits in preventing stroke, preserving renal function, and slowing the progression of heart failure.

The prehypertension classification (classified as SBP 120-139 mm Hg or DBP 80-89 mm Hg) introduced in JNC 7 is based on the recognition that the virtually linear increase in cardiovascular event risk with increasing blood pressure begins at levels lower than previously recognized (Table 8). It is estimated that about 22% (approximately 46 million) of the adult population falls into the prehypertension category. The intent of the NIH in creating this new designation was to stress lifestyle modification to patients and the need for increased education of healthcare professionals and the public to reduce blood pressure levels and prevent development of hypertension in the general population.

**Goal:** Treatment goals in JNC-7 remain unchanged from JNC-VI for patients with uncomplicated hypertension in that the NIH recommends that blood pressure should be controlled to below 140/90 mm Hg. In patients who have co-morbidities such as diabetes or chronic kidney disease, the NIH now recommends that BP be controlled to < 130/80 mm Hg (< 130/85 mm Hg in the JNC-VI guidelines).

**Treatment:** All patients with hypertension should begin with lifestyle modifications. In addition, those patients classified as having prehypertension with a compelling indication, may be managed with lifestyle modifications and drug therapy as well. However, it has always been recognized that once blood pressure rises above the widely recognized threshold of 140/90 mm Hg, it will be necessary to introduce a program of medical treatment. JNC-7 has recognized the *ARB class* for their positive results in recent outcome trials. With the introduction of the ARBs in 1995, JNC VI had not considered this class for any indications due to lack of evidence. However, JNC-7 has recommended ARBs for the following compelling indications: *heart failure, diabetes and chronic kidney disease*. JNC-7 recommends starting with a thiazide diuretic either alone or in combination with one of the other classes of antihypertensives (ACEIs, ARBs, BBs, CCBs) unless there are compelling indications to use other agents. It is recommended that practitioners begin with a low dose of the drug and titrate upward if necessary. Most patients who are hypertensive will require two or more antihypertensive medications to achieve their BP goals. Addition of a second drug from a different class should be initiated when use of a single drug in adequate doses fails to achieve the BP goal. When BP is more than 20/10 mmHg above goal, consideration should be given to initiating therapy with two drugs, either as separate prescriptions or in fixed-dose combinations. The initiation of drug therapy with more than 1 agent may increase the likelihood of achieving the BP goal in a more timely fashion, but particular caution is advised in those at risk for orthostatic hypotension, such as patients with diabetes, autonomic dysfunction, and some older persons. It should be noted that AZOR™ (amlodipine besylate/olmesartan medoxomil) is indicated for the treatment of hypertension, alone or with other antihypertensive agents. This fixed combination drug is not indicated for the initial therapy of hypertension

AZOR™ (amlodipine and olmesartan medoxomil)  
Formulary Dossier

**Table 8: JNC-7 Treatment Guidelines: Classification and Management of Blood Pressure for Adults Aged 18 Years or Older**

BP Classification (SBP/DBP mm Hg)	Lifestyle Modification	Initial Drug Therapy Without Compelling Indication	Initial Drug Therapy With Compelling Indications
<b>Normal</b> ( $< 120$ and $< 80$ )	Encourage		
<b>Prehypertension</b> (120-139 or 80-89)	Yes	No antihypertensive drug indicated	Drug(s) for the compelling indications*
<b>Stage 1 Hypertension</b> (140-159 or 90-99)	Yes	Thiazide-type diuretics for most; may consider ACEI, ARB, BB, CCB or combination	Drug(s) for compelling indications. Other antihypertensive drugs (diuretics, ACEI, ARB, BB, CCB) as needed
<b>Stage 2 Hypertension</b> ( $\geq 160$ or $\geq 100$ )	Yes	Two-drug combo for most (usually thiazide-type diuretic and ACEI or ARB or BB or CCB)**	

ACEI: angiotensin converting enzyme inhibitor ARB: angiotensin receptor blocker

BB: beta-blocker CCB: calcium channel blocker

\*Treat patients with chronic kidney disease or diabetes to BP goal of less than 130/80 mm Hg

\*\* Initial combined therapy should be used cautiously in those at risk for orthostatic hypotension

**b. Angiotensin Receptor Blockers (ARBs) in Antihypertensive Therapy**

Angiotensin I is converted to angiotensin II via at least two pathways. Best known is the ACE-dependent metabolic pathway, which is generally considered the primary source of systemic angiotensin II. This is the pathway that is blocked by ACE inhibitors. Among the most actively studied alternative metabolic pathways is that mediated by chymase, which is derived from mast cells (Arakawa, 1980). Evidence suggests that chymase-dependent angiotensin II production may be important for some forms of cardiovascular pathology (Uehara, 2000; Warnholtz, 1999; Ihara, 1999).

Regardless of the metabolic pathway involved, angiotensin II activity is mediated by two classes of receptors, known as AT<sub>1</sub> and AT<sub>2</sub>. Stimulation of AT<sub>1</sub> receptors is responsible for all the known clinical and cardiovascular pathological effects of angiotensin II. Selective blockade of angiotensin II at AT<sub>1</sub> receptors by angiotensin II receptor blockers results in the antihypertensive and other pharmacologic effects of these drugs. The role of AT<sub>2</sub> receptors is less well understood, although stimulation of these receptors may help counterbalance some of the effects of angiotensin II mediated by AT<sub>1</sub> receptors (Burnier, 2000).

AT<sub>1</sub> receptor blockade has advantages over ACE inhibition, since ARBs block the effects of angiotensin II produced by both ACE-dependent and non-ACE-dependent pathways. In addition, ACE inhibitors block the breakdown of plasma bradykinin, which is thought to result in cough and angioedema, the most common side effects of these drugs (Burnier, 2000). These effects do not occur as a result of AT<sub>1</sub> receptor blockade and may explain why ARBs provide an equally therapeutic and better-tolerated alternative to ACE inhibitors for many patients with hypertension.

**c. Calcium Channel Blockers (CCBs) in Antihypertensive Therapy**

Calcium channel blockers (or calcium antagonists) represent an important group of agents for the treatment of hypertension. Calcium antagonists exhibit an antihypertensive effect by relaxing vascular smooth muscle through the inhibition of the influx of calcium through the cell membrane via the L-type calcium channels (Elliott 2001). Inhibition of the influx of calcium decreases contraction in myocardial tissue and arteriolar vascular smooth muscle. The vasodilatory properties of calcium channel blockers work to enhance total coronary blood flow due to their ability to decrease vascular tone and resistance in the coronary circulation (Faulkner 2001), thus decreasing total peripheral resistance without significantly affecting cardiac preload.

## AZOR™ (amlodipine and olmesartan medoxomil) Formulary Dossier

Depending on the agent and its subtype (non-dihydropyridine or dihydropyridine), calcium antagonists differ in vascular selectivity and other cardio-vascular parameters, particularly in regards to conduction and contractility. Certain agents act more on the calcium of cardiac channels than on the vascular ones and thus have a pronounced negative effect on atrioventricular conduction and contractility (phenylalkylamines), others act more on the channels of the myocytes membranes of the resistance vessels and are thus more vasculo-selective (dihydropyridines), and some act proportionately at the level of the heart and of the arterioles (benzothiazepines). Those derived from the phenylalkylamine and benzothiazepine (or non-dihydropyridine) subtype decrease the conductivity, excitability and the contractility of the heart (Hernandez-Hernandez, 2002).

A marked decrease in arterial blood pressure and dilation of resistance vessels elicits the activation of sympathetic reflexes that may result in reflex tachycardia. Reflex tachycardia is particularly evident in the short-acting agents with rapid absorption (1<sup>st</sup> generation dihydropyridines). The newer 2<sup>nd</sup> generation agents, including amlodipine, have characteristically slower starting effects and a longer duration of action, thus making them favorable for the use in long-term antihypertensive treatment with a decreased incidence of reflex tachycardia as compared to their shorter-acting predecessors (Hernandez-Hernandez, 2002). Additionally, because amlodipine's affinity for vascular tissue is much greater than its affinity for the myocardium, lower doses may be used to maintain its antihypertensive effect without reducing the contractility of myocardial tissue; a factor that is beneficial for patients whom negative inotropy might be an issue (Faulkner 2001). Thus, calcium antagonists have shown to play an integral role in the chronic treatment of hypertension, proving efficacious when used as monotherapy as well as in combination.

### ***d. Place of Combination Products in Hypertension Therapy***

Guidelines relating to the treatment of hypertension are consistently updated with new blood pressure goal ranges. Goal ranges have been on the decline for decades, making it increasingly more difficult for physicians to adequately treat patients to goal with single agent therapy (Stergiou 2006). Current evidence suggests that the majority of patients treated for hypertension, roughly two thirds of the diagnosed population, will require combination therapy in order to achieve targeted blood pressure goals (Sica 2002, Stergiou 2006, Tejada 2007). Combination therapy is required when a patient is unable to reach goal with the use of a single agent in combination with diet and exercise modifications. Therapy utilizing multiple anti-hypertensive agents, from diverse yet complimentary classes, provides a mechanism to target the disease through multiple physiologic actions (Faulkner 2001, Sica 2002). It is estimated that a therapeutic success rate of 80-90% is achieved with the use of multiple antihypertensive agents as compared to single-agent therapy. The ALLHAT study reported similar estimates, disclosing that 63% of study patients required combination therapy of two or more drugs to reach blood pressure goals (Stergiou 2006). JNC-7 states that combination therapy utilizing two or more agents is generally required to maintain adequate blood pressure control. JNC-7 guidelines recommend initiation of two drugs in patients presenting with systolic blood pressure >20 mmHg above goal or diastolic blood pressure >10 mmHg above goal. This recommendation is made based upon evidence showing that combination therapy increases the probability of achieving blood pressure goals within a shorter time period as compared to initiating single drug therapy (Chobanian 2004).

Combination therapy has been reported to increase clinical efficacy, offset adverse events observed with monotherapy, and increase compliance (Tejada 2007). Monotherapy often results in the stimulation of compensatory mechanisms, thus diminishing overall efficacy (Faulkner 2001). By combining multiple agents, regulatory mechanisms potentiated by one drug may be counter-balanced by the activity of another drug (Sica 2002). Additionally, single agent therapy regularly requires use of the maximum daily dose, resulting in an increased incidence of adverse events. The increased risk of side effects has been associated with decreased compliance rates and subsequent therapeutic failure (Faulkner 2001). A study comparing the use of a fixed combination of ramapril/felodipine with the component monotherapies, reported a significantly greater responder rate, 71.4% for the combination and 41.2% and 28.6% for the respective monotherapies. This finding was attributed to reduced adverse event prevalence in patients receiving the combination agent (Scholze 2006). A separate study looking at adherence patterns, evaluated patients receiving a fixed dose combination (FDC) of amlodipine and benazepril compared to patients taking an ACE inhibitor and a dihydropyridine calcium channel blocker as separate agents, and found adherence rates of 87.9% and 69.2%, respectively. Greater adherence to the fixed dose combination was due to regimen simplification (Germino 2007). Studies have also shown that FDCs, like standard combination therapies, are generally better tolerated as lower doses of the component monotherapies are required, and improved blood pressure lowering over component monotherapy has been reported (Tejada 2007).

### 1.3 EVIDENCE FOR PHARMACOGENOMIC TESTS AND DRUGS

Not applicable.

## 2. SUPPORTING CLINICAL AND ECONOMIC INFORMATION

### 2.1 SUMMARIZING KEY CLINICAL AND ECONOMIC STUDIES

#### **A Randomized, Double-Blind, Placebo-Controlled Factorial Study Evaluating The Efficacy and Safety of Co-Administration of Amlodipine Besylate Plus Olmesartan Medoxomil Compared to Monotherapy in Patients with Mild to Severe Hypertension (Blinded-Study Phase, Phase III, United States)**

**Objectives:** The main objective of the blinded-phase (day 1 to week 8) of this pivotal trial was to determine if co-administration of amlodipine besylate (AML) and olmesartan medoxomil (OM) would have a clinically significant benefit versus the respective monotherapy components in controlling blood pressure in patients with mild to severe hypertension.

The primary efficacy endpoint was the change from baseline in SeDBP at the end of Period II (Week 8). If a patient withdrew from the study prior to Week 8, the last observed value during the randomized, double-blind treatment period was carried forward (LOCF) for the primary efficacy analysis. Change from baseline in SeSBP at the end of Period II with LOCF was the secondary efficacy variable.

Secondary endpoints also included the number (percentage) of patients achieving blood pressure goal (defined as blood pressure <140/90 mm Hg, <130/80 mm Hg for patients with diabetes). In addition, change from baseline in SeDBP and SeSBP at weeks 2, 4, 6, and 8 without LOCF were evaluated.

#### **Methodology:**

The 8-week, multi-center, randomized, double-blind, placebo-controlled, parallel-group factorial study was designed to assess the efficacy and safety of co-administration of amlodipine (AML) plus olmesartan medoxomil (OM) in adult patients (N=1940 patients; 384 patients ≥65 years of age) with mild to severe hypertension, in comparison to the respective monotherapies. This 8-week study was a part of a 52-week trial which consisted of 3 periods: 1) a washout period of approximately 2 weeks, 2) an 8-week, double-blind treatment period, and 3) a 44-week, open-label treatment period. Period I of this study consisted of a single screening visit for patients not on antihypertensive medications and a two-week washout period for patients on antihypertensive medications. To be eligible for randomization, all patients (males and females ≥18 years of age) had to have a mean SeDBP ≥95 mm Hg and ≤120 mmHg, with ≤10 mmHg difference between the pre-randomization and the randomization visit BP measurements.

Period II (double-blind) consisted of an 8-week treatment period. Patients who met all of the inclusion criteria and none of the exclusion criteria were randomized equally to one of the following 12 treatment arms: placebo, monotherapy treatment with amlodipine 5 mg or 10 mg, monotherapy treatment with olmesartan medoxomil 10 mg, 20 mg, or 40 mg, or combination therapy with amlodipine/olmesartan medoxomil at doses of 5/10 mg, 5/20 mg, 5/40 mg, 10/10 mg, 10/20 mg, and 10/40 mg. Patients were discontinued from the study if SeSBP was <90 mm Hg or SeDBP was >120 mmHg at any visit during Period II. The ITT (Intent-to-Treat) population was the primary analysis population for efficacy evaluations. The ITT population included patients who took at least 1 dose of randomized double-blind study medication, had baseline blood pressure measurements, and at least 1 blood pressure measurement after taking randomized double-blind study medication.

Treatment was double-blind, parallel-arm for all randomized patients. To achieve even distribution among treatment groups within each stratum, the randomization process included stratification factors for age group (≥65 years, <65 years) and diabetic status. This study targeted approximately 20% of the patients to be ≥ 65 years of age.

Period III consisted of a 44-week, open-label treatment period to assess long-term safety and efficacy of combination therapy with amlodipine 5 mg plus olmesartan medoxomil 40 mg. During the open-label extension, patients whose blood pressure was not adequately controlled (i.e., did not achieve a blood pressure goal of <140/90 mm Hg, or <130/80 mmHg for those patients with diabetes) on amlodipine 5 mg plus olmesartan medoxomil 40 mg were titrated to amlodipine 10 mg plus olmesartan 40 mg. Patients whose blood pressure was still not adequately controlled were offered hydrochlorothiazide 12.5 mg and subsequently 25 mg as required to achieve adequate blood pressure goal. Patients were discontinued from the study if SeDBP was >120 mm Hg at any visit during Period III while on maximal

AZOR™ (amlodipine and olmesartan medoxomil)  
Formulary Dossier

therapy. If a patient experienced symptoms of hypotension or displayed intolerance to study medication at any time during Period III, the patient was back-titrated at the investigator's discretion. After Week 52, patients were discontinued from the study and treated per investigator's discretion. A follow-up visit two weeks later (Week 54) was scheduled to examine any safety issues.

Safety assessments included adverse events, evaluation of edema, clinical laboratory measurements of hematology and biochemistry, vital signs, physical examinations and 12-lead ECG assessments.

In the double-blind period, subgroup analyses were performed for each of the following groups: age group (<65 years, ≥65 years), diabetic status (yes, no), gender (male, female), race (Black, non-Black), ethnicity (Hispanic/Latino, non-Hispanic/Latino), hypertension class (Stage 1 hypertension, Stage 2 hypertension), prior antihypertensive medication use (naïve to antihypertensive medication, not naïve to antihypertensive medication), and baseline BMI (>30 kg/m<sup>2</sup>, ≤30 kg/m<sup>2</sup>).

**Results (Period II, Double-Blind):**

**Overall Baseline Characteristics:** Of the 4,234 patients screened for the study, 1,940 subjects were randomized equally into the 12 treatment arms for the double blind phase of the study. Of those randomized, 251 subjects were discontinued and 1,923 subjects were considered part of the intent to treat (ITT) population. Of the ITT population, 1,689 subjects completed the study.

Baseline characteristics for this study appear in Table 9, below. The treatment groups were comparable with respect to demographics, with no statistically significant differences among the treatment groups. Weight, height, and Body Mass Index (BMI) were also similar for the treatment groups, with no statistically significant differences among the treatment groups for these baseline characteristics. Treatment groups were also similar with respect to baseline values for blood pressure and heart rate, with no statistically significant differences among the treatment groups.

In regards to baseline hypertension class for the Safety Population, the treatment groups were similar, with over 70% of patients in each treatment group having Stage 2 hypertension. Overall, a total of 1538 (79.3%) patients had Stage 2 hypertension.

**Table 9: Baseline characteristics of randomized subjects (N =1,940)**

Age (years, mean ± SD)	54.0 ± 11.1
≥ 65 y (%)	19.8
≥ 75 y (n, %)	3.2
Males (%)	54.3
Blacks (%)	24.8
Subjects with diabetes (%)	13.5
Subjects anti-hypertensive agent naïve (%)	34.3
BMI (kg/M <sup>2</sup> , mean ± SD)	33.5 ± 7.1
SBP (mm Hg, mean ± SD)	163.8 ± 16.1
DBP (mm Hg, mean ± SD)	101.6 ± 5.2
Weight (kg, mean ± SD)	95.1 ± 21.9
Height (cm, mean ± SD)	170.1 ± 10.4
BMI (kg/m <sup>2</sup> , mean ± SD)	33.5 ± 7.1
Patients with BMI ≥ 30 kg/m <sup>2</sup> (%)	64.7
SD = standard deviation, BMI = body mass index, SBP = systolic blood pressure, DBP = diastolic blood pressure	

**Efficacy Results:** Each combination therapy had significantly greater reductions in SeDBP and SeSBP compared to both of its monotherapy components (Table 10, below).

AZOR™ (amlodipine and olmesartan medoxomil)  
Formulary Dossier

**Table 10: Mean Change in SeDBP and SeSBP (mm Hg) from Baseline to Week 8 with LOCF – Combination Therapy versus Monotherapy Comparisons – Intent to Treat Population**

Treatment Comparison			SeDBP Difference (Tmt 1 – Tmt 2)		SeSBP Difference (Tmt 1 – Tmt 2)	
Tmt 1	vs.	Tmt 2	LS Mean (SE)	p-value	LS Mean (SE)	p-value
AML5/OM10	vs.	OM10	-5.5 (0.99)	<0.0001	-11.7 (1.64)	<0.0001
	vs.	AML5	-4.3 (0.99)	<0.0001	-8.2 (1.63)	<0.0001
AML5/OM20	vs.	OM20	-4.7 (1.00)	<0.0001	-9.9 (1.65)	<0.0001
	vs.	AML5	-4.6 (0.99)	<0.0001	-8.3 (1.64)	<0.0001
AML5/OM40	vs.	OM40	-5.4 (1.00)	<0.0001	-9.7 (1.65)	<0.0001
	vs.	AML5	-6.3 (1.00)	<0.0001	-10.8 (1.65)	<0.0001
AML10/OM10	vs.	OM10	-7.8 (0.99)	<0.0001	-13.9 (1.64)	<0.0001
	vs.	AML10	-3.3 (0.99)	0.0004	-5.9 (1.63)	0.0002
AML10/OM20	vs.	OM20	-7.8 (1.00)	<0.0001	-15.4 (1.65)	<0.0001
	vs.	AML10	-4.4 (0.99)	<0.0001	-9.2 (1.64)	<0.0001
AML10/OM40	vs.	OM40	-8.5 (0.99)	<0.0001	-13.0 (1.64)	<0.0001
	vs.	AML10	-6.1 (0.99)	<0.0001	-9.6 (1.63)	<0.0001

LS Mean, SE, 95% CI, and one-sided p-values were obtained from an Analysis of Covariance model with fixed effects for treatment, age group, and diabetic status, and baseline as a covariate.  
AML = amlodipine, OM = olmesartan medoxomil, LS = least squares, SE = standard error, SeDBP = seated diastolic blood pressure, SeSBP = seated systolic blood pressure, Tmt = treatment.

The table (11) below presents the analysis results for mean change in SeDBP and SeSBP from baseline to Week 8 with LOCF for the ITT population. Each active treatment group had a statistically significant mean reduction in SeDBP and SeSBP from baseline to Week 8 with LOCF ( $p < 0.0001$ ). Mean reductions in SeDBP and SeSBP were greater in the combination therapy groups compared with the monotherapy groups. In the combination groups with AML 5 mg and 10 mg, increasing doses of OM (10 mg, 20 mg, and 40 mg) resulted in greater reductions in SeDBP. Overall, the greatest reductions in seated blood pressure occurred in the group treated with AML 10 mg + OM 40 mg (-30.1/-19.0 mm Hg) followed by the group treated with AML 10 mg + OM 20 mg (-29.2/-17.0 mmHg).

**Table 11: Mean Change in SeDBP and SeSBP (mm Hg) from Baseline to Week 8 with LOCF – Intent to Treat Population**

Treatment	N <sup>1</sup>	SeDBP Baseline to Week 8 with LOCF		SeSBP Baseline to Week 8 with LOCF	
		Change Mean ± SD	p-value <sup>2</sup>	Change Mean ± SD	p-value <sup>2</sup>
Placebo	160	-3.1 ± 10.67	<0.0001	-4.8 ± 18.70	0.0235
OM10	160	-8.3 ± 9.28	<0.0001	-11.5 ± 15.23	<0.0001
OM20	159	-9.2 ± 9.73	<0.0001	-13.8 ± 15.90	<0.0001
OM40	160	-10.2 ± 10.69	<0.0001	-16.1 ± 16.58	<0.0001
AML5	161	-9.4 ± 8.25	<0.0001	-14.9 ± 14.95	<0.0001
AML10	163	-12.7 ± 8.25	<0.0001	-19.7 ± 16.52	<0.0001
AML5/OM10	163	-13.8 ± 7.48	<0.0001	-24.2 ± 13.96	<0.0001
AML5/OM20	160	-14.0 ± 9.07	<0.0001	-23.6 ± 14.86	<0.0001
AML5/OM40	157	-15.5 ± 8.15	<0.0001	-25.4 ± 14.70	<0.0001
AML10/OM10	161	-16.0 ± 8.62	<0.0001	-25.3 ± 14.88	<0.0001
AML10/OM20	158	-17.0 ± 8.04	<0.0001	-29.2 ± 16.72	<0.0001
AML10/OM40	161	-19.0 ± 8.90	<0.0001	-30.1 ± 15.91	<0.0001

<sup>1</sup>N was the number of patients with values at both time points.  
<sup>2</sup>P-values were obtained from an Analysis of Covariance model with fixed effects for treatment, age group, and diabetic status, and baseline as a covariate.  
Week 8 with LOCF was defined as the last available measurement during the double-blind, active treatment period.  
AML = amlodipine, OM = olmesartan medoxomil, SD = standard deviation, SeDBP = seated diastolic blood pressure, SeSBP = seated systolic blood pressure



AZOR™ (amlodipine and olmesartan medoxomil)  
Formulary Dossier

Analysis results for the mean change in SeDBP and SeSBP from baseline to Week 2, Week 4, Week 6 and Week 8 without LOCF for the ITT population were similar to the results observed at Week 8 with LOCF. For all active treatment groups, most of the mean reduction in the SeDBP and SeSBP occurred from baseline to Week 2, and plateaued by Week 4.

The greater blood pressure reductions achieved with the combination treatments, compared with the respective monotherapies, resulted in a comparatively greater percentage of patients on combination therapy achieving their blood pressure treatment goals. The percentage of patients achieving their blood pressure goals by Week 8 with LOCF ranged from 20.0% to 36.3% for the groups treated with monotherapy compared with 35.0% to 53.2% for the groups treated with combination therapy. Approximately 50% of patients treated with one of the higher dose combination therapies (i.e., AML 10mg + OM 10 mg, AML 10 mg + OM 20 mg, AML 5 mg + OM 40 mg, and AML 10 mg + OM 40 mg) reached their blood pressure goal by Week 8 with LOCF.

In addition, the percentage of patients brought to a goal blood pressure of <140/90 mmHg ranged from 22.5% to 38.1% among the 5 monotherapy groups compared to 38.7% to 56.3% among the 6 combination therapy groups.

**Safety:** A total of 1020 (52.6% of 1940) patients experienced a treatment-emergent adverse event (TEAE) during the study. Across the 12 treatment groups the percentage experiencing a TEAE ranged from 45.3% to 58.9%, with no apparent trend in the distribution of TEAEs among the treatment groups (placebo incidence: 56.2%). Drug-related TEAEs were experienced by a total of 521 (26.9%) patients, ranging from 19.6% to 33.1% with the incidence in the placebo group being 29.6%. Across all treatment groups, most TEAEs and drug-related TEAEs were considered mild in intensity.

A total of 114 (5.9%) patients were discontinued from the study due to an adverse event. Edema was the most common drug-related TEAE and was experienced by a total of 277 (14.3%) patients. Edema was proactively assessed, and as a result, the observed incidences were higher than that reported in the package inserts for the components. There was a greater frequency of treatment-emergent edema among the groups that used AML 10 mg as a component of their treatment, with the AML 10 mg monotherapy group (36.8%) having the greatest incidence of edema. There appeared to be a decrease in the incidence when AML 10 mg was combined with OM 10 mg, 20 mg, and 40 mg (26.5%, 25.6%, and 23.5%). This reduction in the incidence of edema was not seen when AML 5mg was combined with OM 10 mg, 20 mg, or 40 mg (20.9%, 18.0%, and 18.5%) compared to AML 5 mg monotherapy group (13.0%). Other more common drug-related TEAEs included headache, experienced by 66 (3.4%) patients and dizziness experienced by 50 (2.6%) patients.

The frequency of hypertension (including the preferred terms of hypertension, systolic hypertension, diastolic hypertension, accelerated hypertension, blood pressure increased, blood pressure diastolic increased, or blood pressure inadequately controlled) reported as an adverse event was greatest in the placebo group (8.0%) and ranged from 0.6% to 5.0% among the active treatment groups. The combination treatment groups had a 0.0% to 1.9% incidence of adverse event of hypertension (lack of efficacy) reported as an adverse event.

A total of 7 patients experienced drug-related hypotension (1 patient in the OM 10 mg group, 2 patients in the OM 10 mg + AML 10 mg group, 1 patient in the OM 20 mg + AML 10 mg group, 1 patient in the OM 40 mg + AML 5 mg group, and 2 patients in the OM 40 mg + AML 10 mg treatment group).

There were no laboratory measurements that signified a safety concern. Furthermore, there were no clinically meaningful changes in heart rates, ECGs, or physical examinations for any treatment group during the study.

AZOR™ (amlodipine and olmesartan medoxomil)  
Formulary Dossier

**A Randomized, Double-Blind, Placebo-Controlled Factorial Study Evaluating The Efficacy and Safety of Co-Administration of Amlodipine Besylate Plus Olmesartan Medoxomil Compared to Monotherapy in Patients with Mild to Severe Hypertension (Subgroup Analyses of Blinded-Study Phase, United States)**

**Subgroup Analyses:** Analyses of prospectively identified subgroups from the 8-week controlled phase of this study. Subgroups were analyzed by age (<65 years vs. ≥65 years), diabetic status (subjects without diabetes vs. those with), and by race (Blacks vs. non-Blacks).

**Analysis by Age**

The mean baseline SeDBP for each subgroup was 102.0 mm Hg for subjects <65 years of age and 100.3 mm Hg for the subgroup ≥65 years of age. For both age subgroups, each active treatment group had a statistically significant mean reduction in SeDBP from baseline to Week 8 with LOCF ( $p < 0.0001$ ). The result of this subgroup analysis appears in Table 12, below.

**Table 12: Mean change in SeDBP (mm Hg) from Baseline to Week 8 with LOCF Stratified by Age Subgroups – Intent to Treat Population**

Treatment (mg)	<65 Years of Age			≥65 Years of Age		
	N <sup>1</sup>	Change Mean ± SD	p-value <sup>2</sup>	N <sup>1</sup>	Change Mean ± SD	p-value <sup>2</sup>
Placebo	128	-2.2 ± 10.69	0.0104	32	-6.4 ± 10.06	<0.0001
OM 10	128	-7.8 ± 9.03	<0.0001	32	-10.1 ± 10.13	<0.0001
OM 20	129	-8.3 ± 9.66	<0.0001	30	-13.2 ± 9.15	<0.0001
OM 40	129	-10.6 ± 10.07	<0.0001	31	-8.8 ± 13.04	<0.0001
AML 5	129	-8.3 ± 7.62	<0.0001	32	-13.7 ± 9.37	<0.0001
AML 10	131	-11.9 ± 8.27	<0.0001	32	-16.1 ± 7.33	<0.0001
AML 5/OM 10	131	-13.8 ± 7.84	<0.0001	32	-13.9 ± 5.85	<0.0001
AML 5/OM 20	126	-13.9 ± 8.97	<0.0001	34	-14.6 ± 9.54	<0.0001
AML 5/OM 40	126	-15.5 ± 8.44	<0.0001	31	-15.8 ± 6.96	<0.0001
AML 10/OM 10	130	-15.8 ± 8.59	<0.0001	31	-16.8 ± 8.83	<0.0001
AML 10/OM 20	126	-17.3 ± 8.07	<0.0001	32	-15.9 ± 7.98	<0.0001
AML 10/OM 40	128	-18.5 ± 9.17	<0.0001	33	-20.9 ± 7.59	<0.0001

<sup>1</sup>N was the number of subjects with values at both time points.  
<sup>2</sup>Two-sided p-values were obtained from an Analysis of Covariance model with treatment, age subgroup, and treatment-by subgroup interaction as fixed effects and baseline blood pressure as a covariate.  
 AML = amlodipine, OM = olmesartan medoxomil, SD = standard deviation.

For both age subgroups, subjects treated with AML 10 mg + OM 40 mg demonstrated the greatest overall mean reductions in SeDBP.

With respect to the systolic BP analysis, the mean baseline SeSBP was 161.4 mm Hg for the subgroup <65 years of age and 173.6 mm Hg for the subgroup ≥65 years of age. The results of this analysis appear in Table 13, below.

AZOR™ (amlodipine and olmesartan medoxomil)  
Formulary Dossier

**Table 13: Mean change in SeSBP (mm Hg) from Baseline to Week 8 with LOCF Stratified by Age Subgroups – Intent to Treat Population**

Treatment (mg)	<65 Years of Age			≥65 Years of Age		
	N <sup>1</sup>	Change Mean ± SD	p-value <sup>2</sup>	N <sup>1</sup>	Change Mean ± SD	p-value <sup>2</sup>
Placebo	128	-4.1 ± 18.54	0.0015	32	-7.9 ± 19.33	0.3430
OM 10	128	-10.9 ± 15.30	<0.0001	32	-13.9 ± 14.92	0.0002
OM 20	129	-12.5 ± 15.07	<0.0001	30	-19.4 ± 18.29	<0.0001
OM 40	129	-16.2 ± 15.63	<0.0001	31	-15.7 ± 20.33	<0.0001
AML 5	129	-13.3 ± 13.44	<0.0001	32	-21.1 ± 18.94	<0.0001
AML 10	131	-18.8 ± 16.39	<0.0001	32	-23.4 ± 16.82	<0.0001
AML 5/OM 10	131	-23.3 ± 14.38	<0.0001	32	-27.5 ± 11.70	<0.0001
AML 5/OM 20	126	-23.5 ± 15.38	<0.0001	34	-24.0 ± 12.95	<0.0001
AML 5/OM 40	126	-25.1 ± 13.62	<0.0001	31	-26.8 ± 18.66	<0.0001
AML 10/OM 10	130	-25.1 ± 14.79	<0.0001	31	-26.3 ± 15.45	<0.0001
AML 10/OM 20	126	-28.9 ± 15.86	<0.0001	32	-30.4 ± 19.98	<0.0001
AML 10/OM 40	128	-29.1 ± 16.30	<0.0001	33	-33.9 ± 13.88	<0.0001

<sup>1</sup>N was the number of subjects with values at both time points.  
<sup>2</sup>Two-sided p-values were obtained from an Analysis of Covariance model with treatment, age subgroup, and treatment-by subgroup interaction as fixed effects and baseline blood pressure as a covariate.  
 AML = amlodipine, OM = olmesartan medoxomil, SD = standard deviation.

For both age subgroups, each active treatment group demonstrated a statistically significant mean reduction in SeSBP from baseline to Week 8 with LOCF ( $p < 0.001$ ). For both age subgroups, the magnitude of the response was similar. The greatest mean reductions in SeSBP occurred in the group of subjects treated with AML 10 mg + OM 40 mg.

With respect to comparisons of combination therapy versus monotherapy for SeDBP and SeSBP (from baseline to Week 8 with LOCF) in patients less than 65 years of age, each combination therapy had a significantly greater mean reduction in SeDBP and SeSBP compared to the respective monotherapy components ( $p < 0.001$  for all comparisons).

For patients greater than or equal to 65, not all comparisons of combination therapy versus monotherapy resulted in significantly greater reductions for mean change in SeDBP and SeSBP from baseline to Week 8 with LOCF. Table 14 provides results of the comparisons between combination therapy versus monotherapy for patients greater than or equal to 65.

AZOR™ (amlodipine and olmesartan medoxomil)  
Formulary Dossier

**Table 14: Mean Change in SeDBP and SeSBP (mm Hg) from Baseline to Week 8 with LOCF – Combination Therapy Versus Monotherapy Comparisons – Patients ≥ 65 Years of Age**

Treatment Comparison			SeDBP Difference (Tmt 1 – Tmt 2)		SeSBP Difference (Tmt 1 – Tmt 2)	
Tmt 1	vs.	Tmt 2	LS Mean (SE)	p-value	LS Mean (SE)	p-value
AML5/OM10	vs.	OM10	-3.9 (2.22)	0.0380	-14.1 (3.68)	<0.0001
	vs.	AML5	-0.2 (2.22)	0.4633	-6.6 (3.68)	0.0367
AML5/OM20	vs.	OM20	-1.3 (2.22)	0.2742	-6.0 (3.69)	0.0523
	vs.	AML5	-0.8 (2.18)	0.3655	-4.2 (3.63)	0.1254
AML5/OM40	vs.	OM40	-7.4 (2.25)	0.0005	-12.9 (3.74)	0.0003
	vs.	AML5	-2.3 (2.23)	0.1467	-7.8 (3.71)	0.0183
AML10/OM10	vs.	OM10	-6.6 (2.23)	0.0017	-14.3 (3.71)	<0.0001
	vs.	AML10	-0.5 (2.23)	0.4174	-3.5 (3.71)	0.1748
AML10/OM20	vs.	OM20	-2.8 (2.25)	0.1057	-12.2 (3.74)	0.0006
	vs.	AML10	0.1 (2.22)	0.4767	-7.0 (3.68)	0.0282
AML10/OM40	vs.	OM40	-11.9 (2.22)	<0.0001	-15.7 (3.69)	<0.0001
	vs.	AML10	-4.5 (2.20)	0.0207	-7.2 (3.66)	0.0250

LS Mean, SE, 95% CI, and one-sided p-values were obtained from an Analysis of Covariance model with fixed effects for treatment, age group, and diabetic status, and baseline as a covariate.  
AML = amlodipine, OM = olmesartan medoxomil, LS = least squares, SE = standard error, SeDBP = seated diastolic blood pressure, SeSBP = seated systolic blood pressure, Tmt = treatment.

For the subgroup of patients <65 years of age, approximately 51% to 56% of patients treated with one of the higher dose combination therapies (i.e., AML 10 mg + OM 10 mg, AML 10 mg + OM 20 mg, AML 5 mg + OM 40 mg, and AML 10 mg + OM 40 mg) reached their blood pressure goal.

The subgroup of patients ≥65 years of age had a comparatively lower percentage of patients who reached their blood pressure goal, which was likely due to the higher baseline SeSBP in this subgroup. The mean baseline blood pressure for patients ≥ 65 was 173.6/100.3 mm Hg. Approximately 21% to 44% of patients treated with one of the higher dose combination therapies (i.e., AML 10 mg + OM 10 mg, AML 10 mg + OM 20 mg, AML 5 mg + OM 40 mg, and AML 10 mg + OM 40 mg) reached their blood pressure goal.

AZOR™ (amlodipine and olmesartan medoxomil)  
Formulary Dossier

### Analysis by Diabetes Status

The mean baseline SeDBP was 101.7 mmHg for the subgroup of subjects without diabetes and 101.1 mm Hg for the subgroup of subjects with diabetes. For both subgroups, each active treatment group had a statistically significant mean reduction in SeDBP from baseline to Week 8 with LOCF ( $p < 0.0001$ ). The results of this subgroup analysis appear in Table 15, below

**Table 15. Mean change in SeDBP (mm Hg) from Baseline to Week 8 with LOCF Stratified by Diabetes Status – Intent to Treat Population**

Treatment (mg)	Without Diabetes			With Diabetes		
	N <sup>1</sup>	Change Mean $\pm$ SD	p-value <sup>2</sup>	N <sup>1</sup>	Change Mean $\pm$ SD	p-value <sup>2</sup>
Placebo	137	-2.2 $\pm$ 9.63	0.0065	23	-8.2 $\pm$ 14.77	<0.0001
OM 10	140	-8.0 $\pm$ 9.50	<0.0001	20	-9.9 $\pm$ 7.50	<0.0001
OM 20	137	-9.2 $\pm$ 9.38	<0.0001	22	-9.4 $\pm$ 11.96	<0.0001
OM 40	139	-10.5 $\pm$ 10.33	<0.0001	21	-8.3 $\pm$ 12.94	<0.0001
AML 5	139	-9.0 $\pm$ 7.86	<0.0001	22	-11.6 $\pm$ 10.36	<0.0001
AML 10	140	-12.9 $\pm$ 8.35	<0.0001	23	-11.7 $\pm$ 7.69	<0.0001
AML 5/OM 10	140	-13.6 $\pm$ 7.65	<0.0001	23	-15.1 $\pm$ 6.34	<0.0001
AML 5/OM 20	138	-14.9 $\pm$ 9.02	<0.0001	22	-8.3 $\pm$ 7.23	<0.0001
AML 5/OM 40	140	-15.6 $\pm$ 8.16	<0.0001	17	-14.6 $\pm$ 8.26	<0.0001
AML 10/OM 10	141	-16.0 $\pm$ 8.49	<0.0001	20	-16.0 $\pm$ 9.75	<0.0001
AML 10/OM 20	137	-17.3 $\pm$ 8.26	<0.0001	21	-15.0 $\pm$ 6.20	<0.0001
AML 10/OM 40	137	-19.1 $\pm$ 9.08	<0.0001	24	-18.4 $\pm$ 7.95	<0.0001

<sup>1</sup>N was the number of subjects with values at both time points.  
<sup>2</sup>Two-sided p-values were obtained from an Analysis of Covariance model with treatment, diabetes status subgroup, and treatment-by subgroup interaction as fixed effects and baseline blood pressure as a covariate.  
 AML = amlodipine, OM = olmesartan medoxomil, SD = standard deviation.

For the subgroup of subjects without diabetes, increases in dose were associated with progressively greater mean reductions in SeDBP across all treatment groups, which was less consistent for the subgroup of subjects with diabetes. These findings may be a consequence of the relatively small number of subjects with diabetes in each treatment group. Overall, the greatest mean reductions in SeDBP occurred in the group of subjects treated with AML 10 mg + OM 40 mg.

The mean baseline SeSBP was 163.1 mmHg for the subgroup of subjects without diabetes and 168.6 mmHg for the subgroup of subjects with diabetes. For both subgroups, each active treatment group showed a statistically significant mean reduction in SeSBP from baseline to Week 8 with LOCF ( $p < 0.003$ ). The results of this subgroup analysis appear in Table 16, below.

AZOR™ (amlodipine and olmesartan medoxomil)  
Formulary Dossier

**Table 16. Mean Change in SeSBP (mm Hg) from Baseline to Week 8 with LOCF Stratified by Diabetes Status – Intent to Treat Population**

Treatment (mg)	Without Diabetes			With Diabetes		
	N <sup>1</sup>	Change Mean ± SD	p-value <sup>2</sup>	N <sup>1</sup>	Change Mean ± SD	p-value <sup>2</sup>
Placebo	137	-3.1 ± 15.50	0.0368	23	-15.3 ± 30.18	0.0003
OM 10	140	-11.1 ± 15.58	<0.0001	20	-14.4 ± 12.49	<0.0001
OM 20	137	-14.2 ± 15.70	<0.0001	22	-11.6 ± 17.28	0.0010
OM 40	139	-16.9 ± 15.88	<0.0001	21	-10.5 ± 20.17	0.0026
AML 5	139	-14.0 ± 14.29	<0.0001	22	-20.3 ± 18.04	<0.0001
AML 10	140	-20.1 ± 16.72	<0.0001	23	-17.7 ± 15.48	<0.0001
AML 5/OM 10	140	-23.9 ± 13.87	<0.0001	23	-25.6 ± 14.79	<0.0001
AML 5/OM 20	138	-25.1 ± 14.67	<0.0001	22	-14.2 ± 12.67	<0.0001
AML 5/OM 40	140	-25.5 ± 14.23	<0.0001	17	-25.0 ± 18.65	<0.0001
AML 10/OM 10	141	-25.0 ± 14.33	<0.0001	20	-27.9 ± 18.49	<0.0001
AML 10/OM 20	137	-29.7 ± 16.92	<0.0001	21	-26.3 ± 15.44	<0.0001
AML 10/OM 40	137	-30.1 ± 16.40	<0.0001	24	-30.3 ± 13.08	<0.0001

<sup>1</sup>N was the number of subjects with values at both time points.  
<sup>2</sup>Two-sided p-values were obtained from an Analysis of Covariance model with treatment, diabetes status subgroup, and treatment-by subgroup interaction as fixed effects and baseline blood pressure as a covariate.  
 AML = amlodipine, OM = olmesartan medoxomil, SD = standard deviation.

As observed for SeDBP, increases in dose were associated with progressively greater mean reductions in SeSBP for the subgroup of subjects without diabetes across all treatment groups, which was less consistent for the subgroup of subjects with diabetes. Similarly, these findings may be a consequence of the relatively small number of subjects with diabetes in each treatment group. In addition, the greatest mean reductions in SeDBP occurred in the group of subjects treated with AML 10 mg + OM 40 mg.

With respect to comparisons of combination therapy versus monotherapy for SeDBP and SeSBP (from baseline to Week 8 with LOCF) in patients without diabetes, each combination therapy had a significantly greater mean reduction in SeDBP and SeSBP compared to both of its monotherapy components (p<0.0016 for all comparisons).

For patients with diabetes, not all comparisons of combination therapy versus monotherapy resulted in significantly greater reductions for mean change in SeDBP and SeSBP from baseline to Week 8 with LOCF. Table 17 provides results of the comparisons between combination therapy versus monotherapy for patients with diabetes.

AZOR™ (amlodipine and olmesartan medoxomil)  
Formulary Dossier

**Table 17: Mean Change in SeDBP and SeSBP (mm Hg) from Baseline to Week 8 with LOCF – Combination Therapy Versus Monotherapy Comparisons – Patients With Diabetes**

Treatment Comparison			SeDBP Difference (Tmt 1 – Tmt 2)		SeSBP Difference (Tmt 1 – Tmt 2)	
Tmt 1	vs.	Tmt 2	LS Mean (SE)	p-value	LS Mean (SE)	p-value
AML5/OM10	vs.	OM10	-4.5 (2.72)	0.0494	-7.1 (4.48)	0.0566
	vs.	AML5	-3.5 (2.65)	0.0921	-4.2 (4.37)	0.1655
AML5/OM20	vs.	OM20	1.0 (2.68)	0.3563	-3.7 (4.41)	0.1982
	vs.	AML5	3.2 (2.68)	0.1141	4.1 (4.42)	0.1764
AML5/OM40	vs.	OM40	-6.5 (2.90)	0.0121	-14.8 (4.78)	0.0010
	vs.	AML5	-3.1 (2.87)	0.1375	-6.2 (4.73)	0.0935
AML10/OM10	vs.	OM10	-5.5 (2.82)	0.0265	-10.2 (4.63)	0.0138
	vs.	AML10	-4.4 (2.72)	0.0529	-9.6 (4.48)	0.0162
AML10/OM20	vs.	OM20	-5.9 (2.71)	0.0154	-14.5 (4.47)	0.0006
	vs.	AML10	-3.5 (2.69)	0.0935	-8.9 (4.42)	0.0219
AML10/OM40	vs.	OM40	-10.0 (2.66)	<0.0001	-18.1 (4.38)	<0.0001
	vs.	AML10	-6.5 (2.60)	0.0061	-11.8 (4.27)	0.0030

LS Mean, SE, 95% CI, and one-sided p-values were obtained from an Analysis of Covariance model with fixed effects for treatment, age group, and diabetic status, and baseline as a covariate.  
AML = amlodipine, OM = olmesartan medoxomil, LS = least squares, SE = standard error, SeDBP = seated diastolic blood pressure, SeSBP = seated systolic blood pressure, Tmt = treatment.

For the subgroup of patients without diabetes, approximately 55% to 60% of patients treated with one of the higher dose combination therapies (i.e., AML 10 mg + OM 10 mg, AML 10 mg + OM 20 mg, AML 5 mg + OM 40 mg, and AML 10 mg + OM 40 mg ) reached their blood pressure goal.

The subgroup with diabetes, which started with a higher mean baseline blood pressure of 168.6/101.1, had a lower percentage of patients who reached their blood pressure goal. For the subgroup of patients with diabetes, approximately 10% to 13% of patients treated with one of the higher dose combination therapies (i.e., AML 10 mg + OM 10 mg, AML 10 mg + OM 20 mg, AML 5 mg + OM 40 mg, and AML 10 mg + OM 40 mg) reached their blood pressure goal.

AZOR™ (amlodipine and olmesartan medoxomil)  
Formulary Dossier

### Analysis by Race

The mean baseline SeDBP was 102.4 mm Hg for the subgroup of Black subjects and 101.4 mm Hg for the subgroup of non-Black subjects. For both race subgroups, each active treatment group had a statistically significant mean reduction in SeDBP from baseline to Week 8 with LOCF ( $p < 0.01$ ). The result of this subgroup analysis appear in Table 18, below.

**Table 18: Mean Change in SeDBP (mm Hg) from Baseline to Week 8 with LOCF Stratified by Race (Black vs. non-Black) – Intent to Treat Population**

Treatment	Black Subjects			Non-Black Subjects		
	N <sup>1</sup>	Change Mean $\pm$ SD	p-value <sup>2</sup>	N <sup>1</sup>	Change Mean $\pm$ SD	p-value <sup>2</sup>
Placebo	45	-1.3 $\pm$ 9.55	0.4587	115	3.8 $\pm$ 11.04	<0.0001
OM 10	32	-5.3 $\pm$ 8.44	0.0012	128	-9.0 $\pm$ 9.35	<0.0001
OM 20	34	-4.5 $\pm$ 9.98	0.0032	125	-10.5 $\pm$ 9.30	<0.0001
OM 40	44	-5.5 $\pm$ 9.51	<0.0001	116	-12.0 $\pm$ 10.61	<0.0001
AML 5	42	-8.3 $\pm$ 8.66	<0.0001	119	-9.7 $\pm$ 8.11	<0.0001
AML 10	39	-13.4 $\pm$ 8.40	<0.0001	124	-12.5 $\pm$ 8.22	<0.0001
AML 5/OM 10	34	-9.4 $\pm$ 6.94	<0.0001	129	-15.0 $\pm$ 7.20	<0.0001
AML 5/OM 20	43	-12.4 $\pm$ 9.17	<0.0001	117	-14.6 $\pm$ 9.00	<0.0001
AML 5/OM 40	38	-13.9 $\pm$ 8.35	<0.0001	119	-16.0 $\pm$ 8.06	<0.0001
AML 10/OM 10	43	-15.5 $\pm$ 8.45	<0.0001	118	-16.2 $\pm$ 8.71	<0.0001
AML 10/OM 20	46	-15.2 $\pm$ 7.92	<0.0001	112	-17.8 $\pm$ 8.01	<0.0001
AML 10/OM 40	34	-15.7 $\pm$ 9.05	<0.0001	127	-19.9 $\pm$ 8.68	<0.0001

<sup>1</sup>N was the number of subjects with values at both time points.  
<sup>2</sup>Two-sided p-values were obtained from an Analysis of Covariance model with treatment, race subgroup, and treatment-by-subgroup interaction as fixed effects and baseline blood pressure as a covariate.  
 AML = amlodipine, OM = olmesartan medoxomil, SD = standard deviation.

There was a general association between increases in dose and greater mean reductions in SeDBP across the treatment groups for both the Black and non-Black subgroups. Across the range of combination therapies, the non-Black subgroup had numerically greater mean reductions in SeDBP compared with the Black subgroup. For both race subgroups, the greatest mean reductions in SeDBP occurred in the cohort of subjects treated with AML 10 mg + OM 40 mg.

The mean baseline SeSBP was 163.9 mmHg for the subgroup of Black subjects and 163.8 mmHg for the subgroup of non-Black subjects. All active treatment groups had statistically significant ( $p < 0.05$ ) mean reductions in SeSBP from baseline to Week 8 with LOCF for both race subgroups. The results of this subgroup analysis appear in Table 19, below.



AZOR™ (amlodipine and olmesartan medoxomil)  
Formulary Dossier

**Table 19: Mean Change in SeSBP (mm Hg) from Baseline to Week 8 with LOCF Stratified by Race (Black vs. non-Black) – Intent to Treat Population**

Treatment	Black Subjects			Non-Black Subjects		
	N <sup>1</sup>	Change Mean ± SD	p-value <sup>2</sup>	N <sup>1</sup>	Change Mean ± SD	p-value <sup>2</sup>
Placebo	45	-4.3 ± 21.29	0.2150	115	-5.0 ± 17.69	0.0017
OM 10	32	-6.0 ± 12.30	0.0322	128	-12.9 ± 15.62	<0.0001
OM 20	34	-5.5 ± 17.06	0.0139	125	-16.1 ± 14.84	<0.0001
OM 40	44	-8.2 ± 16.07	0.0008	116	-19.1 ± 15.83	<0.0001
AML 5	42	-11.9 ± 13.40	<0.0001	119	-15.9 ± 15.39	<0.0001
AML 10	39	-22.1 ± 15.12	<0.0001	124	-19.0 ± 16.93	<0.0001
AML 5/OM 10	34	-18.8 ± 12.53	<0.0001	129	-25.6 ± 14.02	<0.0001
AML 5/OM 20	43	-23.7 ± 12.57	<0.0001	117	-23.5 ± 15.66	<0.0001
AML 5/OM 40	38	-24.7 ± 13.84	<0.0001	119	-25.7 ± 15.01	<0.0001
AML 10/OM 10	43	-24.1 ± 16.10	<0.0001	118	-25.8 ± 14.45	<0.0001
AML 10/OM 20	46	-25.3 ± 13.76	<0.0001	112	-30.9 ± 17.59	<0.0001
AML 10/OM 40	34	-28.7 ± 14.85	<0.0001	127	-30.5 ± 16.22	<0.0001

<sup>1</sup>N was the number of subjects with values at both time points.  
<sup>2</sup>Two-sided p-values were obtained from an Analysis of Covariance model with treatment, race subgroup, and treatment-by subgroup interaction as fixed effects and baseline blood pressure as a covariate.  
 AML = amlodipine, OM = olmesartan medoxomil, SD = standard deviation.

For both race subgroups, mean reductions in SeSBP were, in general, numerically greater in the combination therapy groups compared with the monotherapy groups. Across most combination therapies, the non-Black subgroup had numerically greater mean reductions in SeSBP compared with the Black subgroup. For both race subgroups, the greatest mean reductions in SeSBP occurred in the groups treated with AML 10 mg + OM 40 mg.

With respect to comparisons of combination therapy versus monotherapy for SeDBP and SeSBP (from baseline to Week 8 with LOCF) in the subgroup of non-Black patients, each combination therapy had a significantly greater mean reduction in SeDBP and SeSBP compared to both of its monotherapy components (p<0.001 for all comparisons).

For the subgroup of Black patients, not all comparisons of combination therapy versus monotherapy resulted in significantly greater reductions for mean change in SeDBP and SeSBP from baseline to Week 8 with LOCF. Table 20 provides results of the comparisons between combination therapy versus monotherapy for the subgroup of Black patients.

AZOR™ (amlodipine and olmesartan medoxomil)  
Formulary Dossier

**Table 20: Mean Change in SeDBP and SeSBP (mm Hg) from Baseline to Week 8 with LOCF – Combination Therapy Versus Monotherapy Comparisons – Black Patients**

Treatment Comparison			SeDBP Difference (Tmt 1 – Tmt 2)		SeSBP Difference (Tmt 1 – Tmt 2)	
Tmt 1	vs.	Tmt 2	LS Mean (SE)	p-value	LS Mean (SE)	p-value
AML5/OM10	vs.	OM10	-4.0 (2.17)	0.0322	-13.1 (3.58)	0.0001
	vs.	AML5	-1.0 (2.03)	0.3066	-5.2 (3.36)	0.0605
AML5/OM20	vs.	OM20	-7.6 (2.02)	<0.0001	-16.6 (3.34)	<0.0001
	vs.	AML5	-4.1 (1.91)	0.0163	-9.3 (3.16)	0.0016
AML5/OM40	vs.	OM40	-8.5 (1.95)	<0.0001	-17.1 (3.22)	<0.0001
	vs.	AML5	-5.8 (1.97)	0.0018	-11.1 (3.26)	0.0004
AML10/OM10	vs.	OM10	-10.4 (2.05)	<0.0001	-19.7 (3.40)	<0.0001
	vs.	AML10	-2.0 (1.94)	0.1549	-1.9 (3.22)	0.2759
AML10/OM20	vs.	OM20	-10.8 (1.99)	<0.0001	-20.0 (3.29)	<0.0001
	vs.	AML10	-1.7 (1.91)	0.1851	-2.8 (3.17)	0.1848
AML10/OM40	vs.	OM40	-9.9 (2.01)	<0.0001	-19.9 (3.32)	<0.0001
	vs.	AML10	-1.7 (2.07)	0.2023	-3.9 (3.42)	0.1255
LS Mean, SE, 95% CI, and one-sided p-values were obtained from an Analysis of Covariance model with fixed effects for treatment, age group, and diabetic status, and baseline as a covariate. AML = amlodipine, OM = olmesartan medoxomil, LS = least squares, SE = standard error, SeDBP = seated diastolic blood pressure, SeSBP = seated systolic blood pressure, Tmt = treatment.						

For the subgroup of Black patients, approximately 38% to 51% of patients treated with one of the higher dose combination therapies (i.e., AML 10 mg + OM 10 mg, AML 10 mg + OM 20 mg, AML 5 mg + OM 40 mg, and AML 10 mg + OM 40 mg) reached their blood pressure goal.

For these same higher dose combinations, the subgroup of non-Black patients had a percentage of patients who reached their blood pressure goal that ranged from approximately 48% to 56%.

Generally, the response to all therapy groups, monotherapy and combination, was found to be greater in the non-Black population, except for the AML 10 mg monotherapy.

AZOR™ (amlodipine and olmesartan medoxomil)  
Formulary Dossier

**A Randomized, Double-Blind, Placebo-Controlled Factorial Study Evaluating The Efficacy And Safety of Co-Administration of Amlodipine Besylate Plus Olmesartan Medoxomil Compared to Monotherapy in Patients with Mild to Severe Hypertension (Long-Term, Open-Label Study Phase, United States)**

**Objectives:** The main objective of the open-label Period III (Week 8 through Week 52) phase of the pivotal trial was to test long-term safety, and the durability of effect of co-administration of amlodipine (AML) and olmesartan medoxomil (OM) (plus the addition of hydrochlorothiazide [HCTZ], if needed) while minimally treating patients to blood pressure goal (<140/90 mm Hg, <130/80 mm Hg for diabetic patients). A further objective was to evaluate the number (percentage) of patients achieving blood pressure goal (defined as blood pressure <140/90 mm Hg, <130/80 mm Hg for diabetic patients).

**Methodology:** Period III consisted of a 44-week, open-label treatment period to assess long-term safety and efficacy of various treatment combinations. After completing Period II, all patients were switched to the combination of AML 5 mg + OM 40 mg. Those patients whose blood pressure was not adequately controlled (i.e., did not achieve a blood pressure goal of <140/90 mm Hg, or <130/80 mm Hg for those patients with diabetes) on AML 5 mg + OM 40 mg were titrated to AML 10 mg + OM 40 mg. Patients whose blood pressure was still not adequately controlled were additionally offered HCTZ 12.5 mg and subsequently 25 mg as required to achieve this blood pressure goal.

Safety assessments included adverse events, evaluation of edema, clinical laboratory measurements, vital signs, physical examinations, 12-lead ECG assessments, and special chemistry analytes.

**Efficacy Results for the Open-Label Period:** A total of 1684 patients entered Period III of the trial. After 2 weeks of the open-label treatment period (Week 10), 1640 patients remained on AML 5 mg + OM 40 mg with a mean SeDBP of 86.0 mmHg and a mean SeSBP of 137.9 mmHg. A total of 48.3% (792 of 1640) of patients on AML 5 mg + OM 40 mg reached their blood pressure treatment goal within 2 weeks of treatment. As patients were titrated to more intensive treatment regimens, an overall greater percentage of patients reached their blood pressure treatment goal.

From the mean baseline blood pressure of 163.6/101.5 mmHg from Period II, blood pressure reductions were observed across all combination treatment regimens to Week 52. At Week 52 or ET (early termination), the mean SeDBP in patients who remained on AML 5 mg + OM 40 mg was 81.0 mmHg and the mean SeSBP was 127.6 mmHg. A total of 80.0% of patients who remained on AML 5 mg + OM 40 mg achieved the blood pressure goal. For patients whose treatment regimen was titrated to more intensive antihypertensive regimens, mean SeDBP and SeSBP at Week 52 were slightly higher than for patients who remained on the AML 5 mg + OM 40 mg treatment regimen. Despite more intensive antihypertensive regimens, lower percentages of patients achieved blood pressure goals. The groups of patients who required titration of the amlodipine dose or the addition of hydrochlorothiazide were more severe hypertensive patients and/or were more resistant to the antihypertensive effects of treatment.

Table 21, below, presents the BP reductions observed at Week 52 or Early Termination, and the percentage of patients who achieved BP goal (i.e., <140/90 mm Hg or <130/80 for patients with diabetes) in Period III of this study. Note that HCTZ 12.5 mg or 25 mg was added to those patients who did not reach goal on one of the treatment regimens, or where edema persisted.

**Table 21: SeSBP and SeDBP at Week 52/Early Termination and Blood Pressure Goal Rates**

Treatment (mgs)	No. of Patients	Mean SeSBP (mmHg)	Mean SeDBP (mmHg)	% of Subjects Reaching BP Goal*
AML 5/OM 40	525	127.6	81.0	80.0
AML 10/OM 40	378	130.9	82.4	70.6
AML 10/OM 40 + HCTZ 12.5	287	130.7	81.0	66.6
AML 10/OM 40 + HCTZ 25	419	136.8	83.4	46.3

\*<140/90 mmHg or <130/80 for patients with diabetes

Sixty-three (63) patients received other non-standard treatment regimens that included HCTZ 25 mg, AML 5 mg, AML 10 mg + OM 20 mg, AML 2.5 mg + OM 20 mg, AML 5 mg + OM 20 mg, AML 5 mg + OM 20 mg + HCTZ 12.5 mg, AML 10 mg + OM 20 mg + HCTZ 12.5 mg, AML 2.5 mg + OM 40 mg + HCTZ 12.5 mg, AML 5 mg + OM 40 mg + HCTZ 12.5 mg, and AML 5 mg + OM 40 mg + HCTZ 25 mg. Taken as a separate cohort, these patients had a mean SeDBP of 79.4 mmHg and a mean SeSBP of 126.2 mmHg, with 68.3% of these patients reaching their BP goal.

AZOR™ (amlodipine and olmesartan medoxomil)  
Formulary Dossier

The treatment-to-goal approach used during the open-label treatment period resulted in an increase in the overall number and percentage of patients who reached their blood pressure goal over time. A total of 48.2% (792 of 1640 patients) reached their blood pressure goal at Week 10. By Week 52/ET, the total percentage who reached their blood pressure goal increased to 66.7% (1115 of 1672 patients).

Each titration to a more intensive treatment regimen resulted in additional mean reductions in seated blood pressure. Titration from AML 5 mg + OM 40 mg to AML 10 mg + OM 40 mg resulted in a mean seated blood pressure reduction of 7.3/4.8 mmHg. The addition of HCTZ 12.5 mg to the AML 10 mg + OM 40 mg combination resulted in a mean seated blood pressure reduction of 7.7/4.5 mmHg. A further reduction in seated blood pressure of 9.9/6.0 mmHg resulted from the titration in HCTZ from 12.5 mg to 25 mg for patients on the maximum-strength treatment regimen of AML 10 mg + OM 40 mg + HCTZ 25 mg.

**Safety Results for the Open-Label Period:** No new safety issues were identified during the course of this study with any of the combination therapies. During the open-label period, adverse events were experienced by 622 (37.0%) patients on AML 5 mg + OM 40 mg, 455 (40.5%) patients on AML 10 mg + OM 40 mg, 312 (42.4%) patients on AML 10 mg + OM 40 mg + HCTZ 12.5 mg, and 248 (56.4%) patients on AML 10 mg + OM 40 mg + HCTZ 25 mg.

Drug-related adverse events were experienced by 221 (13.2%) patients on AML 5 mg + OM 40 mg, 195 (17.3%) patients on AML 10 mg + OM 40 mg, 124 (16.8%) patients on AML 10 mg + OM 40 mg + HCTZ 12.5 mg, and 89 (20.2%) patients on AML 10 mg + OM 40 mg + HCTZ 25 mg.

Across all treatment regimens, most adverse events and drug-related adverse events were considered mild to moderate in severity. There were no apparent trends among the treatment groups with regard to the maximum severity of either adverse events or drug-related adverse events.

One patient on AML 10 mg + OM 40 mg died as a result of a gunshot wound to the head during the open-label treatment period. This death was not related to use of the study medication. Serious adverse events (SAEs) were experienced by 31 (1.8%) patients on AML 5 mg + OM 40 mg, 23 (2.0%) patients on AML 10 mg + OM 40 mg, 15 (2.0%) patients on AML 10 mg + OM 40 mg + HCTZ 12.5 mg, and 18 (4.1%) patients on AML 10 mg + OM 40 mg + HCTZ 25 mg. Four patients experienced SAEs while on non-standard treatment regimens. One patient on AML 5 mg + OM 40 mg had a drug-related SAE (non-cardiac chest pain).

A total of 77 (4.6%) patients were discontinued during the open-label treatment period due to an adverse event. Seventy (4.2%) patients were discontinued from the study for adverse events that started during the open-label treatment period. Adverse event discontinuations were experienced by 28 (1.7%) patients on AML 5 mg + OM 40 mg, 17 (1.5%) patients on AML 10 mg + OM 40 mg, 11 (1.5%) patients on AML 10 mg + OM 40 mg + HCTZ 12.5 mg, and 11 (2.5%) patients on AML 10 mg + OM 40 mg + HCTZ 25 mg. Three (3.5%) patients experienced an adverse event discontinuation while on a non-standard treatment regimen. Seven (0.4%) other patients were discontinued from the study during the open-label period for an adverse event that started during the double-blind treatment period. Thirty-five (2.1%) patients were discontinued due to drug-related adverse events.

A review of the frequency and specific types of events, as well as an assessment of potential relationships between the events and study medication, suggest that there was no greater incidence of SAEs or adverse event discontinuations due to the combination of OM and AML, or the combination of OM, AML, and HCTZ.

Edema (including the preferred terms of edema, edema peripheral, and pitting edema) was the most common drug-related adverse event and was experienced by a total of 318 (18.9%) patients. Across the 4 standard treatment regimens, the frequency of drug-related edema ranged from 7.0% to 11.1%. Dizziness (ranging from 1.4% to 2.0%) was the other most common drug-related, treatment-emergent adverse event.

In addition to assessing the incidence of adverse events categorized as edema, shifts in the severity of peripheral edema were also evaluated. Throughout the open-label treatment period, there appeared to be an approximate doubling in the frequency of patients with shifts toward worsening peripheral edema grade with AML 10 mg + OM 40 mg treatment (12.2%) compared to AML 5 mg + OM 40 mg treatment (7.2%). The apparent worsening in peripheral edema grade associated with the increased dose of amlodipine from 5 mg to 10 mg was somewhat ameliorated when HCTZ was added to the dosing regimen. The percentage of patients with shifts toward worsening peripheral edema grade was 10.3% for patients treated with AML 10 mg + OM 40 mg + HCTZ 12.5 mg, and 9.1% for patients treated with AML 10 mg + OM 40 mg + HCTZ 25 mg.

AZOR™ (amlodipine and olmesartan medoxomil)  
Formulary Dossier

Overall, there was a low incidence of hypotension across the treatment regimens. Hypotension was experienced by 0.8% to 1.6% of patients on the 3 lowest dose treatment regimens and by 0.7% on AML 10 mg + OM 40 mg + HCTZ 25 mg.

There were no laboratory measurements that signaled a safety concern. Furthermore, there were no clinically meaningful changes in heart rates, ECGs, or physical examinations for any treatment group during the study.

AZOR™ (amlodipine and olmesartan medoxomil)  
Formulary Dossier

**Efficacy and Safety of Amlodipine Used as Add-On Therapy in Moderately to Severely Hypertensive Patients Not Adequately Controlled by Olmesartan Medoxomil 20 mg Monotherapy (Phase III, Europe)**

**Objectives:** The primary objective was to demonstrate the additional antihypertensive efficacy in lowering trough sitting diastolic blood pressure (DBP) gained by adding amlodipine (AML) 5 mg or 10 mg to the treatment regimen in patients with hypertension not adequately controlled on olmesartan medoxomil (OM) 20 mg alone as assessed by conventional blood pressure (BP) measurements after 8 weeks of double-blind treatment. The primary efficacy endpoint was the mean change from baseline to the end of the double-blind treatment period, using last observation carried forward (LOCF) in trough sitting DBP.

The secondary objectives of this study consisted of an evaluation of the trough SeSBP after 4 weeks and 8 weeks of double-blind treatment in comparison to OM 20 mg. Additionally, SeDBP was also evaluated after 4 weeks of double-blind treatment. Ambulatory blood pressure monitoring (ABPM) was utilized to evaluate antihypertensive efficacy (daytime, nighttime, and 24 hour) in DBP and SBP after 8 weeks of treatment and blood pressure goal (defined as BP <140/90 mm Hg, <130/80 mmHg for diabetic patients) was also evaluated after 4 and 8 weeks of double-blind therapy. Lastly, safety and tolerability were assessed for the combinations of OM and AML versus monotherapy with OM 20 mg.

**Methodology:** This was a multi-center, multi-national, randomized, double-blind, parallel-group trial of 538 patients with moderate to severe hypertension (SBP  $\geq$  160 mm Hg and DBP  $\geq$  100 mm Hg) or not controlled on olmesartan monotherapy (BP  $\geq$  140/90 mm Hg). The study consisted of a 1-2 week taper off period for eligible patients previously on antihypertensive treatment (other than olmesartan monotherapy), followed by 16 weeks of active treatment. For the first 8 weeks (Period I), all patients received open-label once-daily doses of olmesartan 20 mg. At the end of the 8 weeks, only non-responders were then randomized to a double-blind, 8-week period (Period II) during which they received once-daily doses of placebo + OM 20 mg, AML 5 mg + OM 20 mg, or AML 10 mg + OM 20 mg. Patients controlled on olmesartan 20mg at the end of week 8 were discontinued from the study and not enrolled into the double-blind period.

The BP requirements for entering the open-label monotherapy treatment period included a mean sitting BP of  $\geq$ 160/100 mm Hg, a mean 24-hour DBP of  $\geq$ 84 mmHg, and at least 30% of daytime DBP readings  $>$ 90 mm Hg. Patients treated with either OM 20 mg or OM 40 mg at the beginning of the trial had to have a mean sitting BP of  $\geq$ 140/90 mm Hg, a mean 24-hour DBP of  $\geq$ 80 mm Hg, and at least 30% of daytime DBP readings  $>$ 85 mm Hg.

To enter the double-blind treatment period, patients needed to be non-responders to OM 20 mg. A non-responder was defined as mean trough sitting DBP  $\geq$ 90 mm Hg; mean trough sitting SBP  $\geq$ 140 mm Hg; and mean 24-hour DBP  $\geq$ 80 mm Hg with at least 30% of daytime DBP readings  $>$ 85 mm Hg. In addition to the BP requirements, patients should have met all other entry qualifications based on medical history, physical examination, electrocardiogram (ECG), and laboratory tests.

The Intent-to-Treat population (ITT) included all randomized patients who received at least 1 dose of double-blind study treatment and had at least 1 post-baseline efficacy assessment.

Sphygmomanometers were used for conventional BP measurements throughout the trial. After a 10-minute rest period, 3 separate sitting BP measurements were taken at least 1 minute apart. In addition, 24-hour ABPM was performed 3 times during the study (1 day prior to Visits 2, 4, and 6).

Safety assessments included adverse events, clinical laboratory measurements (hematology, biochemistry, and urinalysis), vital signs, physical examinations, and 12-lead ECG assessments.

Subgroup analyses were performed on the following subgroups: age (< 65 years,  $\geq$  65 years,  $\geq$  65 and < 75 years, and  $\geq$  75 years), gender (male, female), and hypertension severity (moderate or severe).

**Efficacy Results:** Of the 1,519 patients screened, 722 patients entered the open-label treatment period. There were 184 patients who discontinued from Period I, of which 148 patients were not eligible to continue to the double-blind treatment period. Therefore, a total of 538 patients were randomized in the double-blind treatment period. Of the 538 patients, 263 (48.9%) were male and all were Caucasian. The mean age was 56.8 years with 145 (27%) patients  $\geq$  65 years of age. A total of 43 (8.0%) patients had diabetes. Also, 68.6% of patients were classified with a BP of moderate severity and 26.2% were classified as severe. The treatment groups were similar with respect to BP

AZOR™ (amlodipine and olmesartan medoxomil)  
Formulary Dossier

at the start of the open-label treatment period. For patients randomized in the double-blind treatment period, mean sitting BP was 170.9/104.3 at the beginning of the monotherapy period.

The primary efficacy analysis demonstrated that 8 weeks of double-blind treatment with the combination of AML + OM (AML 5 mg + OM 20 mg and AML 10 mg + OM 20 mg) reduced mean sitting DBP to a significantly greater extent than treatment with placebo + OM 20. Table 22 presents the results for mean change and adjusted mean change in SeDBP from baseline (week 8) to week 16 (end of double-blind treatment) with LOCF for the ITT population. Treatment with AML + OM combination therapy resulted in statistically significant reductions in adjusted mean sitting DBP when compared with placebo + OM 20 mg therapy.

**Table 22: Mean Change in SeDBP (mm Hg) from Baseline (Week 8) to Week 16 with LOCF – ITT population**

Week 16 LOCF Analysis Variable	Placebo/OM20 (N = 179)	AML5/OM20 (N = 182)	AML10/OM20 (N = 177)
N <sup>[1]</sup>	179	182	177
Baseline mean (SD) <sup>[2]</sup>	97.2 (4.89)	97.5 (4.34)	97.1 (4.22)
Week 16 LOCF mean (SD) <sup>[3]</sup>	89.4 (8.54)	86.9 (7.39)	86.0 (7.59)
Mean change (SD)	-7.8 (7.86)	-10.6 (7.20)	-11.1 (8.01)
Adjusted mean change (SE) <sup>[4]</sup>	-7.6 (0.55)	-10.4 (0.55)	-10.9 (0.56)
Treatment comparison with Placebo/OM20			
Adjusted mean change (SE) <sup>[4]</sup>		-2.7 (0.75)	-3.2 (0.76)
95% confidence interval <sup>[4]</sup>		-4.4, -1.1	-4.9, -1.5
P-value <sup>[4]</sup>		0.0006	<0.0001

1. N = the number of patients with values at both time points.  
2. Baseline = Week 8.  
3. Week 16 LOCF was defined as the last available measurement during the double-blind treatment period.  
4. Statistics were based on an Analysis of Covariance model, including treatment, pooled centre, and baseline value as a covariate. All comparisons are with Placebo/OM20 using Dunnett's test to adjust for multiple testing. Adjusted means (least squares mean) and corresponding standard errors were derived from the ANCOVA model.  
AML = amlodipine; LOCF = last observation carried forward; OM = olmesartan medoxomil; SD = standard deviation; SE = standard error.

Similar results, as shown in Table 23, were observed for adjusted mean sitting SBP, and 24-hour BP by ABPM. For mean sitting SBP, the adjusted mean change from baseline (week 8) to week 16 with LOCF was -10.2 mm Hg for the placebo + OM 20 mg treatment group, -16.1 mm Hg for the AML 5 mg + OM 20 mg treatment group, and -16.7 mm Hg for the AML 10 mg + OM 20 mg treatment group. Treatment with AML + OM combination therapy resulted in statistically significant reductions in adjusted mean sitting SBP from baseline (week 8) to week 16 with LOCF when compared with placebo + OM 20 mg therapy: -5.8 mm Hg for AML 5 mg + OM 20 mg (p<0.0001) and -6.4 mm Hg for AML 10 mg + OM 20 mg (p<0.0001).

AZOR™ (amlodipine and olmesartan medoxomil)  
Formulary Dossier

**Table 23: Mean Change in SeSBP (mm Hg) from Baseline (Week 8) to Week 16 with LOCF – ITT population**

Week 16 LOCF Analysis Variable	Placebo/OM20 (N = 179)	AML5/OM20 (N = 182)	AML10/OM20 (N = 177)
N <sup>[1]</sup>	179	182	177
Baseline mean (SD) <sup>[2]</sup>	155.9 (10.26)	155.8 (10.12)	154.9 (9.79)
Week 16 LOCF mean (SD) <sup>[3]</sup>	145.3 (13.76)	139.5 (12.38)	138.4 (12.02)
Mean change (SD)	-10.6 (12.89)	-16.2 (10.66)	-16.5 (12.93)
Adjusted mean change (SE) <sup>[4]</sup>	-10.2 (0.86)	-16.1 (0.86)	-16.7 (0.87)
Treatment comparison with OM20/Placebo			
Adjusted mean change (SE) <sup>[4]</sup>		-5.8 (1.18)	-6.4 (1.18)
95% confidence interval <sup>[4]</sup>		-8.4, -3.2	-9.1, -3.8
P-value <sup>[4]</sup>		<0.0001	<0.0001

1. N = the number of patients with values at baseline and the given timepoint.  
2. Baseline = Week 8.  
3. Week 16 LOCF was defined as the Week 16 measurement or the last available measurement during the double-blind treatment period if the Week 16 value was missing.  
4. Statistics are based on an Analysis of Covariance model, including treatment, pooled centre, and baseline value as a covariate. All comparisons are with OM20/Placebo using Dunnett's test to adjust for multiple testing. Adjusted means (least squares mean) and corresponding standard errors were derived from the ANCOVA model.  
AML = amlodipine; LOCF = last observation carried forward; OM = olmesartan medoxomil; SD = standard deviation; SE = standard error.

The adjusted mean change from baseline (week 8) to week 16 in 24-hour mean ambulatory DBP was -4.5 mmHg for the placebo + OM 20 mg treatment group, -7.3 mmHg for the AML 5 mg + OM 20 mg treatment group, and -8.4 mmHg for the AML 10 mg + OM 20 mg treatment group. Treatment with AML + OM combination therapy resulted in statistically significant reductions in 24-hour adjusted mean ambulatory DBP from baseline (week 8) to week 16 LOCF when compared with placebo + OM 20 mg therapy: -2.8 mmHg for AML 5 mg + OM 20 mg (p=0.0031) and -3.9 mmHg for AML 10 mg + OM 20 mg (p<0.0001).

The adjusted mean change from baseline (week 8) to week 16 in 24-hour mean ambulatory SBP was -6.5 mmHg for the placebo + OM 20 mg treatment group, -11.4 mm Hg for the AML 5 mg + OM 20 mg treatment group, and -12.4 mm Hg for the AML 10 mg + OM 20 mg treatment group. Treatment with AML + OM combination therapy resulted in statistically significant reductions in 24-hour adjusted mean ambulatory SBP when compared with placebo + OM 20 mg therapy: -4.9 mm Hg for AML 5 mg + OM 20 mg (p=0.0020) and -5.8 mm Hg for AML 10 mg + OM 20 mg (p=0.0003).

Results were similar for mean changes in daytime mean DBP and SBP and nighttime mean DBP and SBP.

The adjusted mean change in SeDBP at Week 12 for patients treated with AML 10 mg + OM 20 mg combination therapy (-2.8 mm Hg) was statistically significant when compared to placebo + OM 20 mg therapy (p=0.0002). The adjusted mean change in sitting DBP at Week 12 for patients treated with AML 5 mg + OM 20 mg was -0.9 mm Hg when compared to placebo + OM 20 mg (p=0.3703). For SeSBP, the adjusted mean change for both the AML 5 mg + OM 20 mg treatment group (-2.9 mm Hg; p=0.0220) and the AML 10 mg + OM 20 mg treatment group (-6.3 mm Hg; p<0.0001) were statistically significant in comparison to the placebo + OM 20 mg. The time course of BP reductions demonstrated that in the group of patients that received AML 10 mg + OM 20 mg, earlier reductions in mean sitting DBP and SBP were achieved compared to patients that received AML 5 mg + OM 20 mg. At the week 12 visit, the difference in BP reduction between the placebo + OM 20 mg treatment group and the AML + OM combination treatment groups was greater in the AML 10 mg + OM 20 mg treatment group, compared to the AML 5 mg + OM 20 mg treatment group. However, with time, the differences between the 2 combination regimens narrowed and by week 16, the reductions in BP for both AML + OM combination treatment regimens were very similar.

The greater reductions in BP observed with AML and OM combination treatment translated into significantly more patients achieving pre-defined BP goals in both AML + OM combination treatment groups compared to the placebo + OM 20 mg treatment group. Compared to patients treated with placebo + OM 20 mg (28.5% achieving goal), the percentage of patients achieving BP goal at week 16 with LOCF was significantly higher in the AML 5 mg + OM 20 mg treatment group (44.5%; p=0.0011) and in the AML 10 mg + OM 20 mg treatment group (45.8%; p=0.0004).



AZOR™ (amlodipine and olmesartan medoxomil)  
Formulary Dossier

In the subgroup analyses, the efficacy of the AML + OM combination treatment regimens compared to placebo + OM 20 mg was similar for all age groups, for both males and females, and for both categories of hypertension severity.

**Safety Results:**

There were no new safety issues identified during the course of this study with either placebo + OM 20 mg, AML 5 mg + OM 20 mg, or AML 10 mg + OM 20 mg.

A total of 38 (21.2%) patients on placebo + OM 20 mg therapy experienced a treatment-emergent adverse event (TEAE) during Period II; 16 (8.9%) of these patients were considered to have had a drug-related TEAE. Thirty-two (17.6%) patients on AML 5 mg + OM 20 mg therapy experienced a TEAE; 14 (7.7%) of these patients were considered to have had a drug-related TEAE. A total of 35 (19.8%) patients on AML 10 mg + OM 20 mg therapy experienced a TEAE; 20 (11.3%) of these patients were considered to have a drug-related TEAE by the Investigator. Across the 3 treatment groups, most TEAEs and drug-related TEAEs were considered mild in severity. The differences in the incidence of TEAEs or drug-related TEAEs were not considered clinically meaningful when comparing the AML + OM combination regimens to the placebo + OM 20 mg treatment group.

There did not appear to be any meaningful differences in the incidence of adverse events in the AML + OM combination regimens compared to the placebo + OM 20 mg treatment group. Outside of peripheral edema, which had a slightly higher incidence in the AML 10 mg + OM 20 mg treatment group, there were no clinically meaningful patterns of TEAE incidence that signified that there might be a safety issue in a particular treatment group.

Overall, the most commonly reported drug-related TEAEs in the placebo + OM 20 mg treatment group were headache (1.7%), dizziness (1.1%), and hyperkalemia (1.1%). The incidence of drug-related treatment-emergent peripheral edema in the OM 20 mg + placebo group was 0.6%. The most commonly reported drug-related TEAEs in the AML 5 mg + OM 20 mg treatment group were headache (3.3%), peripheral edema (1.1%), and hyperkalemia (1.1%). The most commonly reported drug-related TEAEs in the AML 10 mg + OM 20 mg treatment group were headache (2.3%), peripheral edema (2.3%), increased blood potassium (1.1%), and increased gamma-glutamyltransferase ( $\gamma$ -GT) (1.1%).

During Period II, 5 (0.9%) patients discontinued due to an adverse event: 2 patients from the placebo + OM 20 mg treatment group, of which 1 (pain in joints) was considered by the Investigator to be related to study medication; 1 patient from the AML 5 mg + OM 20 mg treatment group (dizziness) which was considered to be related to study medication by the Investigator; and 2 patients from the AML 10 mg + OM 20 mg treatment group; both of whom were considered by the Investigator to have had an event (peripheral edema) related to study medication.

There were no laboratory measurements that signified a safety concern. There were no clinically meaningful changes in potassium levels or in renal function in any of the 3 treatment groups. In the AML + OM combination regimens, there were similar increases in platelet counts; a decrease in platelet counts occurred in the placebo + OM 20 mg treatment group. These increases were not considered clinically meaningful. Furthermore, there were no clinically meaningful changes in heart rates, ECGs, or physical examinations when the combinations of AML and OM were compared to treatment with placebo + OM 20 mg.

AZOR™ (amlodipine and olmesartan medoxomil)  
Formulary Dossier

**Add-on Study of Olmesartan Medoxomil in Patients with Moderate to Severe Hypertension not Achieving Target Blood Pressure on Amlodipine 5 mg Alone (Phase III, Europe)**

**Objectives:** The primary objective was to demonstrate the additional antihypertensive efficacy in lowering trough sitting diastolic blood pressure (DBP) gained by adding olmesartan medoxomil (OM) 10 mg, 20 mg, or 40 mg to the treatment regimen in patients with moderate to severe hypertension not adequately controlled on amlodipine (AML) 5 mg alone as assessed by conventional blood pressure (BP) measurements after 8 weeks of double-blind treatment. The primary efficacy endpoint was the mean change from baseline to the end of the double-blind treatment period, using last observation carried forward (LOCF) in trough sitting DBP.

The secondary objectives of this study consisted of an evaluation of the trough SeSBP from baseline (week 8) to week 12 and 16, from week 16 to week 20 and 24, and at week 28 and 34 in comparison to AML 5 mg. Additionally, SeDBP was evaluated from baseline to week 12, from week 16 to week 20 and 24, and at week 28 and 34. Ambulatory blood pressure monitoring (APBM) was utilized to evaluate antihypertensive efficacy (daytime, nighttime, and 24 hour) in DBP and SBP after 8 and 16 weeks of the double-blind treatment, and blood pressure goal (defined as BP <140/90 mmHg, <130/80 mmHg for diabetic patients) was evaluated during Period II, Period III, and Period IV (discussed below). The effect of titration to various dose combinations of AML and OM on DBP and SBP was evaluated. Safety and tolerability were assessed for the combinations of OM and AML versus monotherapy with AML 5 mg. Long-term safety and efficacy of the various combinations were also evaluated.

**Methodology:** This was a 52-week, randomized, parallel-group, multi-center, multi-national study consisting of 4 periods with patients who had hypertension (SBP  $\geq$  160 mm Hg and DBP  $\geq$  100 mm Hg) or were not controlled on amlodipine monotherapy (BP  $\geq$  140/90 mm Hg). For 8 weeks (Period I), patients received open-label once-daily doses of amlodipine 5 mg. Non-responders from Period I were then randomized to a double-blind, 8-week period (Period II) during which they received once-daily AML 5 mg + placebo, AML 5mg + OM 10mg, AML 5 mg + OM 20 mg, or AML 5 mg + OM 40 mg. A non-responder was defined as a patient with a mean trough sitting SBP and DBP  $\geq$  140/90 mmHg; and a mean 24-hour DBP  $\geq$  80 mmHg with at least 30% of daytime DBP readings  $>$ 85 mm Hg. Patients who responded adequately to AML 5mg monotherapy during Period I were discontinued from the study. At week 16, patients with uncontrolled blood pressure (BP  $\geq$  140/90 mmHg) were titrated to higher doses for an additional double-blind 8-week period (Period III). Patients who had been randomized to combination therapy with AML 5mg + OM 10mg, AML 5 mg + OM 20 mg, and AML 5 mg + OM 40 mg had their doses titrated to AML 5mg + OM 20 mg, AML 5 mg + OM 40 mg, or AML 10 mg + OM 40 mg, respectively. Patients randomized to AML 5mg + placebo had their dose titrated to AML 5 mg + OM 20 mg. Patients controlled from Period II, continued on their assigned treatment regimen during Period III. At week 24, patients continued into a 28-week, open-label extension study (Period IV) during which they all received AML 5 mg + OM 40 mg once-daily. If blood pressure was not controlled (BP  $\geq$  140/90 mmHg), patients were titrated to AML 10 mg + OM 40 mg; if blood pressure was still not controlled, hydrochlorothiazide 12.5 mg or 25 mg were added.

Sphygmomanometers were used for conventional BP measurements throughout the trial. After a 10-minute rest period, 3 separate sitting BP measurements were taken at least 1 minute apart. In addition, 24-hour ABPM was performed 4 times during the study (1 day prior to the start of Periods I, II, III, and IV)

The Full Analysis Set 1 included all randomized patients who received at least 1 dose of double-blind study medication and had a baseline sitting DBP measurement and at least 1 post-randomization sitting DBP measurement during Period II. This analysis population comprised 746 of the 755 randomized patients in Period II. The Full Analysis Set 2 included all randomized patients who received at least 1 dose of double-blind study medication during Period III and had a sitting DBP measurement at week 16 and at least 1 sitting DBP measurement during Period III (week 20 and/or week 24).

Safety assessments included treatment-emergent adverse events (TEAEs), clinical laboratory test results, vital signs, physical examinations, and 12-lead ECG assessments.

**Results:** The results provided include data from Period I, II, and III and the interim results from Period IV based on all data through week 34 of the study.

In total, 1017 patients entered Period I and received AML 5 mg monotherapy. Of the 1017 patients, 755 (74.2%) completed Period I and 262 (25.8%) discontinued during Period I, or were withdrawn at the end of Period I because they were responders. A total of 755 patients were randomized into Period II. Of the 755 randomized patients, 49 (6.5%) discontinued and 706 (93.5%) completed Period II. Subsequently, 706

AZOR™ (amlodipine and olmesartan medoxomil)  
Formulary Dossier

patients entered Period III, the double-blind up-titration period, of which 402 patients continued on their randomized treatment regimen from Period II, and 304 patients whose BP was not adequately controlled at the end of Period II had their doses titrated. For Period IV, 692 patients initially received AML 5 mg + OM 40 mg. As of week 34, 550 patients were on AML 5 mg + OM 40 mg, 120 patients were on AML 10 mg + OM 40 mg, 6 patients were on OM 40 mg + AML 10 mg + HCTZ 12.5 mg, and 1 patient was on OM 40 mg + AML 10 mg + HCTZ 25 mg.

The treatment groups for Period II were comparable with respect to demographic characteristics. Of the 755 patients, 61.1% were male, 99.7% were Caucasian, and the mean age was 55.8 years. Mean baseline BP was 164/102 mm Hg at the start of the study.

Table 24 presents the results for the primary efficacy parameter, the mean change in trough sitting DBP from baseline (Week 8) to Week 16 with LOCF (Period II), for the Full Analysis Set 1. Compared with placebo + AML 5 mg, treatment with OM + AML resulted in statistically significant reductions in sitting DBP.

**Table 24: Mean Change in Sitting Diastolic Blood Pressure – Period II**

Time Point Statistic	AML5/Placebo (N = 184)	AML5/OM 10 (N = 189)	AML5/OM20 (N = 187)	AML5/OM40 (N = 186)
<b>Week 16 LOCF</b>				
N [1]	184	189	187	186
Baseline mean (SD) [2]	97.2 (5.03)	96.9 (5.37)	96.9 (5.10)	97.0 (4.96)
Week 16 LOCF mean (SD) [3]	91.5 (8.39)	89.5 (7.58)	87.6 (8.63)	87.5 (7.22)
Mean change (SD)	-5.7 (7.66)	-7.4 (7.14)	-9.3 (7.74)	-9.5 (6.64)
Adjusted mean change (SE) [4]	-5.7 (0.53)	-7.7 (0.52)	-9.5 (0.52)	-9.6 (0.52)
Treatment comparison with AML5/Placebo				
Adjusted mean change (SE) [4]		-2.0 (0.73)	-3.7 (0.73)	-3.8 (0.73)
Adjusted 95% confidence interval [4]		(-3.7 , -0.2)	(-5.4 , -2.0)	(-5.5 , -2.1)
Adjusted p-value [4]		0.0207	<0.0001	<0.0001
1. N = number of patients with values at both baseline and the specified time point. 2. Baseline was the Visit 4 (Week 8) measurement. 3. Week 16 with LOCF was the Week 16 measurement, or the last available measurement during Period II if the Week 16 value was missing. 4. Statistics are from an Analysis of Covariance (ANCOVA) model with treatment and centre as effects and baseline as a covariate. Adjusted p-value is from Dunnett's test. AML = amlodipine; LOCF = last observation carried forward; OM = olmesartan medoxomil; SD = standard deviation; SE = standard error.				

Treatment with AML + OM (all evaluated dose regimens) demonstrated statistically significantly larger mean reductions in sitting DBP than placebo + AML 5 mg treatment at both the Week 12 and Week 16 time points. The DBP-lowering effect of AML 5 mg + OM 40 mg treatment was realized earlier than that of AML 5 mg + OM 20 mg treatment. At Week 16, the 2 highest dose regimens (AML 5 mg + OM 20 mg and AML 5 mg + OM 40 mg) demonstrated numerically larger mean reductions in sitting DBP than AML 5 mg + OM 10 mg.

Table 25 presents the results for the mean change in trough sitting SBP from baseline (Week 8) to Week 16 with LOCF (Period II) for the Full Analysis Set 1. Compared with AML 5 mg + placebo, treatment with AML + OM resulted in statistically significant reductions in sitting SBP.

AZOR™ (amlodipine and olmesartan medoxomil)  
Formulary Dossier

**Table 25: Mean Change in Sitting Systolic Blood Pressure – Period II**

Time Point Statistic	AML5/Placebo (N = 184)	AML5/OM10 (N = 189)	AML5/OM20 (N = 187)	AML5/OM40 (N = 186)
<b>Week 16 LOCF</b>				
N [1]	184	189	187	186
Baseline mean (SD) [2]	155.2 (11.53)	154.6 (10.29)	154.0 (11.32)	153.7 (10.21)
Week 16 LOCF mean (SD) [3]	145.3 (14.26)	141.5 (12.84)	138.7 (14.45)	137.0 (13.23)
Mean change (SD)	-9.9 (12.43)	-13.1 (11.64)	-15.3 (13.32)	-16.7 (12.00)
Adjusted mean change (SE) [4]	-9.7 (0.88)	-13.2 (0.87)	-15.4 (0.87)	-16.8 (0.87)
Treatment comparison with AML5/Placebo				
Adjusted mean change (SE) [4]		-3.5 (1.21)	-5.8 (1.22)	-7.1 (1.22)
Adjusted 95% confidence interval [4]		(-6.4 , -0.7)	(-8.6 , -2.9)	(-10.0 , -4.3)
Adjusted p-value [4]		0.0103	<0.0001	<0.0001
1. N = number of patients with values at both baseline and the specified time point. 2. Baseline was the Visit 4 (Week 8) measurement. 3. Week 16 with LOCF was the Week 16 measurement, or the last available measurement during Period II if the Week 16 value was missing. 4. Statistics are from an Analysis of Covariance (ANCOVA) model with treatment and centre as effects and baseline as a covariate. Adjusted p-value is from Dunnett's test. AML = amlodipine; LOCF = last observation carried forward; OM = olmesartan medoxomil; SD = standard deviation; SE = standard error.				

Table 26 summarizes the percentage of patients in each treatment group who reached BP goal during Period II for the Full Analysis Set 1. Compared with AML 5 mg + placebo, treatment with AML + OM resulted in a statistically significantly higher proportion of patients who reached BP goal at Week 16 with LOCF.

**Table 26: Number (%) of Patients Who Reached Blood Pressure Goal – Period II**

Time Point BP Goal Achieved	AML5/Placebo (N = 184) n (%)	AML5/OM10 (N = 189) n (%)	AML5/OM20 (N = 187) n (%)	AML5/OM40 (N = 186) n (%)
<b>Week 16 LOCF</b>				
Yes	55 (29.9)	74 (39.2)	100 (53.5)	94 (50.5)
P-value [1]		0.0286	<0.0001	<0.0001
1. The Cochran-Mantel-Haenszel test, stratified by centre, was used to compare the percentages for combination therapies with placebo + AML 5 mg therapy. AML = amlodipine; LOCF = last observation carried forward; OM = olmesartan medoxomil.				

The results for mean changes in ambulatory BP during Period II support the results for mean changes in sitting BP during Period II.

Table 27 presents the results for mean changes in trough sitting DBP and SBP during Period III for the Full Analysis Set 2. For patients who remained on their randomized treatment regimen during Period III, mean changes in sitting DBP and SBP from Week 16 to Week 24 with LOCF were not meaningful. Titration resulted in further mean reductions in sitting DBP and SBP during Period III.

AZOR™ (amlodipine and olmesartan medoxomil)  
Formulary Dossier

**Table 27: Mean Changes in Sitting Blood Pressure – Period III**

Treatment	N [1]	Mean Change (SD) in DBP	Mean Change (SD) in SBP	n (%) to BP goal
AML5/Placebo	68	0.7 (6.75)	1.0 (10.11)	40 (58.8)
AML5/Placebo to AML5/OM20	107	-8.2 (6.55)	-12.6 (11.47)	41 (38.3)
AML5/OM10	97	-0.7 (5.99)	-0.9 (9.39)	72 (74.2)
AML5/OM10 to AML5/OM20	82	-5.6 (7.02)	-7.5 (10.42)	23 (28.0)
AML5/OM20	118	-0.2 (6.76)	-0.9 (9.26)	89 (75.4)
AML5/OM20 to AML5/OM40	58	-6.2 (7.47)	-10.6 (12.76)	21 (36.2)
AML5/OM40	117	-0.6 (6.37)	-0.4 (9.39)	86 (72.9)
AML5/OM40 to AML10/OM 40	57	-8.2 (7.34)	-12.3 (11.12)	27 (47.4)

N = number of patients with values at both Week 16 and Week 24 with LOCF.

AML = amlodipine; BP = blood pressure; DBP = diastolic blood pressure; LOCF = last observation carried forward;

OM = olmesartan medoxomil; SBP = systolic blood pressure; SD = standard deviation.

For patients who remained on their randomized treatment regimen during Period III, the proportion who reached BP goal at Week 24 with LOCF was higher with AML + OM treatment than with AML 5 mg + placebo treatment. For patients whose dose regimen was titrated, successively higher proportions reached BP goal with each increase in dose combination of AML + OM. The results for mean changes in ambulatory BP during Period III support the results for mean changes in sitting BP during Period III.

In total, 692 patients entered the open-label Period IV and initially received AML 5 mg + OM 40 mg. For the 563 patients on AML 5 mg + OM 40 mg at Week 34 or Early Termination, the mean sitting DBP was 83.6 mm Hg and the mean sitting SBP was 132.2 mm Hg. For the 121 patients on AML 10 mg + OM 40 mg at Week 34 or Early Termination, the mean sitting DBP was 90.3 mmHg and the mean sitting SBP was 143.0 mmHg. For the 6 patients on AML 10 mg + OM 40 mg + HCTZ 12.5 mg at Week 34 or Early Termination, the mean sitting DBP was 89.3 mm Hg and the mean sitting SBP was 147.6 mm Hg. For the 1 patient on AML 10 mg + OM 40 mg + HCTZ 25 mg at Week 34 or Early Termination, sitting DBP was 92.0 mm Hg and sitting SBP was 155.3 mm Hg.

Titration from AML 5 mg + OM 40 mg to AML 10 mg + OM 40 mg during Period IV resulted in a mean reduction in sitting DBP of 5.0 mm Hg and a mean reduction in sitting SBP of 8.7 mm Hg. Titration from AML 10 + OM 40 mg to AML 10 mg + OM 40 mg + HCTZ 12.5 mg resulted in a mean reduction in sitting DBP of 3.7 mm Hg and a mean reduction in sitting SBP of 3.1 mm Hg.

Of the 692 patients exposed to AML 5 mg + OM 40 mg during Period IV, 502 (72.5%) reached BP goal. Of the 127 patients exposed to AML 10 mg + OM 40 mg, 46 (36.2%) reached BP goal. Of the 6 patients exposed to AML 10 mg + OM 40 mg + HCTZ 12.5 mg, 2 (33.3%) reached BP goal. The 1 patient exposed to AML 10 mg + OM 40 mg + HCTZ 25 mg did not reach BP goal.

**Safety Results:** The incidence of TEAEs during double-blind treatment was similar for the treatment groups. During Period II, 175 (23.2%) patients had a TEAE: 42 (22.3%) in the AML 5 mg + placebo group, 48 (25.1%) in the AML 5 mg + OM 10 mg group, 39 (20.6%) in the AML 5 mg + OM 20 mg group, and 46 (24.6%) in the AML 5 mg + OM 40 mg group. Fifty-three (7.0%) patients had a TEAE that was considered by the investigator to be related to study treatment: 15 (8.0%) in the AML 5 mg + placebo group, 13 (6.8%) in the AML 5 mg + OM 10 mg group, 11 (5.8%) in the AML 5 mg + OM 20 mg group, and 14 (7.5%) in the AML 5 mg + OM 40 mg group. Most of the TEAEs were mild or moderate in severity. The distributions of TEAEs by maximum severity were similar for the treatment groups.

During Period II, the most frequently reported TEAEs in the AML 5 mg + placebo group were peripheral edema (2.1%) and vertigo (2.1%). The most frequently reported TEAEs in the AML 5 mg + OM 10 mg group were headache (3.1%) and dizziness (3.1%). The most frequently reported TEAEs in the AML 5 mg + OM 20 mg group were back pain (2.6%), dizziness (1.6%), and influenza (1.6%). The most frequently reported TEAEs in the AML 5 mg + OM 40 mg group were headache (3.7%) and dizziness (3.2%). During Period II, the most frequently reported drug-related TEAEs in the AML 5 mg + placebo group were peripheral edema (2.1%) and vertigo (1.1%). The most frequently reported drug-related TEAEs in the AML 5 mg + OM 10 mg group were headache (1.6%) and dizziness (1.6%). The most frequently reported drug-related TEAEs in the OM 20 mg + AML 5 mg group were peripheral edema (1.1%), dizziness (1.1%), and hypotension (1.1%). The most frequently reported drug-related TEAEs in the AML 5 mg + OM 40 mg group were headache

AZOR™ (amlodipine and olmesartan medoxomil)  
Formulary Dossier

(2.7%) and dizziness (1.6%). The incidence of drug-related treatment-emergent peripheral edema in AML 5 mg + OM 40 group was 1.1%.

No patients died during Period II. Four (0.5%) patients had a serious adverse event (SAE) during Period II, however none of the SAEs were considered by the investigator to be related to study treatment.

Thirteen (1.7%) patients discontinued from the study during Period II due to a drug-related TEAE: 1 (0.5%) patient in the AML 5 mg + placebo (malaise), 3 (1.6%) patients in the AML 5 mg + OM 10 mg group (1 patient with dizziness and headache, 1 patient with decreased blood pressure, and 1 patient with vomiting and lethargy), 4 (2.1%) patients in the AML 5 mg + OM 20 mg group (1 patient with hypotension, 1 patient with myalgia, 1 patient with increased blood pressure, and 1 patient with peripheral edema), and 5 (2.7%) patients in the AML 5 mg + OM 40 mg group (1 patient with vertigo, 1 patient with dizziness, 2 patients with hypotension, and 1 patient with cold sweat, asthenia, and dizziness).

During Period III, 117 (16.6%) patients had a TEAE. No meaningful differences among the treatment regimens in the incidence of TEAEs during Period III were observed. Thirty-two (4.5%) patients had a TEAE that was considered by the investigator to be related to study treatment. Most of the TEAEs were mild or moderate in severity. The distributions of TEAEs by maximum severity were similar for patients on the various treatment regimens.

Four (0.6%) patients had an SAE during Period III, however, none of the SAEs were considered by the investigator to be related to study treatment. Two (0.3%) patients discontinued from the study during Period III due to a TEAE: 1 (0.8%) patient on AML 5 mg + OM 20 mg and 1 (0.8%) patient on AML 5 mg + OM 40. The TEAEs that led to discontinuation for the 1 (0.8%) patient on AML 5 mg + OM 40 mg (fatigue, angina pectoris, dizziness, headache, and visual disturbance) were considered by the investigator to be related to study treatment.

During Period IV through Week 34, a total of 37 (5.3%) patients had a TEAE that was considered by the investigator to be related to study treatment: 32 (4.6%) patients on AML 5 mg + OM 40 mg and 5 (3.9%) patients on AML 10 mg + OM 40 mg. Most of the TEAEs were mild or moderate in severity. The distribution of TEAEs by maximum severity was similar for patients on the various treatment regimens.

No patients died during Period IV. Seven (1.0%) patients had an SAE during Period IV, however, none of the SAEs were considered by the investigator to be related to study treatment.

Six (0.9%) patients discontinued from the study during Period IV due to a TEAE: 5 (0.7%) patients on AML 5 mg + OM 40 mg and 1 (0.8%) patient on AML 10 mg + OM 40 mg. For 4 (0.6%) patients on AML 5 mg + OM 40 mg, the TEAE that led to discontinuation was considered by the investigator to be related to study treatment (3 patients with dizziness and 1 patient with hypotension).

Mean changes in laboratory parameters and shifts in laboratory abnormalities were not clinically meaningful for any of the treatment groups during Period II or patients on the various treatment regimens during Periods III and IV. No clinically meaningful differences in these parameters between AML + OM combination treatment and AML + placebo treatment were noted. In addition, mean changes in pulse rate and ECG parameters during the study were not clinically meaningful. No clinically meaningful differences in these parameters were observed among the various treatment regimens. No clinically meaningful physical examination findings were noted, as well.

## B. PUBLISHED AMLODIPINE/OLMESARTAN MEDOXOMIL ABSTRACTS

### American Society of Hypertension 2007

#### **A Randomized, Double-Blind, Placebo-Controlled Factorial Study Evaluating The Efficacy And Safety Of Co-Administration Of Amlodipine Besylate (AML) Plus Olmesartan Medoxomil (OM) Compared To Monotherapy In Patients (Pts) With Mild To Severe Hypertension (HTN)**

Steven G Chrysant, MD<sup>1</sup>; Michael Melino, PhD<sup>2</sup>; Sulekha Karki, BAMS<sup>2</sup>; James Lee, PhD<sup>2</sup>; Reinilde Heyrman, MD<sup>2</sup>

<sup>1</sup>Oklahoma Cardiovascular and Hypertension Center and the University of Oklahoma School of Medicine, Oklahoma City, OK; Daiichi Sankyo, Inc., Parsippany, NJ, USA<sup>2</sup>


Most pts will need  $\geq 2$  antihypertensive agents to attain a blood pressure (BP) goal of  $<140/90$  mmHg. Concomitant administration of the calcium channel blocker AML and the angiotensin receptor blocker OM has the potential to improve BP-lowering efficacy without additional safety concerns. We conducted a randomized, double-blind, placebo-controlled factorial study in 1940 pts with mild to severe HTN, defined as seated diastolic BP (SeDBP) between  $\geq 95$  mmHg and  $\leq 120$  mmHg, to determine if co-administration of AML 5-10 mg/day and OM 10-40 mg/day for 8 weeks has a significant benefit versus respective monotherapy components. Primary and secondary endpoints were mean change from baseline in SeDBP and seated systolic BP (SeSBP), respectively, at week (Wk) 8, with last observation carried forward if pts withdrew prior to Wk 8. Each combination had significantly greater reductions in SeDBP (Fig.) and SeSBP compared with both of its monotherapy components ( $P < 0.0001$  for all comparisons). The greatest observed mean reductions in seated SeSBP and SeDBP occurred with AML 10mg + OM 40mg ( $-30.1/-19.0$  mmHg vs  $-4.8/-3.1$  mmHg with placebo, and vs  $-19.7/-12.7$  mmHg with AML 10 mg). Adverse event (AE) incidence for the AML + OM combination treatment groups was 52.7% (511/970 pts); the AE incidence in the placebo group was 56.2% (91/162 pts). Most AEs were mild in intensity. The AE profile for each of the combinations was similar in nature to the monotherapy components. In conclusion, the combination of AML + OM produced greater reductions in BP compared with placebo and respective monotherapy. BP reductions were dose-related and all regimens were well tolerated.

AZOR™ (amlodipine and olmesartan medoxomil)  
Formulary Dossier

**Figure**

## Reduction in SeDBP with All Tested Doses

*Week 8 LOCF in Total Population*



Values are expressed as LS mean ± SE.  $P < 0.0001$  for all active treatment groups.

AML or amlodipine=amlodipine besylate; LOCF=last observation carried forward; OM or olmesartan=olmesartan medoxomil; SeDBP=sitting diastolic blood pressure. Baseline diastolic blood pressure (DBP) = 101.6 mm Hg for safety population.



AZOR™ (amlodipine and olmesartan medoxomil)  
Formulary Dossier

American Society of Hypertension 2007 & European Society of Hypertension 2007

**A Fixed-Dose Combination of Olmesartan Medoxomil (OM) and Amlodipine Besylate (AM) is Bioequivalent (BEQ) to Free Combination of the Agents**

Stephen Haworth, MD<sup>1</sup>; Mohinder Singh Bathala, PhD<sup>1</sup>; James Lee, PhD<sup>1</sup>; Mark Allison, MD<sup>2</sup>; Shashank Rohatagi, PhD<sup>1</sup>; Jean-Francois Marier, PhD<sup>3</sup>; Igor Rubets, PhD<sup>3</sup>; Daniel Salazar, PhD<sup>1</sup>

<sup>1</sup>Daiichi Sankyo Inc., Parsippany, NJ, USA; <sup>2</sup>MDS Pharma Services, Phoenix, AZ, USA; <sup>3</sup>MDS Pharma Services, Montreal, Quebec, Canada

The combination of an angiotensin receptor blocker and calcium channel blocker may have additive BP-lowering effects vs either monotherapy due to complementary mechanisms of action. A fixed-dose combination of OM + AM offers the potential for increased BP-lowering, convenient dosing, and improved compliance. The pharmacokinetics/bioavailability of a novel fixed-dose combination were assessed in a phase I study; 28 healthy subjects enrolled in a single-dose, open-label, 2-way crossover study were randomized in each dosing period to receive under fasting conditions the test treatment, a fixed-dose combination of OM/AM 40/10mg, or the reference treatment of free combination of OM 40mg and AM 10mg. Results showed the fixed-dose and free combinations were BEQ as the criteria of the 90% CI for the area under the concentration-time curve from time 0 to time t ( $AUC_{0-t}$ ),  $AUC$  from time 0 to infinity ( $AUC_{0-\infty}$ ) and  $C_{max}$  ratios (Test/Reference) for OM and AM were within 80.0 to 125.0% limit. [table1] All doses were well tolerated. Mean terminal elimination half-life ( $t_{1/2}$ ) was similar after the test and reference treatments for OM (11h and 12h respectively) and AM (54h and 52h respectively). Of the 30 mild treatment-emergent AEs in 11 subjects (most frequently headache), 11 were possibly drug-related occurring in 4 subjects in the test and 2 subjects in the reference treatment groups. No clinically significant trends in abnormal laboratory findings occurred. The fixed-dose combination of OM/AM 40/10mg is BEQ to OM 40mg and AM 10mg coadministered separately.

**Table**

Parameter	Geometric LSM		LSM ratio (T/R)	90% CI
	Test [T] (n=26)	Reference [R] (n=27)		
<b>OM</b>				
AUC <sub>0-t</sub> (ng·h/mL)	5374.2	5418.6	99.2	93.4, 105.3
AUC <sub>0-∞</sub> (ng·h/mL)	5407.5	5468.3	98.9	93.2, 104.9
C <sub>max</sub> (ng/mL)	833.3	810.3	102.9	94.8, 111.6
<b>AM</b>				
AUC <sub>0-t</sub> (ng·h/mL)	424.8	410.9	103.4	100.1, 106.8
AUC <sub>0-∞</sub> (ng·h/mL)	505.3	482.2	104.8	100.8, 109.0
C <sub>max</sub> (ng/mL)	7.6	7.4	103.9	99.9, 108.2
LSM=least squares mean; t=time to last measurable concentration.				

AZOR™ (amlodipine and olmesartan medoxomil)  
Formulary Dossier

**American Society of Hypertension 2007 & European Society of Hypertension 2007**

**The Bioavailability (BAV) of a Fixed-Dose Combination of Olmesartan Medoxomil (OM) and Amlodipine Besylate (AM) is Unaffected by Food**

Stephen Haworth, MD<sup>1</sup>; Reinilde Heyrman, MD<sup>1</sup>; Mohinder Singh Bathala, PhD<sup>1</sup>; James Lee, PhD<sup>1</sup>; Robert Noveck, MD, PhD<sup>2</sup>; Shashank Rohatagi, PhD<sup>1</sup>; Jean-Francois Marier, PhD<sup>3</sup>; Igor Rubets, PhD<sup>3</sup>; Daniel Salazar, PhD<sup>1</sup>

<sup>1</sup>Daiichi Sankyo Inc., Parsippany, NJ, USA; <sup>2</sup>MDS Pharma Services, Neptune, NJ, USA <sup>3</sup>MDS Pharma Services, Montreal, Quebec, Canada

Most patients require  $\geq 2$  antihypertensive agents to achieve BP control. The combination of an angiotensin receptor blocker, e.g., OM, and a calcium channel blocker, e.g. AM, may provide increased BP-lowering efficacy compared with either monotherapy and a fixed-dose formulation of these agents simplifies dosing and may increase compliance. This study evaluated the effect of food on the BAV of an OM/AM fixed-dose combination using a randomized, open-label, 2-way crossover study in 28 healthy subjects who received a single fixed dose of OM/AM 40/10mg 30 min after a high-fat breakfast (treatment A) and after 10-hr fasting conditions (treatment B), with a 21-day washout period between treatments. Absence of a food effect was concluded if the 90% CIs for the treatment A:treatment B ratios for the geometric least squares mean of AUC time 0 to t (AUC<sub>0-t</sub>), AUC time 0 to infinity (AUC<sub>0-∞</sub>) and C<sub>max</sub> of AM and OM were 80-125.0%. All dosing regimens were well tolerated. Ratios of LSM of AUC<sub>0-t</sub>, AUC<sub>0-∞</sub> and C<sub>max</sub> for OM were 87%, 88%, 94%, with 90% CIs all within 80-125%. For AM, ratios of LSM of AUC<sub>0-t</sub>, AUC<sub>0-∞</sub> and C<sub>max</sub> were 103%, 103%, 99%, with 90% CIs all within 80-125%. The mean terminal elimination half-life for OM (14h) and AM (40h) was unaffected by food. Most treatment-emergent adverse events were of mild severity, occurring in 5 subjects. The only drug-related AE was headache (2 subjects). In conclusion, the BAV of OM and AM after a single fixed-dose combination tablet was equivalent either with food or under fasting conditions.

**Table**

Parameter	Geometric LSM		LSM ratio [A/B] (%)	90%CI (%)
	Treatment A - OM/AM 40/10mg (n=28) Fed	Treatment B - OM/AM 40/10mg (n=27) Fasting		
<b>OM</b>				
AUC <sub>0-t</sub> (ng·h/mL)	5259.6	6034.3	87	82.5, 92.1
AUC <sub>0-∞</sub> (ng·h/mL)	5366.5*	6111.7	88	83.0, 92.9
C <sub>max</sub> (ng/mL)	881.9	939.5	94	87.4, 100.8
<b>AM</b>				
AUC <sub>0-t</sub> (ng·h/mL)	307.0	299.2	103	99.6, 105.7
AUC <sub>0-∞</sub> (ng·h/mL)	334.3	326.1	103	99.2, 106.0
C <sub>max</sub> (ng/mL)	6.4	6.4	99	96.0, 102.7
LSM=least squares mean; * n=27				

**American Society of Hypertension 2007 & European Society of Hypertension 2007**

**Safety and Tolerability of the Combination of Olmesartan Medoxomil (OM) and Amlodipine Besylate (AM)**

Shashank Rohatagi, PhD<sup>1</sup>; Stephen Haworth, MD<sup>1</sup>; Reinilde Heyrman, MD<sup>1</sup>; James Lee, PhD<sup>1</sup>;  
Robert Noveck, MD<sup>2</sup>; Jean-Francois Marier, PhD<sup>3</sup>; Igor Rubets, PhD<sup>3</sup>; Daniel Salazar, PhD<sup>1</sup>

<sup>1</sup>Daiichi Sankyo, Inc., Parsippany, NJ; USA <sup>2</sup>MDS Pharma Services, Neptune, NJ, USA, <sup>3</sup>MDS Pharma Services, Montreal, Quebec, Canada

Many hypertensive patients need >1 antihypertensive agent; combining agents with different mechanisms of action may improve efficacy vs monotherapy. The pharmacokinetics of a novel fixed-dose combination of OM and AM were compared with free combination of these agents (OM + AM) or monotherapy in phase 1 studies in 200 healthy volunteers (VOLS). Safety/tolerability data for OM/AM single- or multiple-dose studies of randomized, open-label, crossover design are presented. [table1] Most treatment-emergent adverse events (TEAEs) were of mild severity; the most common was headache. In one large study (Study 112; n=60), the most frequent TEAEs were headache, dizziness, cough and pharyngolaryngeal pain, reported in 37%, 13%, 12% and 12% of VOLS. In 2 studies (111, 110), headache was the only possibly drug-related AE; headache and dizziness were the only possibly or probably drug-related AEs in Study 112. There were no severe AEs for the fixed or free dose combination of OM + AM. 1 VOL given a single dose of OM 40mg + AM 10mg discontinued due to a non-drug related AE (Study 111). In Study 109, a VOL discontinued due to a positive serum pregnancy test and later had a spontaneous abortion classified as a serious, unlikely drug-related AE. There were few clinically significant laboratory abnormalities (0–6 VOLS in individual studies) and only 1 treatment-related laboratory abnormality (elevated liver enzymes) in 1 VOL. Generally, there were no clinically significant ECG abnormalities. The fixed-dose combinations of OM/AM appear to be safe and well tolerated in healthy VOLS.

AZOR™ (amlodipine and olmesartan medoxomil)  
Formulary Dossier

**Table**

Regimen (mg)	n	TEAE, # VOLS	Drug related AE, # VOLS <sup>‡</sup>
<b>Study 101</b>			
OM 40+AM 10	24	8	5*
OM 40	23	10	7*
AM 10	23	14	7*
<b>Study 111</b>			
OM/AM 10/5	30	7	3*
OM/AM 40/10	29	3	1*
OM 10+AM 5	30	5	0*
OM 40+AM 10	29	3	0*
<b>Study 112</b>			
OM/AM 40/10	30	15	7 <sup>¶</sup>
OM/AM 20/5	29	12	5 <sup>¶</sup>
OM/AM 10/10	29	11	7 <sup>¶</sup>
OM/AM 40/5	30	9	5 <sup>¶</sup>
OM/AM 20/10	29	10	5 <sup>¶</sup>
OM/AM 10/5	30	12	5 <sup>¶</sup>
<b>Study 110</b>			
OM/AM 40/10	28	5	2*
<b>Study 109</b>			
OM/AM 40/10	27	6	4*
OM 40+AM 10	27	5	2*
*Possibly; <sup>¶</sup> Possibly or Probably			
<sup>‡</sup> Pts can appear in >1 group within a study			

**American Society of Hypertension 2007 & European Society of Hypertension 2007**

**Evaluation of Population (POP) Pharmacokinetics (PK) After Coadministration of Olmesartan Medoxomil (OM) and Amlodipine Besylate (AM)**

Shashank Rohatagi, PhD<sup>1</sup>; Timothy Carrothers, ScD<sup>2</sup>; Smita Kshirsagar, PhD<sup>2</sup>; Tatiana Khariton, PhD<sup>2</sup>; James Lee, PhD<sup>1</sup>; Daniel Salazar, PhD<sup>1</sup>

<sup>1</sup>Daiichi Sankyo, Inc., Parsippany, NJ; <sup>2</sup>Pharsight Corporation, Mountain View, CA

A novel fixed-dose combination of OM and AM may increase efficacy over monotherapy in patients (pts) with hypertension (HTN). The single formulation may also improve ease of administration and potentially compliance. Data from phase I studies in 170 healthy volunteers (HV) (studies CS8663-U101, 110, 111, 112) and from a subset (n≈546) of pts with HTN in a phase III trial (CS8663-U301) were used to make POP PK models and subsequent analyses of OM and AM. Both OM and AM were characterized by 1<sup>st</sup> order elimination/absorption and an absorption time lag in a 2 compartmental model for OM and a 1 compartmental model for AM. In the OM model, covariate analyses indicated decreased oral clearance (CL) for pts with HTN vs HV, female pts, pts with higher baseline serum creatinine (SeCr) vs lower SeCr or those with lower vs higher bodyweight (WT). For AM, CL decreased for pts with higher vs lower baseline ALT, older vs younger pts or pts with lower vs higher WT (Table). Gender-based differences were independent of WT and other covariates. Similarly, the association between age and AM exposure was due to an effect on AM CL that was independent of age-related changes in WT or other covariates. Combination therapy did not modify the effects of covariate status on CL of OM or AM. In conclusion, neither compound had a clinically significant impact on the CL of the other. The impact of covariates on the CL of OM and AM did not change between monotherapy and combination therapy.

AZOR™ (amlodipine and olmesartan medoxomil)  
Formulary Dossier

**Table**

	POP mean	
	Estimate	Coefficient of variation [100 * SE <sub>estimate</sub> /estimate] (%)
<b>OM</b>		
CL <sub>TYP</sub> (L/h)	5.9	4.4
V (L)	32.1	2.5
K <sub>a</sub> (per h)	2.0	5.1
ALAG1 (h)	0.374	1.5
CL <sub>HV</sub>	1.7	20
CL <sub>SEX</sub>	-0.878	34
CL <sub>WT</sub>	0.326	31
CL <sub>SeCr</sub>	-0.278	25
<b>AM</b>		
CL <sub>TYP</sub> (L/h)	22.9	1.7
V (L)	1530	2.0
K <sub>a</sub> (per h)	0.640	2.3
ALAG1 (h)	0.390	10
CL <sub>WT</sub>	0.207	36
CL <sub>AGE</sub>	-0.373	14
CL <sub>ALT</sub>	-0.138	32
ALAG1=absorption lag time; CL <sub>TYP</sub> =oral CL of typical pt; K <sub>a</sub> =absorption rate constant; V=volume of central distribution. Effects on CL are linear for discrete covariates, CL (HV)=CL <sub>TYP</sub> +CL <sub>HV</sub> , and multiplicative for continuous covariates, CL (WT of 90)=CL <sub>TYP</sub> * (90/WT <sub>median</sub> ) <sup>CL<sub>WT</sub></sup>		

AZOR™ (amlodipine and olmesartan medoxomil)  
Formulary Dossier

**American Society of Hypertension 2007 & European Society of Hypertension 2007**

**Olmesartan Medoxomil (OM) and Amlodipine Besylate (AM) Show Dose-Proportional Drug Exposure When Used in Combination in Healthy Volunteers (VOLS)**

Shashank Rohatagi, PhD<sup>1</sup>; Stephen Haworth, MD<sup>1</sup>; Daniel Salazar, PhD<sup>1</sup>; Mohinder Singh Bathala, PhD<sup>1</sup>; Igor Rubets, PhD<sup>3</sup>; Jean-Francois Marier, PhD<sup>3</sup>; Dennis Swearingen, MD<sup>2</sup>; James Lee, PhD<sup>1</sup>

<sup>1</sup>Daiichi Sankyo Inc., Parsippany, NJ USA; <sup>2</sup>MDS Pharma Services, Phoenix, AZ, USA; <sup>3</sup>MDS Pharma Services, Montreal, Quebec Canada

JNC 7 recommends combination drug therapy if BP is >20/10mmHg above goal. In accordance, a fixed-dose combination of the angiotensin receptor blocker OM and the calcium channel blocker AM is being evaluated for antihypertensive efficacy. This study assessed the dose-proportionality of OM/AM fixed-dose combinations in VOLS in 2 cohorts of a randomized, open-label, 3-way crossover design. VOLS received a single dose of oral OM/AM under fasting conditions as follows: 40/10mg (treatment A), 20/5mg (B) and 10/10mg (C) in cohort 1 (n=30) or 40/5mg (D), 20/10mg (E) and 10/5mg (F) in cohort 2 (n=30). All dosing regimens were well tolerated. Noncompartmental pharmacokinetics (PK) by dose level (data pooled across treatments) are presented. [table1] A dose-proportional increase in OM AUC<sub>0-t</sub> and AUC<sub>0-∞</sub> was observed as the 95% CI of the regression coefficient for the ln-transformed PK parameters were within 0.75-1.25. The C<sub>max</sub> for OM 10, 20, and 40mg doses increased in a slightly less than dose-proportional manner; considered to be not clinically significant. Dose-proportionality of AM was demonstrated as the 90% CI for dose-normalized parameters were within the 80–125% range. Mean terminal elimination half-life of OM (10 to 40mg) and AM (5 to 10mg) ranged from 14.0-15.1 h and 48.4-51.6 h respectively. OM 10, 20 or 40mg was well tolerated when given in fixed-dose combination with either AM 5 or 10mg. Most TEAEs were of mild severity and none considered treatment-related. There were no serious AEs. In conclusion, the exposure of OM and AM for the fixed-dose combinations increased in a dose-proportional manner.

**Table**

PK Parameters*	OM dose			AM dose		
	40mg [A and D] (n=59)	20mg [B and E] (n=54-57)	10mg [C and F] (n=58)		10mg [A,C,E] (n=87)	5mg [B,D,F] (n=87)
AUC <sub>0-t</sub> (ng·h/mL)	6006.4	3512.4	1885.0	AUC <sub>0-t</sub> (ng·h/mL)	385.8 <sup>†</sup>	172.5
AUC <sub>0-∞</sub> (ng·h/mL)	6096.2**	3573.8	1921.0	AUC <sub>0-∞</sub> (ng·h/mL)	455.0 <sup>†</sup>	200.3
C <sub>max</sub> (ng/mL)	928.2	574.9	337.2	C <sub>max</sub> (ng/mL)	7.7	3.6

\*Arithmetic means are presented; \*\*n=58; <sup>†</sup>n=86

AZOR™ (amlodipine and olmesartan medoxomil)  
Formulary Dossier

**American Society of Hypertension 2007 & European Society of Hypertension 2007**

**LOW AND HIGH DOSES OF A FIXED-DOSE COMBINATION OF OLMESARTAM MEDOXOMIL (OLM) AND AMLODIPINE BESYLATE (AML) ARE BIOEQUIVALENT TO FREE COMBINATION OF THE AGENTS**

Shashank Rohatagi, PhD<sup>1</sup>; Stephen Haworth, MD<sup>1</sup>; Daniel E. Salazar, PhD<sup>1</sup>; Reinilde Heyrman, MD<sup>1</sup>; Mohinder Singh Bathala, PhD<sup>1</sup>; James Lee, PhD<sup>1</sup>; Robert Noveck, MD<sup>2</sup>

<sup>1</sup>Daiichi Sankyo Pharma Development, Edison, NJ; <sup>2</sup>MDS Pharma, Neptune, NJ

Two or more antihypertensive agents from different pharmacologic classes may be needed to reach a target BP of <140/90 mmHg in a majority of patients. The use of a fixed-dose combination such as the angiotensin receptor blocker OLM and the calcium channel blocker AML provides increased BP-lowering efficacy with good tolerability and ease of administration. The bioequivalence under fasting conditions of a fixed-dose combination of OLM and AML in low (OLM/AML 10/5mg) and high (OLM/AML 40/10mg) doses to that of the drugs coadministered separately was investigated in 2 cohorts of a randomized, open-label, 2-way crossover single-dose study in healthy subjects (n=60). Noncompartmental analyses showed the bioavailability of both OLM and AML after administration of the fixed-dose combination (Test) was similar to that of the agents coadministered separately (Reference), at both the low and high doses (Table). Mean  $t_{1/2}$  for the low dose Test and Reference treatments were 14.3 and 13.6 hrs and for the high dose 15.6 and 17.3 hrs, respectively. Overall, treatment emergent adverse events (TEAEs) were mild; none were considered definitely or probably drug-related and the most common was headache. In conclusion, a single fixed-dose combination tablet of OLM/AML 10mg/5mg or 40mg/10mg was well tolerated and is bioequivalent to coadministration of single doses of OLM and AML under fasted conditions in healthy subjects.



AZOR™ (amlodipine and olmesartan medoxomil)  
Formulary Dossier

**Table**

	Geometric LSM ratio (%) [Test/Reference]	90%CI (%)
<b>Low dose (n=30)</b>		
<b>OM</b>		
AUC <sub>0-t</sub>	107.6	99.7, 116.1*
AUC <sub>0-∞</sub> <sup>†</sup>	107.4	99.4, 116.0*
C <sub>max</sub>	114.3	106.6, 122.5*
<b>AM</b>		
AUC <sub>0-t</sub>	101.6	99.1, 104.2*
AUC <sub>0-∞</sub>	101.6	99.0, 104.3*
C <sub>max</sub>	99.0	95.7, 102.5*
<b>High dose (n=29)</b>		
<b>OM</b>		
AUC <sub>0-t</sub>	112.1	103.3, 121.6*
AUC <sub>0-∞</sub> <sup>‡</sup>	113.5	104.7, 123.0*
C <sub>max</sub>	109.7	101.8, 118.3*
<b>AM</b>		
AUC <sub>0-t</sub>	101.6	97.3, 106.2*
AUC <sub>0-∞</sub>	101.2	96.6, 106.0*
C <sub>max</sub>	108.3	103.2, 113.6*
*Criteria for bioequivalence, i.e., 90%CI for the ratio of geometric LSM to be within 80.0 to 125.0%, were met		
LSM=least squares mean, <sup>†</sup> n=29, <sup>‡</sup> n=27		

AZOR™ (amlodipine and olmesartan medoxomil)  
Formulary Dossier

American Society of Hypertension 2007 & European Society of Hypertension 2007

**Lack of Pharmacokinetic (PK) Drug Interaction Between Olmesartan Medoxomil (OM) and Amlodipine Besylate (AM) During Coadministration**

Daniel Salazar, PhD<sup>1</sup>; James Lee, PhD<sup>1</sup>; Magdy Shenouda, MD<sup>2</sup>; Shashank Rohatagi, PhD<sup>1</sup>

<sup>1</sup>Daiichi Sankyo Inc., Parsippany, NJ, USA; <sup>2</sup>MDS Pharma Services, Neptune, NJ, USA

The majority of hypertensive patients require multiple drug regimens to achieve blood pressure (BP) goal. The angiotensin receptor blocker OM and the calcium channel blocker AM are each efficacious monotherapies and combining these agents adheres to JNC 7 guidelines for using combination therapies with complementary mechanisms of action. The PK interaction between these agents was investigated in an open-label, multiple-dose, 3-way crossover study in healthy subjects under fasting conditions. Subjects were assigned randomly to receive 1 of the following 10-day regimens on 3 occasions (each regimen separated by ≥21 days): treatment A - OM 40mg/d; treatment B - AM 10mg/d and treatment C - OM 40mg/d + AM 10mg/d. Noncompartmental, steady-state (day 10) PK parameters of OM following treatments A and C and of AM following treatments B and C are shown. [table1] All dosing regimens were well tolerated. There was no significant PK interaction between OM and AM. The 90% CIs for the ratio of geometric least squares mean for AUC<sub>0-τ</sub> and C<sub>max</sub> of OM and AM were within the 80.0-125.0% limit (combined treatment vs each monotherapy). Coadministration had no effect on the t<sub>1/2</sub> of either agent. The tolerability profile of combination therapy was similar to that of each monotherapy. There were no serious AEs or AE-related discontinuations. Most TEAEs were mild; the most frequent was headache (n=10/24). Drug-related AEs occurred in 5 subjects during combination therapy, and in 7 subjects with each monotherapy. In conclusion, coadministration of OM and AM did not affect the rate or extent of exposure of OM or AM and was well tolerated.

	OLM			AML		
PK parameters (mean)	Treatment A n=23	Treatment C n=24	C/A ratio of geometric LSM (%) (90%CI)	Treatment B n=23	Treatment C n=24	C/B ratio of geometric LSM (%) (90%CI)
AUC <sub>τ</sub> (ng·h/mL)	6793.9	6890.9	101.1 (93.5, 109.4)	359.2	388.7	107.7 (100.1, 115.9)
C <sub>max</sub> (ng/mL)	1083.8	1038.1	95.2 (87.2, 103.9)	19.8	20.1	100.7 (91.3, 111.1)
t <sub>max</sub> (h) <sup>1</sup>	1.5 (1.00, 2.52)	2.0 (1.02, 2.98)		8.00 (5.00, 14.00)	8.0 (0.0, 16.1)	
t <sub>1/2</sub> (h)	13.7	13.48		51.2	50.6	

<sup>1</sup>Median (min, max) values presented

AZOR™ (amlodipine and olmesartan medoxomil)  
Formulary Dossier

AZOR™ (amlodipine and olmesartan medoxomil)  
Formulary Dossier

**C. EVIDENCE TABLE SPREADSHEETS (NOTED ABOVE) OF ALL PUBLISHED AND UNPUBLISHED TRIALS:**

Study No. and Study Dates	Study Design/ Sample Size/Treatments	Inclusion & Exclusion Criteria	Endpoints/Results
<p>CS-8663-A-U301 A randomized, double-blind, placebo-controlled, factorial study evaluating the efficacy and safety of co-administration of olmesartan medoxomil plus amlodipine compared to monotherapy in patients with mild to severe hypertension. (Data on File 2007, Daiichi Sankyo, Inc.)</p> <p>Q2 2005 – Q1 2006</p>	<p><b>Design:</b> Randomized Double-blind Placebo-Controlled Multi-center Parallel-group factorial trial (Period II), 8 Weeks</p> <p><b>Patient type and Sample Size</b> Mild to severe hypertension N= 1940</p> <p><b>Treatment Arms</b> Placebo OM 10 mg OM 20 mg OM 40 mg AML 5 mg AML 10 mg AML 5 mg + OM 10 mg AML 5 mg + OM 20 mg AML 5 mg + OM 40 mg AML 10 mg + OM 10 mg AML 10 mg + OM 20 mg or AML 10 mg + OM 40 mg</p> <p><i>Subgroup Analyses of Age (&lt;65 vs ≥ 65 years old); Diabetes Status; and Race (Black vs Non-Black) on ITT</i></p>	<p><b>Inclusion:</b></p> <ul style="list-style-type: none"> <li>Male or female patients 18 yrs and older</li> <li>Female patients had to have a negative urine pregnancy test at screening, not lactating, and either post-menopausal for 1 year, surgically sterile, or using effective contraceptive methods</li> <li>Mean SeDBP of ≥ 95 mm Hg and ≤ 120 mmHg by Visit 2 or 2.1 (pre-randomization) and Visit 3 (randomization)</li> <li>Mean difference in SeDBP between the pre-randomization visit and randomization of ≤ 10 mm Hg</li> </ul> <p><b>Exclusion:</b></p> <ul style="list-style-type: none"> <li>SeDBP &gt; 120 mm Hg</li> <li>Uncontrolled Type 1 or Type 2 DM defined as HbA1c &gt; 9.0%</li> <li>History of heart failure</li> </ul>	<p><b>Primary</b></p> <ul style="list-style-type: none"> <li>Mean Change in SeDBP from baseline to Week 8 with LOCF – Intent to Treat <ul style="list-style-type: none"> <li>Each active treatment group had a statistically significant mean reduction in SeDBP from baseline to Week 8 with LOCF (p&lt;0.0001)</li> <li>Mean reductions in SeDBP were significantly greater in the combination therapy groups compared with the respective monotherapy groups. In the combination groups with AML 5 mg and 10 mg, increasing doses of OM (10 mg, 20 mg, and 40 mg) resulted in greater reductions in SeDBP.</li> <li>Overall, the greatest reductions in seated diastolic blood pressure occurred in the group treated with OM 40 mg + AML 10 mg (-19.0 mm Hg) followed by the group treated with OM 20 mg + AML 10 mg (-17.0 mmHg).</li> </ul> </li> </ul> <p><b>Secondary</b></p> <ul style="list-style-type: none"> <li>Mean Change in SeSBP from baseline to Week 8 with LOCF – Intent to Treat <ul style="list-style-type: none"> <li>Each active treatment group had a statistically significant mean reduction in SeSBP from baseline to Week 8 with LOCF (p&lt;0.0001)</li> <li>Mean reductions in SeSBP were significantly greater in the combination therapy groups compared with the respective monotherapy groups.</li> <li>Overall, the greatest reductions in seated systolic blood pressure occurred in the group treated with OM 40 mg + AML 10 mg (-30.1 mm Hg) followed by the group treated with OM 20 mg + AML 10 mg (-29.2 mm Hg).</li> </ul> </li> </ul>

AZOR™ (amlodipine and olmesartan medoxomil)  
Formulary Dossier

**CONTINUED - EVIDENCE TABLE SPREADSHEETS (NOTED ABOVE) OF ALL PUBLISHED AND UNPUBLISHED TRIALS**

Study No. and Study Dates	Study Design/ Sample Size/Treatments	Inclusion & Exclusion Criteria	Endpoints/Results
		<ul style="list-style-type: none"> <li>History of MI, percutaneous transluminal coronary revascularization, coronary artery bypass graft, and/or unstable angina within the past 6 months</li> <li>Hypertensive encephalopathy, stroke, or TIA within the past 6 months</li> <li>History of secondary hypertension</li> <li>Evidence of symptomatic resting bradycardia, hemodynamically significant cardiac valvular disease, and liver disease</li> <li>Presence of heart block greater than 1<sup>st</sup>-degree SA block, chronic atrial fibrillation, or flutter</li> <li>Uncorrected coarctation of the aorta, bilateral RAS, or unilateral RAS in a solitary kidney</li> </ul>	<ul style="list-style-type: none"> <li>Change in SeDBP and SeSBP from Baseline to Weeks 2, 4, 6, and 8 without LOCF               <ul style="list-style-type: none"> <li>Analysis results for the mean change in SeDBP and SeSBP from baseline to Week 2, Week 4, Week 6 and Week 8 without LOCF for the ITT population were similar to the results observed at Week 8 with LOCF.</li> <li>For all active treatment groups, most of the mean reduction in SeDBP and SeSBP occurred from baseline to Week 2, which plateaued by Week 4</li> </ul> </li> <li>Blood Pressure Goals               <ul style="list-style-type: none"> <li>The percentage of patients achieving their blood pressure goals by Week 8 with LOCF ranged from 20.0% to 36.3% for the groups treated with monotherapy compared with 35.0% to 53.2% for the groups treated with combination therapy.</li> <li>Approximately 50% of patients treated with one of the higher dose combination therapies (i.e., AML 10mg + OM 10 mg, AML 10 mg + OM 20 mg, AML 5 mg + OM 40 mg, and AML 10 mg + OM 40 mg) reached their blood pressure goal by Week 8 with LOCF</li> </ul> </li> </ul> <p>Subgroup analyses were conducted on the ITT population for age (&lt;65 vs ≥ 65 years old); diabetes status; and race (Black vs Non-Black). For each analyses, all active treatment arms had a statistically significant mean reduction in SeDBP and SeSBP from baseline. Combination therapy versus monotherapy resulted in significantly greater reductions in mean BP for those &lt;65 yo, patients without diabetes, and non-Black patients, in each of the respective subgroups analyzed. However, for the comparative subgroups (≥ 65 yo, patients with diabetes, and Black patients), the results were not consistent (see pages 26 - 33).</p>

AZOR™ (amlodipine and olmesartan medoxomil)  
Formulary Dossier

**CONTINUED - EVIDENCE TABLE SPREADSHEETS (NOTED ABOVE) OF ALL PUBLISHED AND UNPUBLISHED TRIALS**

Study No. and Study Dates	Study Design/ Sample Size/Treatments	Inclusion & Exclusion Criteria	Endpoints/Results																														
<p>CS-8663-A-U301</p> <p>A randomized, double-blind, placebo-controlled, factorial study evaluating the efficacy and safety of co-administration of olmesartan medoxomil plus amlodipine compared to monotherapy in patients with mild to severe hypertension. (Data on File 2007, Daiichi Sankyo, Inc.)</p> <p>Q2 2005 – Q1 2007</p> <p>(Phase III, US Study)</p>	<p><b>Design:</b> <b>Open-Label Treatment Week 8 to Week 52</b> (Period III) (Results from the double-blind period (Period II – Day 1 to Week 8) are above)</p> <p><b>Sample Size:</b> N=1684 (these patients completed Period II and entered Period III)</p> <p><b>Treatments:</b> Treat-to-goal All patients started on:</p> <ul style="list-style-type: none"><li>○ AML 5 mg + OM 40 mg</li></ul> <p>Titrated if goal of &lt;140/90 mm Hg or &lt;130/80 mm Hg for those patients with DM was not achieved:</p> <ul style="list-style-type: none"><li>○ AML 10 mg + OM 40 mg</li><li>○ AML 10 mg + OM 40 mg + HCTZ 12.5 mg</li><li>○ AML 10 mg + OM 40 mg + HCTZ 25 mg</li></ul> <p>If SeDBP was &gt;120 mm Hg while on maximal therapy, the patient was discontinued from the study</p>	<p><b>Inclusion:</b></p> <ul style="list-style-type: none"><li>• Male or female patients 18 yrs and older</li><li>• Female patients had to have a negative urine pregnancy test at screening, not lactating, and either post-menopausal for 1 year, surgically sterile, or using effective contraceptive methods</li><li>• Mean SeDBP of ≥ 95 mm Hg and ≤ 120 mm Hg by Visit 2 or 2.1 (pre-randomization) and Visit 3 (randomization)</li><li>• Mean difference in SeDBP between the pre-randomization visit and randomization of ≤ 10 mm Hg</li></ul> <p><b>Exclusion:</b></p> <ul style="list-style-type: none"><li>• SeDBP &gt; 120 mm Hg</li><li>• Uncontrolled Type 1 or Type 2 DM defined as HbA<sub>1c</sub> &gt; 9.0%</li><li>• History of heart failure</li><li>• History of MI, percutaneous transluminal coronary revascularization, coronary artery bypass graft, and/or unstable angina within the past 6 months</li></ul>	<p><b><u>Mean SeDBP and SeSBP and percentage of patients achieving blood pressure goals by week for each treatment</u></b></p> <p>Mean SeDBP and SeSBP and % of Patients Reaching BP Goal at Week 52 or Early Termination and by Dosing Regimen</p> <table><tr><th>Treatment (mgs)</th><th>n</th><th>Mean SeDBP (mm Hg)</th><th>Mean SeSBP (mm Hg)</th><th>% of Subjects Reaching BP Goal*</th></tr><tr><td>AML 5/OM 40</td><td>525</td><td>81.0</td><td>127.6</td><td>80.0</td></tr><tr><td>AML 10/OM 40</td><td>378</td><td>82.4</td><td>130.9</td><td>70.6</td></tr><tr><td>AML 10/OM 40 + HCTZ 12.5</td><td>287</td><td>81.0</td><td>130.7</td><td>66.6</td></tr><tr><td>AML 10/OM 40 + HCTZ 25</td><td>419</td><td>83.4</td><td>136.8</td><td>46.3</td></tr><tr><td>Other**</td><td>63</td><td>79.4</td><td>126.2</td><td>68.3%</td></tr></table> <p>*&lt;140/90 mm Hg or &lt;130/80 for patients with diabetes ** Despite the treatment algorithm presented, some investigators elected for a variety of reasons to use alternative antihypertensive medications</p> <ul style="list-style-type: none"><li>• From the mean baseline BP of 163.6/101.5 mm Hg, BP reductions were observed across all combination treatment regimens to Week 52. Also, each titration to a more intensive treatment regimen resulted in additional mean reductions of SeBP.</li><li>• The groups of patients who required titration of the amlodipine dose or the addition of HCTZ were more severe hypertensive patients and/or were more resistant to the antihypertensive effects of treatment</li></ul>	Treatment (mgs)	n	Mean SeDBP (mm Hg)	Mean SeSBP (mm Hg)	% of Subjects Reaching BP Goal*	AML 5/OM 40	525	81.0	127.6	80.0	AML 10/OM 40	378	82.4	130.9	70.6	AML 10/OM 40 + HCTZ 12.5	287	81.0	130.7	66.6	AML 10/OM 40 + HCTZ 25	419	83.4	136.8	46.3	Other**	63	79.4	126.2	68.3%
Treatment (mgs)	n	Mean SeDBP (mm Hg)	Mean SeSBP (mm Hg)	% of Subjects Reaching BP Goal*																													
AML 5/OM 40	525	81.0	127.6	80.0																													
AML 10/OM 40	378	82.4	130.9	70.6																													
AML 10/OM 40 + HCTZ 12.5	287	81.0	130.7	66.6																													
AML 10/OM 40 + HCTZ 25	419	83.4	136.8	46.3																													
Other**	63	79.4	126.2	68.3%																													

AZOR™ (amlodipine and olmesartan medoxomil)  
Formulary Dossier

**CONTINUED - EVIDENCE TABLE SPREADSHEETS (NOTED ABOVE) OF ALL PUBLISHED AND UNPUBLISHED TRIALS**

Study No. and Study Dates	Study Design/ Sample Size/Treatments	Inclusion & Exclusion Criteria	Endpoints/Results
		<ul style="list-style-type: none"> <li>• Hypertensive encephalopathy, stroke, or TIA within the past 6 months</li> <li>• History of secondary hypertension</li> <li>• Evidence of symptomatic resting bradycardia, hemodynamically significant cardiac valvular disease, and liver disease</li> <li>• Presence of heart block greater than 1<sup>st</sup>-degree SA block, chronic atrial fibrillation, or flutter</li> <li>• Uncorrected coarctation of the aorta, bilateral RAS, or unilateral RAS in a solitary kidney</li> </ul>	<ul style="list-style-type: none"> <li>• As patients were titrated to more intensive treatment regimens, an overall greater percentage of patients reached their BP treatment goal</li> </ul>

AZOR™ (amlodipine and olmesartan medoxomil)  
Formulary Dossier

**CONTINUED - EVIDENCE TABLE SPREADSHEETS (NOTED ABOVE) OF ALL PUBLISHED AND UNPUBLISHED TRIALS**

Study No. and Study Dates	Study Design/ Sample Size/Treatments	Inclusion & Exclusion Criteria	Endpoints/Results
<p>CS-8663-A-E302 Efficacy and safety of Amlodipine used as add-on therapy in moderately to severely hypertensive patients not adequately controlled by olmesartan medoxomil 20 mg monotherapy (Data on File 2007, Daiichi Sankyo, Inc.)</p> <p>Q4 2005 – Q4 2006</p> <p>(Phase III, European Study)</p>	<p><b>Design:</b> 8 week open-label monotherapy (Period I) and 8 week randomized double-blind treatment (Period II); Parallel-group Multi-national Multi-center</p> <p><b>Patient Type and Sample Size:</b> Moderate to Severe hypertension or uncontrolled on OM monotherapy Period I, n = 722 Period II, n=538</p> <p><b>Treatment Arms:</b> <b>Period I:</b> OM 20 mg (all patients received monotherapy, n = 722) <b>Period II:</b> Non-responders from Period I randomized to:</p> <ul style="list-style-type: none"> <li>Placebo + OM 20 mg</li> <li>AML 5 mg + OM 20 mg</li> <li>AML 10 mg + OM 20 mg</li> </ul> <p><i>(Non-responder defined as mean trough SeBP <math>\geq</math> 140/90, mean 24-hour DBP <math>\geq</math> 80 mm Hg, and at least 30% of daytime DBP readings <math>&gt;</math>85 mm Hg)</i></p>	<p><b>Inclusion:</b></p> <ul style="list-style-type: none"> <li>Male or female patients 18 yrs or older</li> <li>History of moderate to severe hypertension (SBP <math>\geq</math> 160 mm Hg &amp; DBP <math>\geq</math> 100 mm Hg; mean 24 hours DBP of <math>\geq</math> 84 mm Hg at Visit 2; and at least 30% of daytime DBP readings <math>&gt;</math> 90 mm Hg</li> <li>Patients treated with OM 20 or 40 mg had to have a previous diagnosis of moderate to severe hypertension, were uncontrolled on monotherapy (<math>\geq</math> 140/90 mm Hg), mean 24 hour DBP of <math>\geq</math> 80 mm Hg at Visit 2; and at least 30% of daytime DBP readings <math>&gt;</math> 85 mm Hg</li> </ul> <p><b>Exclusion</b></p> <ul style="list-style-type: none"> <li>Mean SeDBP <math>&gt;</math> 115 mm Hg</li> <li>Mean SeSBP <math>&gt;</math> 200 mm Hg</li> <li>Mean 24-hour DBP as assessed by 24 hour ABPM <math>&gt;</math>104 mmHg</li> <li>Bradycardia (<math>&lt;</math>50 beats/min)</li> </ul>	<p><b>Primary</b></p> <ul style="list-style-type: none"> <li>Mean change from baseline (Week 8) to Week 16 (end of double-blind treatment period) in trough SeDBP using LOCF <ul style="list-style-type: none"> <li>Combination arms (AML 5/OM 20 &amp; AML 10/OM 20) reduced the mean SeDBP to a significantly greater extent than treatment with OM 20 + placebo; -2.7 mmHg for AML 5/OM 20 (p=0.0006 vs. monox) and -3.2 mmHg for AML 10/OM 20 (p&lt; 0.0001 vs. monox)</li> </ul> </li> </ul> <p><b>Secondary</b></p> <ul style="list-style-type: none"> <li>Mean change from baseline (Week 8) to Week 16 (end of double-blind) in trough SeSBP using LOCF <ul style="list-style-type: none"> <li>Combination arms (AML 5/OM 20 &amp; AML 10/OM 20) resulted in statistically significant reductions in adjusted mean sitting SBP when compared with OM 20 mg + placebo therapy: -5.8 mm Hg for AML 5 mg + OM 20 mg (p&lt;0.0001) and -6.4 mmHg for AML 10 mg + OM 20 mg (p&lt;0.0001)</li> </ul> </li> <li>Mean change from baseline (Week 8) to Week 16 (end of double-blind) in daytime, nighttime, and 24-hour DBP and SBP assessed by 24-hour ABPM <ul style="list-style-type: none"> <li>Combination arms resulted in statistically significant reductions in 24-hour, daytime, and nighttime adjusted mean changes for DBP and SBP when compared with monotherapy</li> </ul> </li> <li>Mean change from baseline (Week 8) to Week 12 in trough SeDBP &amp; SeSBP w/o LOCF <ul style="list-style-type: none"> <li>The adjusted mean change in SeDBP at Week 12 for patients treated with OM 20 mg + AML 10 mg combination therapy (-2.8 mmHg) was statistically significant when compared to OM 20 mg + placebo therapy (p=0.0002). The adjusted mean change in sitting DBP at Week 12 for patients treated with OM 20 mg + AML 5 mg was -0.9 mmHg when compared to OM 20 mg + placebo therapy (p=0.3703)</li> </ul> </li> </ul>



AZOR™ (amlodipine and olmesartan medoxomil)  
Formulary Dossier

**CONTINUED - EVIDENCE TABLE SPREADSHEETS (NOTED ABOVE) OF ALL PUBLISHED AND UNPUBLISHED TRIALS**

Study No. and Study Dates	Study Design/ Sample Size/Treatments	Inclusion & Exclusion Criteria	Endpoints/Results
			<ul style="list-style-type: none"> <li>○ Adjusted mean change in SeSBP at Week 12: The differences from OM 20 mg + placebo in adjusted mean change in SeSBP for both the OM 20 mg + AML 5 mg treatment group (-2.9 mm Hg; p=0.0220) and the OM 20 mg + AML 10 mg treatment group (-6.3 mm Hg; p&lt;0.0001) were statistically significant.</li> <li>○ The difference in BP reduction between the OM 20mg + placebo arm and combination arms was greater in the OM 20 mg + AML 10 mg arm, compared to the OM 20 mg + AML 5 mg arm. However, with time, the differences between the 2 combinations were narrowed by week 16.</li> <li>• Blood Pressure Goals Compared to patients treated with OM 20 mg + placebo (28.5% achieving goal), the percentage of patients achieving BP goal at week 16 with LOCF was significantly higher in the OM 20 mg + AML 5 mg treatment group (44.5%; p=0.0011) and in the OM 20 mg + AML 10 mg treatment group (45.8%; p=0.0004)</li> </ul>

AZOR™ (amlodipine and olmesartan medoxomil)  
Formulary Dossier

**CONTINUED - EVIDENCE TABLE SPREADSHEETS (NOTED ABOVE) OF ALL PUBLISHED AND UNPUBLISHED TRIALS**

Study No. and Study Dates	Study Design/ Sample Size/Treatments	Inclusion & Exclusion Criteria	Endpoints/Results
<p>CS-8663-A-E303 Add-on Study of olmesartan medoxomil in patients with moderate to severe hypertension not achieving target blood pressure on amlodipine 5mg alone (Data on File 2007, Daiichi Sankyo, Inc.)</p> <p>Q4 2005 – Q4 2006 (to Week 34)</p> <p>Phase III, European Study</p>	<p><b>Design:</b> 52 week study consisting of 4 periods: Period I: Open-label monotherapy Period II: Double-blind treatment Period III: Double-blind treatment with dose up-titration Period IV: Open-label Combination Treatment Randomized Parallel-group Multi-national Multi-center</p> <p><b>Patient type and Sample Size</b> Moderate to severe hypertension (<math>\geq 160/100</math>) or uncontrolled on AML monotherapy (<math>\geq 140/90</math>) Period 1: n=1017 Period 2: n=755 Period 3: n=706 Period 4: n=677</p> <p>Treatment Arms: Period I: AML 5 mg (8 weeks) Period II: Non-responders from Period I randomized to (8 weeks):</p> <ul style="list-style-type: none"> <li>• AML 5 mg + placebo</li> <li>• AML 5 mg + OM 10 mg</li> <li>• AML 5 mg + OM 20 mg</li> <li>• AML 5 mg + OM 40 mg</li> </ul>	<p><b>Inclusion:</b></p> <ul style="list-style-type: none"> <li>• Male or female patients 18 yrs or older</li> <li>• History of moderate to severe hypertension (SBP <math>\geq 160</math> mm Hg &amp; DBP <math>\geq 100</math> mm Hg; mean 24 hours DBP of <math>\geq 84</math> mm Hg at Visit 2; and at least 30% of daytime DBP readings <math>&gt; 90</math> mm Hg</li> <li>• Patients treated with AML 5 mg or 10 mg had to have a previous diagnosis of moderate to severe hypertension, were uncontrolled on monotherapy (<math>\geq 140/90</math> mm Hg), mean 24 hour DBP of <math>\geq 80</math> mm Hg at Visit 2; and at least 30% of daytime DBP readings <math>&gt; 85</math> mm Hg</li> </ul> <p><b>Exclusion</b></p> <ul style="list-style-type: none"> <li>• Mean SeDBP <math>&gt; 115</math> mm Hg</li> <li>• Mean SeSBP <math>&gt; 200</math> mm Hg</li> <li>• Mean 24-hour DBP as assessed by 24 hour ABPM <math>&gt; 104</math> mm Hg</li> <li>• Bradycardia (<math>&lt; 50</math> beats/min)</li> </ul>	<p><b>Primary</b></p> <ul style="list-style-type: none"> <li>• Mean change in trough sitting DBP from baseline (Week 8) to Week 16 – Period II <ul style="list-style-type: none"> <li>◦ Combination arms (AML 5/OM 10, AML 5/OM 20, &amp; AML 5/OM 40) reduced the mean SeDBP to a significantly greater extent than treatment with AML 5mg + placebo; -2.0 mm Hg for AML 5/OM 10 (<math>p=0.0207</math> vs. monox); -3.7 mm Hg for AML 5/OM 20 (<math>p&lt;0.0001</math> vs. monox); and -3.8 mm Hg for AML 5/OM 40 (<math>p&lt;0.0001</math> vs. monox)</li> </ul> </li> </ul> <p><b>Secondary</b></p> <ul style="list-style-type: none"> <li>• Mean change in trough SeSBP from baseline (Week 8) to Week 16 – Period II <ul style="list-style-type: none"> <li>◦ Compared with placebo + AML 5 mg, treatment with AML + OM resulted in statistically significant reductions in sitting SBP at Week 16 with LOCF (-3.5 mm Hg, <math>p=0.0103</math> for OM 10 mg + AML 5 mg; -5.8 mm Hg, <math>p&lt;0.0001</math> for OM 20 mg + AML 5 mg; and -7.1 mm Hg, <math>p&lt;0.0001</math> for OM 40 mg + AML 5 mg)</li> </ul> </li> <li>• Mean change in trough SeDBP and SeSBP from baseline to Week 12, from Week 16 to Week 20 and 24; and at Week 34 <ul style="list-style-type: none"> <li>◦ Combination arms demonstrated significantly larger mean reductions in SeDBP and SeSBP than AML 5 mg + placebo at Week 12 and Week 16; the DBP lowering effect of AML 5/OM 40 occurred earlier than that of AML 5/OM 20, however at Week 16 both arms had numerically larger mean reductions in SeDBP than AML 5/OM 10</li> <li>◦ Mean change in trough SeDBP and SeSBP from Week 16 to Week 20 and 24 was not meaningful for patients not titrated; however, for those who were titrated further mean reductions in SeDBP and SeSBP occurred</li> <li>◦ At Week 34, mean SeBP for the AML 5/OM 40 arm was 132.2/83.6 mm Hg; for the AML 10/OM 40 arm was 143/90.3 mm Hg; for the AML 10/OM 40/HCTZ 12.5 arm was 147.6/89.3; and for the AML 10/OM 40/HCTZ 25 arm was 155.3/92 mm Hg</li> </ul> </li> </ul>

AZOR™ (amlodipine and olmesartan medoxomil)  
Formulary Dossier

**CONTINUED EVIDENCE TABLE SPREADSHEETS (NOTED ABOVE) OF ALL PUBLISHED AND UNPUBLISHED TRIALS**

Study No. and Study Dates	Study Design/ Sample Size/Treatments	Inclusion & Exclusion Criteria	Endpoints/Results
	<p><i>(Non-responder defined as mean trough SeBP <math>\geq</math> 140/90, mean 24-hour DBP <math>\geq</math> 80 mm Hg, and at least 30% of daytime DBP readings <math>&gt;</math>85 mm Hg)</i></p> <p>Period III: Patients uncontrolled in Period II had doses titrated (all others continued on assigned treatment regimen) (8 weeks):</p> <ul style="list-style-type: none"> <li>• AML 5 mg + placebo to AML 5 mg + OM 20 mg</li> <li>• AML 5 mg + OM 10 mg to AML 5 mg + OM 20 mg</li> <li>• AML 5 mg + OM 20 mg to AML 5 mg + OM 40 mg</li> <li>• AML 5 mg + OM 40 mg to AML 10 mg + OM 40 mg</li> </ul> <p>Period IV: Open-label Combination Treatment (28 weeks):</p> <ul style="list-style-type: none"> <li>• Initial treatment with AML 5 mg + OM 40 mg</li> <li>• Then: AML 10 mg + OM 40 mg (if needed)</li> <li>• Then: AML 10 mg + OM 40 mg + HCTZ 12.5 mg (if needed)</li> <li>• Then: AML 10 mg + OM 40 mg + HCTZ 12.5 mg (if needed)</li> </ul>		<ul style="list-style-type: none"> <li>• Mean change from baseline (Week 8) to Week 16 (end of double-blind) and from Week 16 to Week 24 in daytime, nighttime, and 24-hour DBP and SBP assessed by 24-hour ABPM <ul style="list-style-type: none"> <li>◦ Combination arms resulted in statistically significant reductions in 24-hour, daytime &amp; nighttime adjusted mean changes for DBP and SBP when compared with monotherapy</li> </ul> </li> </ul> <p><b><u>Blood Pressure Goals</u></b></p> <ul style="list-style-type: none"> <li>• After double-blind treatment, compared with AML 5 mg + placebo, treatment with the combinations resulted in a statistically significantly higher proportion of patients who reached BP goal at Week 16 with LOCF (50.5% - AML 5/OM 40; 53.5% - AML 5/OM 20; and 39.2% - AML 5/OM 10; versus 29.9% - AML 5/placebo)</li> <li>• In Period III, for patients who remained on their randomized treatment regimen, the proportion who reached BP goal at Week 24 with LOCF was higher with AML + OM treatment than with AML 5mg + placebo. For patients whose dose regimen was titrated, successively higher proportions reached BP goal with each increase in dose combination of AML + OM.</li> </ul>

### 3 SUPPORTING INFORMATION

#### 3.1 REFERENCES

##### a. REFERENCES IN DOSSIER

- American Heart Association. *Heart Disease and Stroke Statistics – 2007 Update*. Dallas, Texas: American Heart Association; 2007.
- Arakawa K, Maruta H. Ability of kallikrein to generate angiotensin II-like pressor substance and a proposed kinin-tensin enzyme system. *Nature*. 1980;288:705-706.
- Beevers G, Lip GYH, O'Brien E. ABC of Hypertension: The Pathophysiology of Hypertension. *BMJ*. 2001;322:912-916.
- Brunner HR, Laragh JH, Baer L, et al. Essential hypertension: renin and aldosterone, heart attack and stroke. *N Engl J Med*. 1972;286:441-449.
- Burnier M, Brunner HR: Angiotensin II receptor antagonists. *Lancet*. 2000;355:637-645.
- Burnier M: Angiotensin II type 1 receptor blockers. *Circulation*. 2001;103:904-912.
- Chobanian, A, Bakris, G, Black H, et al. The Sixth Report of the Joint National Committee on Prevention, Detection, and Treatment of High Blood Pressure: The JNC 6 Report. *JAMA*. 1997.
- Chobanian, A, Bakris, G, Black H, et al. The Seventh Report of the Joint National Committee on Prevention, Detection, and Treatment of High Blood Pressure: The JNC 7 Report. *JAMA*. 2003.
- CONSENSUS Trial Study Group. Effects of enalapril on mortality in severe congestive heart failure: results of the Cooperative North Scandinavian Enalapril Survival Study. *N Engl J Med*. 1987;316:1429-1435.
- Cooper R, Cutler J, Desvigne-Nickens P, et al: Trends and disparities in coronary heart disease, stroke, and other cardiovascular diseases in the United States: findings of the national conference on cardiovascular disease prevention. *Circulation*. 2000;102:3137-3147.
- Data on file. Daiichi Sankyo, Inc., Parsippany, NJ.
- Dina R, Jafari M: Angiotensin II-receptor antagonists: an overview. *Am J Health Syst Pharm*. 2000;57:1231-1241.
- Elliot WJ, Black HR. Calcium antagonists in hypertension. *Blood Pressure*. 2001; 10:387-397.
- Faulkner MA, Hilleman DE. Amlodipine/benazepril: fixed dose combination therapy for hypertension. *Expert Opin Pharmacother*. 2001;2:165-178.
- Gavras H, Lever AF, Brown JJ, et al. Acute renal failure, tubular necrosis, and myocardial infarction induced in the rabbit by intravenous angiotensin II. *Lancet*. 1971; 2:19-22.
- Gerbino PP, Shoheiber O. Adherence Patterns Among Patients Treated with Fixed-Dose Combination versus Separate Antihypertensive Agents. *AJHP*. 2007;64(12):1279-1283.
- Heart Outcomes Prevention Evaluation Study Investigators, HOPE. Effects of ramipril on cardiovascular and microvascular outcomes in people with diabetes mellitus: results of the HOPE study and MICRO-HOPE substudy. *Lancet*. 2000;355:253-259.
- Hernandez-Hernandez RM, Velasco MJ, Armas-Hernandez MJ, et al. Update on the use of calcium antagonists on hypertension. *J Hum Hypertens*. 2002; 16, no. Suppl 1:S114-S117.

AZOR™ (amlodipine and olmesartan medoxomil)  
Formulary Dossier

Ihara M, Urata H, Kinoshita A, et al. Increased chymase-dependent angiotensin II formation in human atherosclerosis aorta. *Hypertension*. 1999;33:1399-1405.

Lewis EJ, Hunsicker LG, Bain RP, Rohde RD. The effect of angiotensin-converting-enzyme inhibition on diabetic nephropathy: the Collaborative Study Group. *N Engl J Med*. 1993;329:1456-1462.

Pickering T: Headache and hypertension—something old, something new. *J Clin Hypertens* 2000;2:345–347.

Rudd P, Dzau VJ: Hypertension: evaluation and management. In: Loscalzo J, Creager MA, Dzau VJ, eds. *Vascular medicine*. Boston: Little, Brown, 1996:609–638.

Scholze J, Bida M, Hansen A, et al. Initiation of Hypertensive Treatment with a Fixed-Dose Combination or its Monocomponents – Does it Really Matter? *J Clin Pract*. 2006;60(3):265-274.

Sica DA. Rationale for Fixed-Dose Combinations in the Treatment of Hypertension: The Cycle Repeats. *Drugs*. 2002;62(3):443-462.

SOLVD Investigators. Effect of enalapril on survival in patients with reduced left ventricular ejection fractions and congestive heart failure. *N Engl J Med*. 1991;325:294-302.

SOLVD Investigators. Effect of enalapril on mortality and the development of heart failure in asymptomatic patients with reduced left ventricular ejection fractions. *N Engl J Med*. 1992;327:685-691.

Stergiou GS. Combination Pharmacotherapy in Hypertension. *International Urology and Nephrology*. 2006;38:673-682.

Stern MD. Theory of Excitation-Contraction Coupling in Cardiac Muscle. *Biophys J*. 1992;63:497-517.

Tejada T, Fornoni A, Lenz O, et al. Combination Therapy with Rennin-Angiotensin System Blockers: Will Amlodipine Replace Hydrochlorothiazide? *Current Hypertension Reports*. 2007;9:284-290.

Uehara Y, Urata H, Sasaguri M, et al. Increased chymase activity in internal thoracic artery of patients with hypercholesterolemia. *Hypertension*. 2000;35:55-60.

Warnholtz A, Nickenig G, Schulz E, et al. Increased NADH-oxidase-mediated superoxide production in the early stages of atherosclerosis: evidence for involvement of the renin-angiotensin system. *Circulation*. 1999;99:2027-2033.

Yusuf S, Sleight P, Pogue J, et al: Effects of an angiotensin-converting-enzyme inhibitor, ramipril, on cardiovascular events in high-risk patients. The Heart Outcomes Prevention Evaluation Study Investigators. *N Engl J Med*. 2000;342:145-153.

AZOR™ (amlodipine and olmesartan medoxomil)  
Formulary Dossier

**b. AMLODIPINE AND OLMESARTAN MEDOXOMIL CLINICAL BIBLIOGRAPHY**

Haworth S, Bathala M, Lee J, et al. A Fixed-Dose Combination of Olmesartan Medoxomil (OM) and Amlodipine Besylate (AM) is Bioequivalent (BEQ) to Free Combination of the Agents. *Journal of Clinical Hypertension*. 2007;9(Suppl. A):181.

Haworth S, Heyrman R, Bathala M, et al. The Bioavailability (BAV) of a Fixed-Dose Combination of Olmesartan Medoxomil (OM) and Amlodipine Besylate (AM) is Unaffected by Food. *Journal of Clinical Hypertension*. 2007;9(Suppl. A):181-182.

Rohatagi S, Haworth S, Heyrman R, et al. Safety and Tolerability of the Combination of Olmesartan Medoxomil (OM) and Amlodipine Besylate (AM). *Journal of Clinical Hypertension*. 2007; 9 (Suppl. A):179 - 180.

Rohatagi S, Carrothers T, Kshirsagar S, et al. Evaluation of Population (POP) Pharmacokinetics (PK) After Coadministration of Olmesartan Medoxomil (OM) and Amlodipine Besylate (AM). *Hypertension*. 2007;9 (Suppl. A):181-182.

Rohatagi S, Haworth S, Salazar D, et al. Olmesartan Medoxomil (OM) and Amlodipine Besylate (AM) Show Dose-Proportional Drug Exposure When Used in Combination in Healthy Volunteers (VOLS). *Journal of Clinical Hypertension*. 2007;9 (Suppl. A):180.

Rohatagi S, Haworth S, Heyrman R, et al. Low and High Doses of a Fixed-Dose Combination of Olmesartan Medoxomil (OM) and Amlodipine Besylate (AM) are Bioequivalent (BEQ) to Free Combination of the Agents. *Journal of Clinical Hypertension*. 2007; 9(Suppl. A):178 – 179.

Salazar D, Lee J, Shenouda M, et al. Lack of Pharmacokinetic (PK) Drug Interaction Between Olmesartan Medoxomil (OM) and Amlodipine Besylate (AM) During Coadministration. *Journal of Clinical Hypertension*. 2007;9 (Suppl. A):180-181.

AZOR™ (amlodipine and olmesartan medoxomil)  
Formulary Dossier

**C. OLMESARTAN MEDOXOMIL CLINICAL BIBLIOGRAPHY**

Agabiti-Rosei E, Rodicio JL. Introduction. *Clinical Therapeutics*. 2004;26(suppl a):a1-a2.

Alain T, Mallat Z. Hypertension: A novel regulator of adaptive immunity in atherosclerosis? *Hypertension*. 2004;44:257-258.

Alexander LM. Desirable therapeutic characteristics of an optimal antihypertensive agent. *Drugs* 2006;66(9):1239-1252.

Almendral J, Shick VI, Atlas SA, et al. Transforming growth factor - 1: A marker of left ventricular hypertrophy in hypertension. *AJH*. (Abstract/posters: antihypertensive drugs and pharmacology) 2005;18(5): part 2.

Almendral JL, Atlas SA, Rosendorff C. Tgf-β1 is a marker for left ventricular hypertrophy in hypertensive patients. 21st Scientific Meeting of International Society of Hypertension (ISH), October 2006. [Abstract]

Almendral JL, Atlas SA, Rosendorff C. Transforming growth factor-β1 (tgf-β1) predicts the severity of left ventricular hypertrophy in hypertensive patients. November 2004. [Abstract]

Andros V, Egger A, Dua U. Blood pressure goal attainment according to JNC 7 guidelines and utilization of antihypertensive drug therapy in MCO patients with type 1 or type 2 diabetes. *J Manag Care Pharm*. 2006;12(4):303-09.

Anonymous. Changed guidelines of the German High Blood Pressure League. AT(1) blocker in the first league now. *MMW Fortschritte der Medizin*. 2002 ;144(44):55.

Anonymous. New AT1 receptor antagonist. Rapid and effective blood pressure decrease. *MMW Fortschritte der Medizin*. 2003 Jan 30;145(5):53.

Anonymous. New AT1-blocker is approved; "me too" or progress? *MMW Fortschritte der Medizin*. 2002; 144(40):51.

Anonymous. The Medical Letter. Olmesartan (Benicar) for Hypertension. *The Medical Letter on Drugs and Therapeutics*. 2002; 44: 69-71.

Anonymous. Therapy of Hypertension. AT1 blockers soon to be the first choice? *MMW Fortschritte der Medizin*. 2002;144(19):64.

Anonymous. Therapy of Hypertension. A new rapidly effective AT1 blocker. *MMW Fortschritte der Medizin*. 2002;144(25): 55.

Arakawa K. Development of combined drugs (ARB and diuretics). *Nippon Rinsho*. 2004;62(suppl 3):662-5. Review.

Aranda P, Ruilope LM, Esteban A, et al. Long-term comparative antiproteinuric effect of single versus combination therapy based on olmesartan in non-diabetic nephropathies. 65th Annual American Society of Hypertension Conference, May 2006. [Abstract]

Asmar R. Targeting effective blood pressure control with angiotensin receptor blockers. *Int J Clin Pract*. 2006;60(3):315-20.

Bahlmann FH, De Groot K, Mueller O, et al. Stimulation of endothelial progenitor cells: A new putative therapeutic effect of angiotensin ii receptor antagonist. *Hypertension*. 2005;45:526-529.

Bakris G, Winer N, Jones M. Effects of olmesartan medoxomil (OLM) and losartan potassium (LOS) in patients with hypertension (HTN), type 2 diabetes mellitus (T2DM), and microalbuminuria. *American Journal of Hypertension*. 2004;17(5):88A-89A. (P-150)

AZOR™ (amlodipine and olmesartan medoxomil)  
Formulary Dossier

**OLMESARTAN MEDOXOMIL CLINICAL BIBLIOGRAPHY**  
(continued)

Ball K. A Multi-Centre, Double-Blind, Efficacy, Tolerability and Safety Study of the Oral Angiotensin II-Antagonist Olmesartan Medoxomil versus Losartan in Patients with Mild to Moderate Essential Hypertension. *Journal of HTN*. 2001; 19 (suppl 2): P2.176.

Ban E, Mathe L, Major J. Effect of olmesartan and antioxidant C and E on microinflammation and serum lipid levels in patients with hypertension. 15th European Meeting on Hypertension, June 2005.

Barrios V, Boccanelli A, Ewald S, et al. Efficacy and tolerability of olmesartan medoxomil in patients with mild to moderate essential hypertension: the OLMEBEST study. *Clin Drug Invest* 2007;27(8):545-558.

Basile JN, Chrysant S. The importance of early antihypertensive efficacy: The role of angiotensin II receptor blocker therapy. *J Hum Hypertens*. 2006;20:169-175.

Bayés M, Rabasseda X, Prous JR. Gateways to clinical trials. Methods find. *Exp Clin Pharmacol*. 2004;26(7): 587-612.

Bayes M, X Rabasseda, JR Prous. Gateways to Clinical Trials. *Methods & Findings in Experimental & Clinical Pharmacology*. 2002; 24: 37-55.

Bohm M, Ewald S. Blood pressure reduction with olmesartan in mild-to-moderate essential hypertension: A planned interim analysis of an open label sub-study in German patients. *Current Medical Research & Opinion*. 2006;22(7):1375-80.

Brousil JA, Burke JM. Olmesartan medoxomil: An angiotensin II-receptor blocker. *Clinical Therapeutics*. 2003; 25(4):1041-55.

Brunner H, Fujita T. Olmesartan: a new A II antagonist in cardiovascular risk prevention. *Journal of Human Hypertension*. 2002. 16.

Brunner HR, Arakawa K. Antihypertensive efficacy of olmesartan medoxomil and candesartan cilexetil in achieving 24-hour blood pressure reductions and ambulatory blood pressure goals. *Clin Drug Invest*. 2006;26(4):185-193.

Brunner HR, Laeis P. Clinical efficacy of olmesartan medoxomil. *Journal of Hypertension*. 2003 May; suppl21(2):43-6.

Brunner HR, Stumpe KO, Januszewicz A. Antihypertensive Efficacy of Olmesartan Medoxomil and Candesartan Cilexetil Assessed by 24-Hour Ambulatory Blood Pressure Monitoring in Patients with Essential Hypertension. *Clin Drug Invest*. 2003;23(7):419-430.

Brunner HR. The new oral angiotensin II antagonist olmesartan medoxomil: a concise overview. *J Hum HTN*. 2002; 16 (Suppl 2): S13-S16.

Brunner HR. Clinical efficacy and tolerability of olmesartan. *Clinical Therapeutics*. 2004;26(suppl a):a28-a32.

Brunner HR. Olmesartan medoxomil: Current status of its use in monotherapy. *Vasc Health Risk Manag*. 2006;2(4):327-40.



AZOR™ (amlodipine and olmesartan medoxomil)  
Formulary Dossier

**OLMESARTAN MEDOXOMIL CLINICAL BIBLIOGRAPHY**  
(continued)

Burnier M., Brunner HR. Angiotensin II receptor antagonists: antihypertensive agents. *Expert Opinion on Investigational Drugs*. 1997; 6: 489-500.

Celentano C, Prefumo F, di Vera E, Iannicco A, Gallo DP, Liberati M. Reversible Acute Fetal Renal Failure Due to Maternal Exposure to Angiotensin Receptor Blocker. *Pediatr Nephrol*. 2007 Sep 26.

Chan P, Cheng TH, Liu JC. Angiotensin ii antagonist olmesartan inhibits angiotensin ii induces endothelin-1 gene expression via extracellular signal-regulated kinase pathway in rat aortic smooth muscle cells. *Journal of Hypertension*. 2004; 22(s2):s41. [Abstract p1.75]

Charles JA, Jotkowitz S, Byrd LH. Prevention of migraine with olmesartan in patients with hypertension/prehypertension. *Headache*. 2006;46:503-507.

Chilman-Blair K, Rabasseda X. Olmesartan, An AT1-Selective Antihypertensive Agent. *Drugs of Today*. 2003;39(10):745-761.

Chrysant S, Dimas B, Shiraz M. Treatment of hypertension with olmesartan medoxomil, alone and in combination with a diuretic: An update. *J Hum Hypertens*. 2007;21:699-708.

Chrysant S, Dubiel R, Walker F, et al. Efficacy of olmesartan medoxomil and olmesartan/hydrochlorothiazide in achieving blood pressure goals in patients with stage 2 systolic hypertension: Effect of body mass index. 65th Annual American Society of Hypertension Conference, May 2006. [Poster]

Chrysant S, Dubiel R, Walker F, et al. Efficacy of olmesartan medoxomil and olmesartan medoxomil/hydrochlorothiazide in achieving blood pressure goals in patients with stage 2 systolic hypertension: Effect in non-black patients. 65th Annual American Society of Hypertension Conference, May 2006. [Poster]

Chrysant S, Dubiel R, Walker F, et al. Efficacy of olmesartan medoxomil and om/hctz in achieving blood pressure goals in patients with stage 2 systolic hypertension: Effect of body mass index. 65th Annual American Society of Hypertension Conference, May 2006. [Abstract]

Chrysant S, Dubiel R, Xu J. Correlation between blood pressure reduction and baseline blood pressure in patients receiving olmesartan medoxomil and/or hydrochlorothiazide in a factorial-design study. 22nd Annual American Society of Hypertension Scientific Meeting, May 2007. [Poster]

Chrysant S, Dubiel R, Xu J. Correlation between blood pressure reduction and baseline blood pressure in patients receiving olmesartan medoxomil and/or hydrochlorothiazide in a factorial-design study. 22nd Annual American Society of Hypertension Scientific Meeting, May 2007. [Abstract]

Chrysant S, Neutel J, Dubiel R, et al. Efficacy of olmesartan medoxomil plus hydrochlorothiazide in patients stratified by race, gender, age, bmi and isolated systolic hypertension. 17th Annual European Meeting on Hypertension, June 2007. [Poster]

Chrysant S, Neutel J, Dubiel R, et al. Efficacy of olmesartan medoxomil plus hydrochlorothiazide in patients stratified by race, gender, age, bmi and isolated systolic hypertension. 22nd Annual American Society of Hypertension Scientific Meeting, May 2007. [Poster]

AZOR™ (amlodipine and olmesartan medoxomil)  
Formulary Dossier

**OLMESARTAN MEDOXOMIL CLINICAL BIBLIOGRAPHY**  
(continued)

Chrysant S, Neutel J, Dubiel R, et al. Efficacy of treating stage 2 systolic hypertension with olmesartan medoxomil and olmesartan medoxomil/hydrochlorothiazide in black and non-black patients. 17th Annual European Meeting on Hypertension, June 2007. [Poster]

Chrysant S, Neutel J, Xu J, et al. Distribution of blood pressure response in patients receiving an olmesartan medoxomil-based titration regimen. 17th Annual European Meeting on Hypertension, June 2007. [Poster]

Chrysant S, Wang A, Jones M. Efficacy of olmesartan medoxomil (OLM), losartan potassium (LOS) and valsartan (VAL) in a non-black patient (pt) cohort. *AJH*. 2005;18(5): part 2. [Abstract]

Chrysant S, Wang A. Olmesartan medoxomil lowers blood pressure as rapidly as amlodipine besylate in patients with mild to moderate hypertension: results of a randomized, double-blind, placebo-controlled study. *American Journal of Hypertension*. 2004;17(5):102A. (P-188)

Chrysant S, Xu J, Chavanu K. Secondary analysis of the distribution blood pressure reductions in subjects receiving olmesartan medoxomil and hydrochlorothiazide in a factorial-design study. 22nd Annual American Society of Hypertension Scientific Meeting, May 2007. [Poster]

Chrysant S, Xu J, Chavanu K. Secondary analysis of the distribution blood pressure reductions in subjects receiving olmesartan medoxomil and hydrochlorothiazide in a factorial-design study. 22nd Annual American Society of Hypertension Scientific Meeting, May 2007. [Abstract]

Chrysant S, Xu J, Dubiel R. Correlation between blood pressure reduction and baseline blood pressure in patients receiving olmesartan medoxomil and/or hydrochlorothiazide in a factorial-design study. 17th Annual European Meeting on Hypertension, June 2007. [Poster]

Chrysant S, Xu J, Dubiel R. Secondary analysis of the distribution blood pressure reductions in patients receiving olmesartan medoxomil and hydrochlorothiazide in a factorial-design study. 17th annual european meeting on hypertension, June 2007. [Poster]

Chrysant SG, Marbury TC, Robinson TD. Antihypertensive efficacy and safety of olmesartan medoxomil compared with amlodipine for mild-to-moderate hypertension. *Journal of Human Hypertension*. 2003;17(6):425-32.

Chrysant SG, Marbury TC, Silfani TN. Use of 24-h ambulatory blood pressure monitoring to assess blood pressure control: a comparison of olmesartan medoxomil and amlodipine besylate. *Blood Pressure Monitoring*. 2006;11:135-141.

Chrysant SG, Neutel JM, Xu J, et al. Distribution of blood pressure response in patients receiving an olmesartan medoxomil-based titration regimen. 22nd annual american society of hypertension scientific meeting, May 2007. [Abstract]

Chrysant SG, Neutel JM, Xu J, et al. Distribution of blood pressure response in patients receiving an olmesartan medoxomil-based titration regimen. 22nd annual american society of hypertension scientific meeting, May 2007. [Poster]

Chrysant SG, Weber MA, Wang A, Hinman DJ. A Factorial Design to Assess the Safety and Efficacy of Olmesartan Medoxomil and Hydrochlorothiazide Combination Therapy. *American Journal of Hypertension*. 2003;16(5):110A.

Chrysant SG, Weber MA, Wang A, Hinman DJ. Evaluation of antihypertensive therapy with the combination of olmesartan medoxomil and hydrochlorothiazide. *American Journal of Hypertension*. 2004; 17(3): 252-259.

AZOR™ (amlodipine and olmesartan medoxomil)  
Formulary Dossier

**OLMESARTAN MEDOXOMIL CLINICAL BIBLIOGRAPHY**  
(continued)

Chrysant SG,, Chrysant GS. Antihypertensive efficacy of olmesartan medoxomil alone and in combination with hydrochlorothiazide. Expert Opin. *Pharmacother*. 2004; 5(3):657-667.

Cook CA, Brown-Jones K. Efficacy of olmesartan medoxomil, a new angiotensin receptor antagonist, in reducing blood pressure in hypertensive African Americans. *Ethnicity and Disease*. 2002;12:S3-127.

Cushman WC, Basile J. Achieving blood pressure goals: why aren't we? *Journal of Clinical Hypertension*. 2006;8(12):865-872.

Cushman WC. Introduction: Hypertension Control in Clinical Practice With a New Standard in Angiotensin II Receptor Blockade. *American Journal of Hypertension*. 2002; 15: 99S-100S.

Daiana Weiss, Dan Sorescu, W. Robert Taylor. Angiotensin II and atherosclerosis. *Am J Cardiology*. 2001; 87(8A): 25C-32C.

Danilo Fliser and Heyrman Haller. Angiotensin subtype 1-receptor antagonists in the treatment of diabetic nephropathy. *Journal of Hypertension* .2001; 19 (suppl 1):S57-S60.

Danser AH, Schuijt MP. Expert Opinion on Investigational Drugs. 2000; 9: 2419-2423.

De Pablos-Velasco PL, Macias-Batista A, Rosas-Rodriguez H, et al. Effects of olmesartan vs. diltiazem on blood pressure and albumin excretion in type 2 diabetic patients with hypertension and microalbuminuria. 15th European Meeting on Hypertension, June 2005.

De Vinuesa SG, Goicoechea M, Kanter J, et al. Effects of olmesartan on insulin resistance syndrome, adipokines and inflammatory parameters in ckd patients. 65th Annual American Society of Hypertension Conference, May 2006. [Abstract]

De Vinuesa SG, Goicoechea M, Kanter J, et al. Insulin resistance, inflammatory biomarkers, and adipokines in patients with chronic kidney disease: Effects of angiotensin ii blockade. *J Am Soc Nephrol*. 2006;17(12)(suppl 3):s206-12.

De Zeeuw D, Remuzzi G, Hans-Henrick P, et.al. Proteinuria, a target for renoprotection in patients with type 2 diabetic nephropathy: Lessons from RENAAL. *Kidney International*. 2004;65:2309-2320.

Deerochanawong C, Hosp R, Yigit Z, et al. Olmesartan, monotherapy and combination diuretic therapy, for the treatment of hypertension: Findings from a large cross-regional study. 47th Annual Conference on Cardiovascular Disease Epidemiology & Prevention in association with the Council on Nutrition, Physical Activity & Metabolism (Affiliation to the American Heart Association), March 2007. [Poster]

Del Castillo A, Calvo C, Lopez Paz JE. A short-term retrospective economic evaluation for the treatment of mild to moderate essential hypertension with angiotensin 2 receptor antagonist in Spain. [Abstract]

Destro M, Georgatos J, Cognoni F, et al. Comparative efficacy of valsartan 160 mg od and olmesartan 20 mg od in mild-to-moderate hypertension: Results of 24-hour ambulatory blood pressure monitoring. 65th Annual American Society of Hypertension Conference, May 2006. [Abstract]

Destro M, Scabrosetti R, Vanasia A, et al. Comparative efficacy of valsartan and olmesartan in mild-to-moderate hypertension: results of 24-hour ambulatory blood pressure monitoring. *Advances In Therapy*. 2005;22(1): 32-43

Diego M, Celia F, Fernando J, et al. Effectiveness of olmesartan on pro-inflammatory cytokines in patients with essential hypertension. 22nd Annual American Society of Hypertension Scientific Meeting, May 2007. [Abstract]

AZOR™ (amlodipine and olmesartan medoxomil)  
Formulary Dossier

**OLMESARTAN MEDOXOMIL CLINICAL BIBLIOGRAPHY**  
(continued)

Diego M, Celia F, Luis A, et al. Effectiveness of olmesartan on rigidity of great arteries measured by pulse wave velocity. 22nd Annual American Society of Hypertension Scientific Meeting, May 2007. [Abstract]

Douglas E. Vaughan. Angiotensin, fibrinolysis, and vascular homeostasis. *Am J Cardiology*. 2001; 87(8A): 18C-24C.

Elisaf M. Olmesartan. *Curr Opin Cardiol Pulm Ren Invest Drugs*. 2000; 2:378-383.

Ewald S, Bohm M, Stefenelli T. OLMEBEST-study: Reduction of blood pressure in the treatment of patients with mild-to-moderate essential hypertension-interim analysis of the Austrian sub study. 15th European Meeting on Hypertension, June 2005.

Ewald S, Bohm M, Woittiez AJ. OLMEBEST-study: Reduction of blood pressure in the treatment of patients with mild-to-moderate essential hypertension-interim analysis of the Dutch sub study-. 15th European Meeting on Hypertension, June 2005.

Ewald S, Bohm M. OLMEBEST-study: Reduction of blood pressure in the treatment of patients with mild-to-moderate essential hypertension-the onset of efficacy. 15th European Meeting on Hypertension, June 2005.

Ewald S, Bohm M. OLMEBEST-study: Reduction of blood pressure in the treatment of patients with mild-to-moderate essential hypertension-metaanalysis of the interim results of the sub studies in Germany, United Kingdom, the Netherlands and Austria. 15th European Meeting on Hypertension, June 2005.

Ewald S, Böhm M. OLMEBEST-study: Reduction of blood pressure in the treatment of patients with mild-to-moderate essential hypertension - The onset of efficacy. 65th Annual American Society of Hypertension Conference, May 2006. [Poster]

Ewald S, Heagerty T, Bohm M. An interim analysis of the United Kingdom OLMEBEST sub-study: blood pressure reduction in patients with mild-to-moderate essential hypertension. 65th Annual American Society of Hypertension Conference, May 2006. [Abstract]

Ewald S, Heagerty T, Bohm M. OLMEBEST-study: Reduction of blood pressure in the treatment of patients with mild-to-moderate essential hypertension - The onset of efficacy. 15th European Meeting on Hypertension, June 2005.

Ewald S, Heagerty T, Stefenelli T, et al. OLMEBEST - study: Reduction of blood pressure in the treatment of patients with mild-to-moderate essential hypertension. - Interims analysis of the pooled data. 65th annual american society of hypertension conference, May 2006. [Poster]

Ewald S, Krzesinski JM, Lins R, et al. OLMEBEST-study: Reduction of blood pressure in the treatment of patients with mild to moderate essential hypertension - Analysis of the randomised study period. 22nd Annual American Society of Hypertension Scientific Meeting, May 2007. [Abstract]

Ewald S, Krzesinski JM, Lins R, et al. OLMEBEST-study: Reduction of blood pressure in the treatment of patients with mild to moderate essential hypertension - Analysis of the randomised study period. 22nd Annual American Society of Hypertension Scientific Meeting, May 2007. [Poster]

Ewald S, Nickenig G, Bohm M. OLMEBEST - study: Reduction of blood pressure in the treatment of patients with mild-to-moderate essential hypertension- Interim analysis of the German sub study. 15th European Meeting on Hypertension, June 2005.

Ewald S, Vor Dem Esche J, Mendgen T. Frequency of blood pressure self measurement and success of antihypertensive treatment in a telemonitoring study: The OLMETEL study. 65th Annual American Society of Hypertension Conference, May 2007. [Poster]

AZOR™ (amlodipine and olmesartan medoxomil)  
Formulary Dossier

**OLMESARTAN MEDOXOMIL CLINICAL BIBLIOGRAPHY**  
(continued)

Ewald S, Vor Dem Esche J, Mengden T. Number of home measurements needed to identify responders in the OLMETEL study. 65th Annual American Society of Hypertension Conference, May 2006. [Poster]

Ewald S, Vor Dem Esche J, Mengden T. Number of home measurements needed to identify responders in the OLMETEL study. 15th European Meeting on Hypertension, June 2005.

Ewald S, Vor Dem Esche J, Neikes F, et al. Relationship between frequency of blood pressure self-measurement and blood pressure reduction in OLMETEL: A telemonitoring study. 65th Annual American Society of Hypertension Conference, May 2006. [Abstract]

Ewald S, Vor Dem Esche J, Neikes F, et al. Using home BP measurements to identify responders: The OLMETEL study. 65th Annual American Society of Hypertension Conference, May 2006. [Abstract]

Ewald S, Vor Dem Esche J, Uen S, Neikes F, et al. Relationship between the frequency of blood pressure self-measurement and blood pressure reduction with antihypertensive therapy: Results of the OLMETEL (olmesartan telemonitoring blood pressure) study. *Clinical Drug Investigation*. 26(8):439-446, 2006.

Ewald S. OLMEBEST-study: Reduction of blood pressure in the treatment of patients with mild-to-moderate essential hypertension - The onset of efficacy. 9th International Forum for the Evaluation of Cardiovascular Care, February 2006. [Poster]

Ewald S. OLMEBEST-study: Reduction of blood pressure in the treatment of patients with mild-to-moderate essential hypertension - Metaanalysis of the interim results of the sub studies in Germany, United Kingdom, the Netherlands and Austria. 9th international forum for the evaluation of cardiovascular care, February 2006. [Poster]

Ewald S. OLMEBEST - study: Reduction of blood pressure in the treatment of patients with mild to moderate essential hypertension - Analysis of the randomised study period. 17th Annual European Meeting on Hypertension, June 2007. [Poster]

Ewald S. OLMEPAS - study: Results of the OLMER3b-substudy (olmesartan real life responder rate in Belgium). 17th Annual European Meeting on Hypertension, June 2007. [Poster]

Fabia MJ, Abdilla N, Oltra RM, et al. Factors determinant of antihypertensive activity of the ARB's. A meta-analysis from 24 hr ambulatory blood pressure studies. 65th annual american society of hypertension conference, May 2006. [Abstract]

Ferrario CM, Strawn WB. Role of the renin-angiotensin-aldosterone system and proinflammatory mediators in cardiovascular disease. *American Journal of Cardiology*. 2006;98(1):121-128.

Fliser D, Wagner KK, Loos A, et al. Chronic angiotensin ii receptor blockade reduces (intra) renal vascular resistance in patients with type ii diabetes. *J Am Soc Nephrol*. 2005;16:1135-1140

Fliser D, Buchholz K, Haller H. Antiinflammatory effects of angiotensin ii subtype 1 receptor blockade in hypertensive patients with microinflammation. *Circulation*. 2004;110:1103-1107.

Fliser D, Haller H. Anti-inflammatory effects of angiotensin ii subtype 1-receptor blockade in hypertensive patients with micro-inflammation. *Journal of Hypertension*. 2004;22(s2):s284. [Abstract 8a.4]

Fliser F. Olmesartan Medoxomil: A Viewpoint by Danilo Fliser. *Drugs*. 2002; 62(9):1354-6.

Fogari R, Mugellini A, Zoppi A, et al. Telmisartan hydrochlorothiazide combination is more effective than olmesartan hydrochlorothiazide combination in the treatment of moderate hypertension. 17th annual european meeting on hypertension, June 2007. [Poster]

AZOR™ (amlodipine and olmesartan medoxomil)  
Formulary Dossier

**OLMESARTAN MEDOXOMIL CLINICAL BIBLIOGRAPHY**  
(continued)

Fogari R, Preti P, Destro M, et al. Efficacy and tolerability of delapril-manidipine combination vs. olmesartan-hctz fixed combination in elderly diabetic hypertensive patients. 21st Scientific Meeting of International Society of Hypertension (ISH), October 2006. [Abstract]

Gavras I, H Gavras. Angiotensin II as a Cardiovascular Risk Factor. *J Hum HTN*. 2002; 16: S2-S6.

Giles T, Oparil S, Silfani T, et al. Comparison of ascending doses of olmesartan medoxomil (o), losartan potassium (l) and valsartan (v) in patients (pts) with essential hypertension (htn). *AJH*. (Abstract/posters: antihypertensive drugs and pharmacology) 2005;18(5): part 2.

Giles T, Oparil S, Silfani T, et al. Comparison of increasing doses of olmesartan medoxomil, losartan potassium, and valsartan in patients with essential hypertension. *J Clin Hypertens*. 2007;9(3):187-195.

Giles TD, Oparil S, Silfani T, et al. Comparison of ascending doses of olmesartan medoxomil, losartan potassium and valsartan in patients with stage 2 essential hypertension. *Am J Hypertens*. 2005;18(5):59a. [Abstract]

Giles TD, Robinson TD. Reduction of Systolic Blood Pressure and Pulse Pressure with Olmesartan Medoxomil. *American Journal of Hypertension*. 2003; 16(5):116A.

Giles TD, Robinson TD. Effects of olmesartan medoxomil on systolic blood pressure and pulse pressure in the management of hypertension. *American Journal of Hypertension*. 2004;17(8):690-695.

Goette A, Breithardt G, Fetsch T, et al. Angiotensin ii antagonist in paroxysmal atrial fibrillation (antipaf) trial : Rationale and study design. *Clin Drug Investig*. 2007;27(10):697-705.

Gonzalez Albarran O, Lahera M, Calvo S, et al. Effects of olmesartan on insulin resistance and inflammatory cytokines in hypertensive patients with metabolic syndrome. 17th Annual European Meeting on Hypertension, June 2007. [Poster]

Gordeon T, McInnes. Clinical potential: angiotensin converting enzyme inhibitor or angiotensin antagonist. *Journal of Hypertension*. 2001; 19 (suppl 1):S61-S67.

Greathouse M. A review of olmesartan medoxomil monotherapy: antihypertensive efficacy similar to that of other angiotensin II receptor blocker/hydrochlorothiazide combinations? *Congestive Heart Failure*. 2002; 8(6):313-20.

Greathouse M. Olmesartan medoxomil combined with hydrochlorothiazide for the treatment of hypertension. *Vascular Health and Risk Management*. 2006;2(4):401-409.

Greaves P. Urinary tract. In, *Histopathology of Preclinical Toxicity Study: Interpretation and Relevance in Drug Safety Evaluation*. Elsevier (New York) 1990; 497-583.

Grossman E, Elkayam A, Baron T, et al. The efficacy of add-on olmesartan in uncontrolled hypertension. 17th Annual European Meeting on Hypertension, June 2007. [Poster]

Haller H, Viberti GC, Mimran A, et al. Preventing microalbuminuria in patients with diabetes: Rationale and design of the randomised olmesartan and diabetes microalbuminuria prevention (roadmap) study. *J Hypertens* 2006;24:403-408.

Hans R. Brunner and Jurg Nussberger. Relevance of clinical pharmacological models for the evaluation of therapeutic dose range of an AT1 receptor antagonist. *Journal of Hypertension*. 2001; 19 (suppl 1):S15-S20.

AZOR™ (amlodipine and olmesartan medoxomil)  
Formulary Dossier

**OLMESARTAN MEDOXOMIL CLINICAL BIBLIOGRAPHY**  
(continued)

- Hansson L. Hypertension Management in 2002: Where Have We Been? Where Might We Be Going? *American Journal of Hypertension*. 2002; 15: 101S-107S.
- Hassler C, Forclaz A, Maillard M, et.al. 24h blockade of the renin-angiotensin system by once daily doses of olmesartan medoxomil given alone or in association with an ace-inhibitor to healthy men. *Journal of hypertension*. 2004;22(s2):s249 [Abstract p2.347]
- Hassler C. Sustained 24-hour blockade of the renin-angiotensin system: A high dose of a long-acting blocker is as effective as a lower dose combined with an angiotensin-converting enzyme inhibitor. *Clin Pharmacol Ther*. 2005;78(5):501-7.
- Haworth S, Bathala M, Lee J, et al. A fixed-dose combination of olmesartan medoxomil and amlodipine besylate is bioequivalent to free combination of the agents. 17th Annual European Meeting on Hypertension, June 2007. [Poster]
- Haworth S, Heyrman R, Bathala M, et al. The bioavailability of a fixed-dose combination of olmesartan medoxomil and amlodipine besylate is unaffected by food. 17th Annual European Meeting on Hypertension, June 2007. [Poster]
- Hazan L, Kelfkens Y, Rodriguez O, et al. Olmesartan medoxomil safely treats hypertension in children and adolescents: Preliminary results of a blinded dose-ranging study. 22nd Annual American Society of Hypertension Scientific Meeting, May 2007. [Abstract]
- Hazan L, Kelfkens Y, Rodriguez O, et al. Olmesartan medoxomil safely treats hypertension in children and adolescents: Preliminary results of a blinded dose-ranging study. 22nd Annual American Society of Hypertension Scientific Meeting, May 2007. [Poster]
- Heagerty AM, Mallion JM. Effect of age on olmesartan medoxomil plus hydrochlorothiazide. 22nd Annual American Society of Hypertension Scientific Meeting, May 2007. [Abstract]
- Heagerty AM, Mallion JM. Effect of age on olmesartan medoxomil plus hydrochlorothiazide. 22nd Annual American Society of Hypertension Scientific Meeting, May 2007. [Poster]
- Heagerty T, Ewald S, Boehm M. OLMEBEST - study: Reduction of blood pressure in the treatment of patients with mild to moderate essential hypertension. - Interims analysis of the British part. 65th Annual American Society of Hypertension Conference, May 2006. [Abstract]
- Heyrman WH, Shahinfar S, Carides GW, et.al. Losartan reduces the costs associated with diabetic end-stage renal disease. *Diabetes Care*. 2003. 26 (4) 683-687.
- Hermida R, Calvo C, Ayala D, et al. Comparison of the efficacy of morning versus evening administration of olmesartan in essential hypertension. 17th Annual European Meeting on Hypertension, June 2007. [Poster]
- Hermida RC, Smolensky MH, Portaluppi F. Comparison of the efficacy of morning versus evening administration of olmesartan in uncomplicated essential hypertension. 22nd Annual American Society of Hypertension Scientific Meeting, May 2007. [Abstract]
- Hidaka H, Kokubu S, Nakazawa T, et al. New angiotensin ii type 1 receptor blocker olmesartan improves portal hypertension in patients with cirrhosis. *Hepatology Research*. April 2007.
- Hirai A, Ito K. Clinical evaluation of an angiotensin ii receptor antagonist in essential hypertensive patients. *Journal of Clinical Therapeutics and Medicine*. 2004;20(2):161(77)-162(78).
- Horinaka S, Yagi H, Hara H, et al. Olmesartan medoxomil improves cardio ankle vascular index. 21st Scientific Meeting of International Society of Hypertension (ISH), October 2006. [Abstract]

AZOR™ (amlodipine and olmesartan medoxomil)  
Formulary Dossier

**OLMESARTAN MEDOXOMIL CLINICAL BIBLIOGRAPHY**  
(continued)

Huber M, Bolbrinker J, Kreutz R. Pharmacokinetics and safety of olmesartan medoxomil in combination with glibenclamide in healthy volunteers. *Clin Exp Hypertens*. 2006;28(7):631-43.

Hussar DA. New Drugs 2003, part I. Nursing. 2003; 33(2):57-62.

Ichikawa S, Takayama Y. Long-term treatment by a novel angiotensin II (All) receptor antagonist-CS-866 decreases plasma A II level. *J Hypertens*. 2000;18(suppl. 4):S3.

Ichikawa S, Y Takayama. Long-term effects of olmesartan, an Ang II receptor antagonist, on blood pressure and the renin-angiotensin-aldosterone system in hypertensive patients. *Hypertension Research - Clinical Experimental*. 2001; 24(6):641-6.

Igarashi M, Ishikawa Y, Fujita T, et al. The study design of olmesartan mega study to determine the relationship between cardiovascular events and blood pressure control in hypertensive patients in japan (omega study). 21st Scientific Meeting of International Society of Hypertension (ISH), October 2006. [Abstract]

Imai E, Ito S, Haneda M, et al. Olmesartan reducing incidence of endstage renal disease in diabetic nephropathy trial (orient): Rationale and study design. *Hypertens Res*. 2006;29(9):703-9.

Inukai T, Yoshida N, Wakabayashi S, et al. Angiotensin-converting enzyme inhibitors and angiotensin ii receptor blockers effectively and directly potentiate superoxide scavenging by polymorphonuclear leukocytes from patients with type 2 diabetes mellitus. *Am J Med Sci*. 2005;329(5):222-227.

Iseki K, Kohagura K, Tozawa M, et al. Multicenter, randomized, parallel study of angiotensin receptor blockade in chronic hemodialysis patients (octopus). 21st Scientific Meeting of International Society of Hypertension (ISH), October 2006. [Abstract]

Ishii M. Clinical profiles of new angiotensin ii receptor blockers: telmisartan, olmesartan medoxomil, and irbesartan. *Nippon Rinsho*. 2004;62(1):193-202.

Izzo J, Neutel J, Dubiel R, et al. Metabolic effects and safety of hydrochlorothiazide (hctz) in combination with olmesartan medoxomil (olm) in stage 2 systolic hypertension (htn). *AJH*. (Abstract/posters: antihypertensive drugs and pharmacology) 2005;18(5): part 2.

Izzo J, Neutel JM, Dubiel R, et al. Efficacy of olmesartan medoxomil and olmesartan/ hydrochlorothiazide in achieving blood pressure control and normalization in stage 2 systolic hypertension. *Am J Hypertens*. 2005;18(5):64a. [Abstract]

Izzo J, Silfani T, Dubiel R, et al. Efficacy of olmesartan medoxomil (om) and om/hydrochlorothiazide (hctz) in patients with isolated systolic hypertension (ish). 65th Annual American Society of Hypertension Conference, May 2006. [Abstract]

Izzo J, Silfani T, Dubiel R, et al. Efficacy of treating stage 2 systolic hypertension with olmesartan medoxomil and olmesartan medoxomil/hydrochlorothiazide in elderly patients and patients with isolated systolic hypertension. 17th Annual European Meeting on Hypertension, June 2007. [Poster]

Izzo JL, Neutel JM, Silfani T, et al. Efficacy and safety of treating stage 2 systolic hypertension with olmesartan and olmesartan/hctz: Results of an open-label titration study. *J Clin Hypertens* (Greenwich). 2007;9(1):36-44.

Izzo JL, Neutel JM, Silfani T, et al. Titration of hctz to 50 mg daily in individuals with stage 2 systolic hypertension pretreated with an angiotensin receptor blocker. *J Clin Hypertens* (Greenwich). 2007;9(1):45-8.



AZOR™ (amlodipine and olmesartan medoxomil)  
Formulary Dossier

**OLMESARTAN MEDOXOMIL CLINICAL BIBLIOGRAPHY**  
(continued)

Izzo JL, Wingertzahn MA. Blood Pressure Responses to Dose Titration of the Angiotensin Receptor Blocker Olmesartan Medoxomil in Essential Hypertension. *American Journal of Hypertension*. 2003;16(5):119A.

Jones M, Lai YL, Sealey J, et al. Effect of olmesartan, irbesartan, and valsartan on renin-angiotensin-aldosterone system (raas) in healthy normal subjects. 65th Annual American Society of Hypertension Conference, May 2006. [Abstract]

Jones M, Lai YL, Sealey J, et al. Effects of olmesartan medoxomil (olm), irbesartan (irb), and valsartan (val) on the renin-angiotensin-aldosterone system (raas) in healthy normal subjects. World Congress of Cardiology, September 2006. [Abstract]

Jones M, Sealey J, Laragh J. Effects of olmesartan medoxomil (olm) and valsartan (val) on the reninangiotensin-aldosterone system (raas) in healthy normal subjects. *AJH*. (Posters: antihypertensive drugs and pharmacology) 2005;18(5): part 2.

Jones M, Sealey J, Laragh J. Effects of olmesartan medoxomil and valsartan on the reninangiotensin-aldosterone system in healthy normal subjects. *Am J Hypertens*. 2005;18(5):65a. [Abstract]

Jones M, Sealey J, Laragh J. Effects of olmesartan medoxomil, irbesartan, and valsartan on the renin-angiotensin-aldosterone system in healthy normal subjects. 65th Annual American Society of Hypertension Conference, May 2006. [Poster]

Jones MR, Sealey JR, Laragh JH. Effects of angiotensin receptor blockers on ambulatory plasma renin activity in healthy, normal subjects during unrestricted sodium intake. *AJH*. 2007; 20:907-916.

Kawaratani T, K. Püchler, P. Laeis, P.U. Witte. Pharmacokinetics of RNH-6270, the active metabolite of the oral angiotensin II-antagonist CS-866, in healthy, male subjects. *Journal of HTN* 1999; 17(suppl 3).

Kawaratani T, P. Laeis, K. Püchler, P.U. Witte. The effect of an antacid (aluminum magnesium hydroxide) on the pharmacokinetics and safety of the oral angiotensin II-antagonist CS-866 in healthy, male subjects. *Journal of HTN*. 1999; 17(suppl 3): A145.

Kawaratani T, Püchler K, Laeis P, et al. Steady state pharmacokinetics of CS-866, a novel angiotensin II (AT1)-receptor antagonist in elderly and young hypertensive patients. 2nd International Symposium on Angiotensin II Antagonism, 1999 [Abstract #9].

Kazuhiko SONOYAMA. The effect of angiotensin II on inward rectifier K channel in atrial myocytes. *J Yonago Med Ass* 2002; 53: 41-53.

Keith J. Ball, Paul A. Williams, et al. Relative efficacy of an angiotensin antagonist compared with other antihypertensive agents. Olmesartan medoxomil versus antihypertensives. *Journal of Hypertension*. 2001; 19 (suppl 1):S49-S56.

Kereiakes D, Neutel JM, Punzi H, et al. Comparative blood pressure-lowering efficacy of olmesartan/hydrochlorothiazide and benazepril/amlodipine in patients with stage 2 hypertension. 22nd Annual American Society of Hypertension Scientific Meeting, May 2007. [Abstract]

Kereiakes D, Neutel JM, Punzi H, et al. Comparative blood pressure-lowering efficacy of olmesartan/hydrochlorothiazide and benazepril/amlodipine in patients with stage 2 hypertension. 22nd Annual American Society of Hypertension Scientific Meeting, May 2007. [Poster]

Kereiakes D, Neutel JM, Punzi HA, et al. What is the optimal combination therapy for hypertension: rationale and design of a study comparing benazepril/amlodipine with olmesartan medoxomil/hydrochlorothiazide. 22nd Annual American Society of Hypertension Scientific Meeting, May 2007. [Poster]

AZOR™ (amlodipine and olmesartan medoxomil)  
Formulary Dossier

**OLMESARTAN MEDOXOMIL CLINICAL BIBLIOGRAPHY**  
(continued)

Kerekiakes D, Neutel J, Punzi H, et al. Comparative blood pressure-lowering efficacy of olmesartan/hydrochlorothiazide and benazepril/amlodipine in patients with stage 2 hypertension. 17th Annual European Meeting on Hypertension, June 2007. [Poster]

Kikuo Arakawa. Introduction. *Am J Cardiology* April 19 2001; 87(8A): 1-2.

Kitamura N, Takahashi Y, Yamadate S, Asai S. Angiotensin II Receptor Blockers Decreased Blood Glucose Levels: A Longitudinal Survey using Data from Electronic Medical Records. *Cardiovasc Diabetol*. 2007 Sep 29;6(1):26.

Kjeldsen SE, Strand A, Julius S, et al. Mechanism of angiotensin ii type i receptor blocker action in the regression of left ventricular hypertrophy. *J Clin Hypertens*. 2006;8(7):487-92.

Klaus von Bergmann, Petra laeis, et al. Olmesartan medoxomil : influence of age, renal and hepatic function on the pharmacokinetics of olmesartan medoxomil. *Journal of Hypertension*. 2001; 19 (suppl 1):S33-S40.

Klebs S, Trenkwalder P, Samer H, et al. The fixed dose combination of valsartan 160/hctz 25 in hypertensive patients not controlled by monotherapy with olmesartan 40 or olmesartan 20 plus hctz 12.5. 15th European Meeting on Hypertension, June 2005.

Kobayashi N, Fujimori I, Watanabe M, Ikeda T. Real-time monitoring of metabolic reactions by microdialysis in combination with tandem mass spectrometry: hydrolysis of CS-866 in vitro in human and rat plasma, livers, and small intestines. *Anal Biochem* 2000 Dec 15;287(2):272-8.

Kostis JB, Silfani T. The combination of olmesartan medoxomil plus hydrochlorothiazide in subjects with stage 2 hypertension: results of a randomized, double-blind, factorial-design study. *American Journal of Hypertension*. 2004;17(5):114A. (P-223)

Kreutz R, Bolbrinker J, Huber M. Pharmacokinetics of olmesartan medoxomil plus hydrochlorothiazide combination in healthy subjects. *Clinical Drug Investigation*. 2006;26(1):29-34.

Kreutz R, Puchler K, Laeis P. Combination therapy of olmesartan medoxomil with amlodipine or atenolol maintains steady state pharmacokinetics of each drug unchanged. *Journal of Hypertension*. 2004;22(s2):s244. [Abstract p2.329]

Kreutz R, Puchler K, Laeis P. The effect of the combination of the oral angiotensin ii-antagonist olmesartan medoxomil and glibenclamide on pharmacokinetics, safety and tolerability in healthy male subjects. 14th European Society of Hypertension, June 2004. [Abstract]

Krzensinski JM, Scheen AJ. Le medicament du mois: olmesartan medoxomil (belsar ou olmetec). *Rev Med Liege*. 2004;59(10):607-611.

Kupecz D, Berardinelli C. Lower the pressure with a new antihypertensive. *Nurse Practitioner*. 2002; 27(8):42-4.

Kushiro T, Saito J, Sato Y, et al. Discrepancy of actual achievement rate of blood pressure target and physicians' assessment of efficacy in elderly hypertensives. 21st Scientific Meeting of International Society of Hypertension (ISH), October 2006. [Abstract]

AZOR™ (amlodipine and olmesartan medoxomil)  
Formulary Dossier

**OLMESARTAN MEDOXOMIL CLINICAL BIBLIOGRAPHY**  
(continued)

Kushiro T, Saito J, Yamakawa M, et al. Comparative study of blood pressure lowering effects of angiotensin ii receptor blocker (arb) monotherapy and in combination with other antihypertensives. 21st Scientific Meeting of International Society of Hypertension (ISH), October 2006. [Abstract]

Laeis P, K Puchler, K von Bergmann. A Comparative Pharmacokinetic, Safety, and Tolerability Trial of the Oral Angiotensin II-Antagonist Olmesartan Medoxomil in Subjects with Varying Degrees of Renal Impairment and in Healthy Volunteers. *Journal of HTN*. 2001; 19 (suppl 2): P2.147.

Laeis P, K Puchler, K von Bergmann. The Effect of Aluminum Magnesium Hydroxide on the Pharmacokinetics of the Oral Angiotensin II-Antagonist Olmesartan Medoxomil in Healthy Male Volunteers. *Journal of HTN*. 2001; 19 (suppl 2): P2.177.

Laeis P, K Puchler, W Kirch. The Effect of the Oral Angiotensin II-Antagonist Olmesartan Medoxomil on the Safety, Tolerability and Pharmacokinetics of Digoxin in Healthy, Male Subjects. *Journal of HTN*. 2001; 19 (suppl 2): P2.179.

Laeis P, Kawaratani T, Puechler K. Clinical pharmacology profile of CS-866. *Kidney & Blood Pressure Research* 1998, 21: 364.

Laeis P, Nussberger J, Puechler K, et al. Dose-Response Relationship of The Oral Angiotensin II-Antagonist CS-866 in Healthy Male Subjects. *Journal of HTN*. 1999; 17(suppl 3): (poster: P1.168).

Laeis P, Nussberger J, Puechler K, et al. Dose Related Effects of CS-866, A New Oral Angiotensin-II (AT1)-Receptor Antagonist, In Normotensive and Hypertensive Human Subjects. 2nd International Symposium on Angiotensin II Antagonism, 1999 [Abstract No. 10].

Laeis P, Puechler K, Witte PU, Woodward A. Pharmacokinetics of the new oral angiotensin II-antagonist CS-866. 2nd Congress of the European Association for Clinical Pharmacology and Therapeutics (EACPT). *European Journal of Clinical Pharmacology*. 1997; 52: A148.

Laeis P, Puechler K, Witte PU, Woodward A. The pharmacokinetic profile of CS-866. 8th European Meeting on Hypertension. *Journal of Hypertension*, 1997; 15 (Supple. 4); S158.

Lee TH. "Me-Too" Products - Friend of Foe? *New England Journal of Medicine*. 2004; 350(3):211-212.

Levi-Derrick V, Neutel J, Smith D, et al. Treatment of stage 1 and stage 2 hypertension (htn) utilizing an olmesartan medoxomil (om)-based treatment algorithm hypertension. 21st National Conference, American Academy of Nurse Practitioners (AANP), June 2006. [Poster]

Lewis EJ. The Role of Angiotensin II Receptor Blockers in Preventing the Progression of Renal Disease in Patients with Type 2 Diabetes. *American Journal of Hypertension*. 2002; 15: 123S-128S.

Liau CS, Lee CM, Sheu SH, et al. Efficacy and safety of olmesartan in the treatment of mild-to-moderate essential hypertension in chinese patients. *Clinical Drug Investigation*. 2005;25(7):473-479.

Liefde IV, Le MT, Pugsley M, et al. A characterization of the molecular interactions of olmesartan and telmisartan with human angiotensin at1 receptors. 65th Annual American Society of Hypertension Conference, May 2006. [Poster]

Liu D, Hua P, Matsushima N, Li X, et al. Quantitative determination of olmesartan in human plasma and urine by liquid chromatography coupled to tandem mass spectrometry. *J Chromatogr B Analyt Technol Biomed Life Sci*. 2007 Jun 16.

AZOR™ (amlodipine and olmesartan medoxomil)  
Formulary Dossier

**OLMESARTAN MEDOXOMIL CLINICAL BIBLIOGRAPHY**  
(continued)

- Lopez J, Hermida R, Calvo C, et al. Administration-time dependent efficacy of olmesartan on ambulatory blood pressure according to gender. 17th Annual European Meeting on Hypertension, June 2007. [Poster]
- Lopez JE, Hermida RC, Calvo C, et al. Administration-time dependent efficacy of olmesartan on ambulatory blood pressure according to gender. 22nd Annual American Society of Hypertension Scientific Meeting, May 2007. [Abstract]
- Lopez-Farre A, Luque M, Marques M, et al. Olmesartan treatment modified the platelet proteome in hypertensive patients. 17th Annual European Meeting on Hypertension, June 2007. [Poster]
- Ludwig M, Von Buquoy M, Von Petzinger-Kruthoff A, et al. Reproducibility of 3d-ultrasound readings in the assessment of carotid plaque volume in patients with cardiovascular risk. 22nd Annual American Society of Hypertension Scientific Meeting, May 2007. [Poster]
- Mallion JM, Heagerty A, Laeis P. Systolic Blood Pressure Reduction with Olmesartan Medoxomil Versus Nitrendipine in Elderly Patients with Isolated Systolic Hypertension. *J Hypertens*. 2007 Oct;25(10):2168-2177.
- Marbury TC, Silfani T. Use of 24-hour ambulatory blood pressure monitoring to assess antihypertensive efficacy: A comparison of olmesartan medoxomil and amlodipine besylate. *Am J Hypertens*. 2005;18(5):68a. [Abstract]
- Marks E, Smith D, Neutel J, et al. Quality of life improvements in patients achieving blood pressure goal with an olmesartan medoxomil based treatment algorithm. 21st National Conference, American Academy of Nurse Practitioners (AANP), June 2006. [Poster]
- Marques M, Luque M, Tamargo J, et al. Effect of olmesartan treatment on cardiovascular risk markers in hypertensive patients. 17th Annual European Meeting on Hypertension, June 2007. [Poster]
- Martinez-Martin FJ, Macias-Batista A, Rodriguez-Rosas H, et al. Effects of olmesartan on insulin sensitivity in non-diabetic hypertensive patients with metabolic syndrome. 15th European Meeting on Hypertension, June 2005.
- Masonson HN, Punzi HA, Neutel JM, et al. CS-866 (Angiotensin-II Receptor Antagonist): A Double-Blind Study Using Ambulatory Blood Pressure Monitoring in Hypertensive Patients. *American Journal of Hypertension*. 1998; 11: 77A.
- McInnes GT. Olmesartan Medoxomil: A Viewpoint by Gordon T. McInnes. *Drugs*. 2002; 62(9):1354-6.
- McInnes GT. How important is optimal blood pressure control? clinical therapeutics. 2004;26(suppl a):a3-a11.
- Mengden T, Ewald S, Kaufmann S, et al. Telemonitoring of blood pressure self measurement in the olmetel study. *Blood Pressure Monitoring*. 2004;9:1-5.
- Mengden T, Ewald S, Vor Dem Esche J. Frequency of blood pressure self management and success of anti-hypertensive treatment in a telemonitoring study: the OLMETEL study. 15th European Meeting on Hypertension, June 2005.
- Milani RV. Reaching for aggressive blood pressure goals: role of angiotensin receptor blockade in combination therapy. *The American Journal of Managed Care*. 2005;11(suppl 7):s220-7.
- Mire D, Silfani T, Pugsley M. A review of the structural and functional features of olmesartan medoxomil, an angiotensin receptor blocker. *J Cardiovasc Pharmacol*. 2005;46:585-593.

AZOR™ (amlodipine and olmesartan medoxomil)  
Formulary Dossier

**OLMESARTAN MEDOXOMIL CLINICAL BIBLIOGRAPHY**  
(continued)

Miura S, Saku K. Do all angiotensin ii type 1 receptor blockers have the same beneficial effects? *British Journal of Pharmacology*. 2007;151:912–913.

Miyata T, Van Ypersele De Strihou C, et al. Angiotensin II receptor Antagonists and Angiotensin-Converting Enzyme Inhibitors Lower in vitro the Formation of Advanced Glycation End Products: Biochemical Mechanisms. *Journal of the American Society of Nephrology*. 2002; 13(10): 2478-87.

Mizutani N, Fukuta M, Yonemoto T, et al. Autonomic and baroreflex function after olmesartan medoxomil treatment in essential hypertensive patients. 21st Scientific Meeting of International Society of Hypertension (ISH), October 2006. [Abstract]

Moniwa N, Yoshida H, Agata J, et al. Olmesartan improves glucose metabolism and insulin sensitivity in patients with essential hypertension more than does a calcium channel blocker. 21st Scientific Meeting of International Society of Hypertension (ISH), October 2006. [Abstract]

Morimoto S, Okaishi K, Nakahashi T, et al. Comparisons of efficacies of olmesartan and telmisartan on elderly hypertensive subjects. 21st Scientific Meeting of International Society of Hypertension (ISH), October 2006. [Abstract]

Naya M, Tsukamoto T, Morita K, Katoh C, Furumoto T, Fujii S, Tamaki N, Tsutsui H. Olmesartan, but not Amlodipine, Improves Endothelium-Dependent Coronary Dilation in Hypertensive Patients. *J Am Coll Cardiol*. 2007 Sep 18;50(12):1144-9.

Neutel J, Lee Y, Silfani T. Olmesartan medoxomil (olm) reduces blood pressure (bp) load and improves goal rates more effectively than losartan potassium (los) and valsartan. *AJH (Antihypertensive Drugs and Pharmacology)*. 2005;18(5): part 2.

Neutel J, Chrysant S, Dubiel R, et al. Efficacy of treating stage 2 systolic hypertension with olmesartan medoxomil and olmesartan medoxomil/hydrochlorothiazide in patients stratified by gender and body mass index. 17th Annual European Meeting on Hypertension, June 2007. [Poster]

Neutel J, Dubiel R, Walker F, et al. Efficacy of olmesartan medoxomil (om) and om/hctz in achieving blood pressure goals in patients with stage 2 systolic hypertension: Effect of gender. 65th Annual American Society of Hypertension Conference, May 2006. [Abstract]

Neutel J, Dubiel R, Walker F, et al. Efficacy of olmesartan medoxomil and olmesartan medoxomil/hydrochlorothiazide in achieving blood pressure control and normalization in stage 2 systolic hypertension: Effect of gender. 65th Annual American Society of Hypertension Conference, May 2006. [Poster]

Neutel J, Dubiel R, Walker F, et al. Efficacy of olmesartan medoxomil and om/hctz in achieving blood pressure goals in patients with stage 2 systolic hypertension: Effect of age. 65th annual american society of hypertension conference, May 2006. [Abstract]

Neutel J, Dubiel R, Walker F, et al. Efficacy of olmesartan medoxomil and olmesartan medoxomil/hydrochlorothiazide in achieving blood pressure goals in patients with stage 2 systolic hypertension: Effect of age. 65th Annual American Society of Hypertension Conference, May 2006. [Poster]

Neutel J, Kereiakes D, Punzi H, et al. Attainment of blood pressure goals in patients with stage 2 hypertension treated with olmesartan medoxomil/hydrochlorothiazide or benazepril/amlodipine. 17th Annual European Meeting on Hypertension, June 2007. [Poster]

Neutel J, Lee Y, Walker F. Efficacy of olmesartan medoxomil (olm), losartan potassium (los) and valsartan in diabetic patients. *AJH (antihypertensive drugs and pharmacology)*. 2005;18(5): part 2. [Poster]

AZOR™ (amlodipine and olmesartan medoxomil)  
Formulary Dossier

**OLMESARTAN MEDOXOMIL CLINICAL BIBLIOGRAPHY**  
(continued)

- Neutel J, Smith D, Silfani T, et al. Treatment of stage 1 and stage 2 hypertension (htn) utilizing an olmesartan medoxomil (om)-based treatment algorithm hypertension. *AJH* (antihypertensive drugs and pharmacology). 2005;18(5): part 2. [Poster]
- Neutel JM, Kereiakes D, Punzi HA, et al. Attainment of blood pressure goals in patients with stage 2 hypertension treated with olmesartan medoxomil/hydrochlorothiazide or benazepril + amlodipine. 22nd Annual American Society of Hypertension Scientific Meeting, May 2007. [Abstract]
- Neutel JM, Kereiakes D, Punzi HA, et al. Attainment of blood pressure goals in patients with stage 2 hypertension treated with olmesartan medoxomil/hydrochlorothiazide or benazepril + amlodipine. 22nd Annual American Society of Hypertension Scientific Meeting, May 2007. [Poster]
- Neutel JM, Kereiakes D, Punzi HA, et al. What is the optimal combination therapy for hypertension: Rationale and design of a study comparing benazepril/amlodipine with olmesartan medoxomil/hydrochlorothiazide. 22nd annual american society of hypertension scientific meeting, May 2007. [Abstract]
- Neutel JM, Smith D, Silfani T, et al. Achievement of blood pressure goals using an angiotensin receptor blocker-based regimen. *Am J Hypertens* 2005;18(5):70a. [Abstract]
- Neutel JM, Smith DH, Silfani TN, et al. Effects of a structured treatment algorithm on blood pressure goal rates in both stage 1 and stage 2 hypertension. *J Hum Hypertens*. 2006;20:255-262
- Neutel JM, Smith DHG, Weber MA, Wang AC, Masonson HN. Use of an Olmesartan Medoxomil-based treatment algorithm for hypertension control. *The Journal of Clinical Hypertension*. 2004; 6:168-174.
- Neutel JM. Clinical studies of CS-866, the newest angiotensin II receptor antagonist. *American Journal of Cardiology*. 2001;87(8A):37C-43C.
- Nissen S et al. Effect of antihypertensive agents on cardiovascular events in patients with coronary disease and normal blood pressure the CAMELOT study: A randomized controlled trial. *JAMA*. 2004;292:2217-2226.
- Norwood D, Branch E, Smith B, Honeywell M. Olmesartan Medoxomil for Hypertension: a clinical review. *P&T*. 2002; 27(12):611-617.
- Nussberger J, Koike H. Antagonizing the angiotensin ii subtype i receptor: A focus on olmesartan medoxomil. *Clinical Therapeutics*. 2004;26(suppl a):a12-a20.
- Ogihara T. Olmesartan Medoxomil: A Viewpoint by Toshio Ogihara. *Drugs*. 2002; 62(9):1354-6.
- Oltra PB, Almenar BC, Torres JNY. Therapeutic interchange standardization for angiotensin ii receptor antagonists in the treatment of hypertension in the hospital setting. *Farmacia Hospitalaria*. 2005;29(2):104-12.
- Onai Y, Iwata O, Okahara M, et al. Blood pressure-lowering effect of angiotensin ii receptor blocker, olmesartan medoxomil, and its relationship with the plasma renin activity (pra) and urinary na excretion rate in hypertensive patients. 21st Scientific Meeting of International Society of Hypertension (ISH), October 2006. [Abstract]
- Oparil S, Masonson HN. Antihypertensive Efficacy of Olmesartan Medoxomil in Hypertensive Men Compared with Hypertensive Women. *American Journal of Hypertension*. 2003;16(5):125A.
- Oparil S, Silfani TN, Walker JF. Role of angiotensin receptor blockers as monotherapy in reaching blood pressure goals. *AJH*. 2005;18:287-294.

AZOR™ (amlodipine and olmesartan medoxomil)  
Formulary Dossier

**OLMESARTAN MEDOXOMIL CLINICAL BIBLIOGRAPHY**  
(continued)

Oparil S, Walker JF. Blood pressure goal attainment: comparative efficacy of olmesartan medoxomil, losartan potassium, valsartan, and irbesartan. *American Journal of Hypertension*. 2004;17(5):118A-119A. (P-235)

Oparil S, Williams D, Chrysant SG, et al. Comparative Efficacy of Olmesartan, Losartan, Valsartan, and Irbesartan in the control of Essential Hypertension. *J Clin Hypertens*. 2001; 5:283-91.

Oparil S. Are there meaningful differences in blood pressure control with current antihypertensive agents? *Am Journal of HTN*. 2002; 15:14S-21S.

Oparil S. Comparative antihypertensive efficacy of olmesartan: comparison with other angiotensin II receptor antagonists. *J Hum HTN*. 2002; 16 (Suppl 2): S17-S23.

Pepine C. What is the optimal blood pressure and drug therapy for patients with coronary artery disease? *JAMA*. 200;292:2271-2273

Porterl E, Rodella L, Rizzoni D, et.al. Effects of olmesartan and enalapril on vascular structure and endothelial function of spontaneously hypertensive rats. *Journal of Hypertension*. 200;22(s2):s153. [Abstract p6c.3]

Preti P, Destro M, Lazzari P, et al. Efficacy and tolerability of delapril-manidipine combination vs. olmesartan-hctz fixed combination in elderly diabetic hypertensive patients. 65th Annual American Society of Hypertension Conference, May 2006. [Abstract]

Puchler K, J Nussberger, PU Witte, P Laeis, et al. Pharmacokinetic, Safety and Tolerability Studies of the Oral Angiotensin II-Antagonist Olmesartan Medoxomil in Young, Elderly and Very Elderly Patients with Hypertension. *Journal of HTN*. 2001; 19 (suppl 2): P2.180.

Puchler K, Laeis P, Guenther A, et al. Dose-Finding Studies with CS-866, a new oral antihypertensive agent. *Kidney & Blood Pressure Research* 1998, 21: 374.

Puchler K, P Laeis, KO Stumpe. A Comparison of the Efficacy and Safety of the Oral Angiotensin II-Antagonist Olmesartan Medoxomil with those of Atenolol in patients with Moderate to Severe Hypertension under Continuous Treatment with Hydrochlorothiazide. *Journal of HTN*. 2001; 19 (suppl 2): P2.175.

Puchler K, P Laeis, KO Stumpe. Dose Related Differences in Blood Pressure Response Rate are not Related to Adverse Event Incidence. *Journal of HTN*. 2001; (poster: P1.191).

Puchler K, P Laeis, W Kirch. The Effect of the Oral Angiotensin II-Antagonist Olmesartan Medoxomil on the Pharmacodynamics, Pharmacokinetics and Safety of Warfarin in Healthy, Male Subjects. *Journal of HTN*. 2001; 19 (suppl 2): P2.178.

Püchler K, P. Laeis, T. Kawaratani, et al. The effect of the combination of the oral angiotensin II-antagonist CS-866 and warfarin on pharmacodynamics, pharmacokinetics and safety in healthy, male subjects. *Journal of HTN* 1999; 17(suppl 3): A 271.

Puchler K, Petra Leis, Klaus O. Stumpe. Blood pressure response, but not adverse event incidence, correlates with dose of angiotensin antagonist. *Journal of Hypertension*. 2001; 19 (suppl 1):S41-S48.

Puechler K, Laeis P, Guenther A, et al. Safety, tolerability and efficacy of the new oral angiotensin II (AT1)-receptor antagonist in patients with mild to moderate hypertension. 2nd International Symposium on Angiotensin II Antagonism, 1999 [Abstract No. 11].

AZOR™ (amlodipine and olmesartan medoxomil)  
Formulary Dossier

**OLMESARTAN MEDOXOMIL CLINICAL BIBLIOGRAPHY**  
(continued)

- Puechler K, Laeis P, Witte PU, Fleck C. Antihypertensive efficacy of CS-866, a novel angiotensin II-antagonist. 8th European Meeting on Hypertension *Journal of Hypertension*, 1997; 15 (Supple. 4); S185.
- Puechler K, Nussberger J, Laeis P, Witte PU, et al. Blood pressure and endocrine effects of single doses of CS-866, a novel angiotensin II-antagonist, in salt restricted hypertensive patients. *Journal of Hypertension*. 1997; 15: 1809-1812.
- Puechler K, Nussberger J, Laeis P, Witte PU, et al. Dose response relationship of the new oral angiotensin II antagonist CS-866. 2nd Congress of the European Association for Clinical Pharmacology and Therapeutics (EACPT). *European Journal of Clinical Pharmacology*. 1997; 52: A150.
- Puechler K, Nussberger J, Laeis P, Witte, et al. Angiotensin II-Antagonists CS-866 in hypertensive patients. 8th European Meeting on Hypertension. *Journal of Hypertension*, 1997; 15 (Supple. 4); S115.
- Pugsley MK, Gupta S. Olmesartan, an angiotensin-ii (at-ii) receptor blocker, dose-dependently reduces both blood pressure and renal damage in pre-clinical and clinical studies of hypertension associated with type-ii diabetes. Forest Research Institute. Jersey City, NJ. [Abstract/Poster]
- Pugsley MK, Mire DE, Silfani TN, et al. The dose-dependent antihypertensive actions of olmesartan medoxomil, an angiotensin ii (at-ii) receptor blocker. *American Journal of Hypertension*. 2004;17(5):121a.
- Pugsley MK, Mire DE, Silfani TN, Gupta S. The dose-dependent antihypertensive actions of olmesartan medoxomil, an angiotensin II (AT-II) receptor blocker. *American Journal of Hypertension*. 2004 ;17(5):121A. (P-242)
- Punzi H, Kereiakes D, Neutel JM, et al. Effect of olmesartan medoxomil and benazepril on plasma renin activity (pra) in patients with stage 2 hypertension. 22nd Annual American Society of Hypertension Scientific Meeting, May 2007. [Abstract]
- Punzi H, Kereiakes D, Neutel JM, et al. Effect of olmesartan medoxomil and benazepril on plasma renin activity (pra) in patients with stage 2 hypertension. 22nd Annual American Society of Hypertension Scientific Meeting, May 2007. [Poster]
- Quan A, Chavanu K, Merkel J. A review of the efficacy of fixed-dose combinations olmesartan medoxomil/hydrochlorothiazide and amlodipine besylate/benazepril in factorial design studies. *Am J Cardiovasc Drugs*. 2006;6(2):103-113.
- Ram CVS, Silfani T. A comparative analysis of the efficacy of olmesartan medoxomil (O)/hydrochlorothiazide (HCTZ), valsartan (V)/HCTZ, irbesartan (I)/HCTZ and telmisartan (T)/HCTZ combinations. *American Journal of Hypertension*. 2004 ;17(5):121A. (P-243)
- Ram CVS, Silfani T. A comparative analysis of the efficacy of olmesartan medoxomil (o)/hydrochlorothiazide (hctz), valsartan (v)/hctz, irbesartan (i)/hctz and telmisartan (t)/hctz combinations. *American Journal of Hypertension*. 2004;17(5):121a. (p-243)
- Resnick LM, Catanzaro D, Sealey JE, et al. Acute vascular effects of the angiotensin ii receptor antagonist olmesartan in normal subjects: Relation to the renin-aldosterone system. *American Journal of Hypertension*. 2004;17(3):203-208.
- Resnick LM, Catanzaro D, Sealey JE, et al. Ionic aspects of acute angiotensin receptor blockade in normal subjects. *AJH*. 2004;17(5): part 2. [Abstract]
- Resnick LM, Catanzaro D, Sealey JE, Laragh JH. Acute vascular effects of the angiotensin II receptor antagonist olmesartan in normal subjects: relation to the renin-aldosterone system. *American Journal of Hypertension*. 2004; 17(3):203-208.



AZOR™ (amlodipine and olmesartan medoxomil)  
Formulary Dossier

**OLMESARTAN MEDOXOMIL CLINICAL BIBLIOGRAPHY**  
(continued)

Resnick LM, Gupta R. Linkage between altered rbc free magnesium and plasma membrane and circulating lipids in hypertension: A 1hand 31p-nmr spectroscopy study. *AJH*. 2004;17(5): part 2.

Reynolds MW, Sercus B, Claflin J, et al. Angiotensin ii receptor blockers (arb), hypertensive treatments and the risk of myocardial infarction: A systematic review of the literature. International Society for Pharmaceutical Engineering (ISPE). [Poster]

Rohatagi S, Carrothers Tt, Kshirsagar S, et al. Evaluation of population pharmacokinetics after coadministration of olmesartan medoxomil and amlodipine besylate. 17th Annual European Meeting on Hypertension, June 2007. [Poster]

Rohatagi S, Haworth S, Heyrman R, et al. Low and high doses of a fixed-dose combination of olmesartan medoxomil and amlodipine besylate are bioequivalent to free combination of the agents. 17th Annual European Meeting on Hypertension, June 2007. [Poster]

Rohatagi S, Haworth S, Heyrman R, et al. Safety and tolerability of the combination of olmesartan medoxomil and amlodipine besylate. 17th Annual European Meeting on Hypertension, June 2007. [Poster]

Rohatagi S, Haworth S, Salazar D, et al. Olmesartan medoxomil and amlodipine besylate show dose-proportional drug exposure when used in combination in healthy volunteers. 17th Annual European Meeting on Hypertension, June 2007. [Poster]

Rosen A, Karter A, Liu J, et al. Use of angiotensin converting enzyme inhibitors and angiotension receptor blockers in high risk clinical and ethnic groups with diabetes. *Journal of Gen Internal Medicine*. 2004;19:669-675.

Rosendorff C, Atlas SA, Almendral JL. Angiotensin receptor blockade and cardiovascular hypertrophy Results from the echocardiographic study of the effect on cardiovascular hypertrophy of olmesartan (the ECHO study). 21st Scientific Meeting of International Society of Hypertension (ISH), October 2006. [Abstract]

Rothermund L, Holleck F, Holleck S, et al. AT1 receptor blockade with olmesartan confers striking protective effects on cardio-renal damage in hypertension due to renal mass reduction. 65th Annual American Society of Hypertension Conference, May 2006. [Abstract]

Ruilope LM, Brunner HR. Combined analysis of the efficacy of olmesartan/hydrochlorothiazide. 22nd Annual American Society of Hypertension Scientific Meeting, May 2007. [Abstract]

Ruilope LM, Brunner HR. Combined analysis of the efficacy of olmesartan/hydrochlorothiazide. 22nd Annual American Society of Hypertension Scientific Meeting, May 2007. [Poster]

Rump LC, Ambrosioni E, Burnier M, et al. Initial combination therapy with olmesartan/ hydrochlorothiazide in moderate-to-severe hypertension. *Journal of Human Hypertension*. 2006 ;20(4):299-301.

Rump LC, Ambrosioni E, Burnier M, et.al. Initial combination therapy with olmesartan/hydrochlorothiazide shows superior blood pressure control rate in moderate to severe hypertension. *Journal of Hypertension*. 2004;22(s2):s381 [Abstract p3.324]

Sada T, Mizuno M. Pharmacological profiles and clinical effects of olmesartan medoxomil, A novel angiotension ii receptor blocker. *Folia Pharmacol. JPN*. 2004;124: 257-269.

Saito J, Kushiro T, Sato Y, et al. Efficacy and safety of angiotensin ii receptor blocker in elderly isolated systolic hypertensive patients. 21st Scientific Meeting of International Society of Hypertension (ISH), October 2006 [Abstract]

Saito J, Kushiro T, Yamakawa M, et al. Olmesartan medoxomil has rapid onset of antihypertensive action when given alone, in combination, and to patients switched from other antihypertensive agents. 21st Scientific Meeting of International Society of Hypertension (ISH), October 2006. [Abstract]

AZOR™ (amlodipine and olmesartan medoxomil)  
Formulary Dossier

**OLMESARTAN MEDOXOMIL CLINICAL BIBLIOGRAPHY**  
(continued)

- Salazar D, Lee J, Senouda M, et al. Lack of pharmacokinetic drug interaction between olmesartan medoxomil and amlodipine besylate during coadministration. 17th Annual European Meeting on Hypertension, June 2007. [Poster]
- Saunders E, Neutel J, Bailey W, et al. Efficacy of olmesartan medoxomil and om/hctz in achieving blood pressure goals in patients with stage 2 systolic hypertension: Effect in african americans. 65th Annual American Society of Hypertension Conference, May 2006. [Abstract]
- Saunders E, Neutel JM, Bailey W, et al. Efficacy of olmesartan medoxomil and olmesartan/hydrochlorothiazide in achieving blood pressure control and normalization in stage 2 systolic hypertension. 65th Annual American Society of Hypertension Conference, May 2006. [Poster]
- Scheen AJ. Prevention of type 2 diabetes mellitus through inhibition of the renin-angiotensin system. *Dugs* 2004; 64(22): 2537-2565.
- Schieffer B, Bunte C. Comparative effects of at1-antagonism and angiotension-converting enzyme inhibition on markers of inflammation and platelet aggregation in patients with coronary artery disease. *Journal of the American College of Cardiology*. 2004;44(2):362-368.
- Schiffrin EL. Beyond Blood Pressure: The Endothelium and Atherosclerosis Progression. *American Journal of Hypertension*. 2002; 12: 115S-122S.
- Schmidt AC, Bramlage P, Lichtenthal A, et al. Socio-economic status and the therapeutic effectiveness of antihypertensive treatment – The design of the LEO study. *Curr Med Res Opin* . 2007;23(8)
- Scholze J, Ewald S. Olmesartan medoxomil for the treatment of hypertension under daily-practice conditions – Results of a post-authorization study. 9th International Forum for the Evaluation of Cardiovascular Care, February 2006. [Poster]
- Schwocho LR, Masonson HN. Pharmacokinetics of CS-866, a new angiotensin II receptor blocker, in healthy subjects. *J Clin Pharmacol*. 2001; 41(5):515-27.
- Seki S, Mochizuki S. Olmesartan attenuates inflammation in essential hypertensive patients. 17th Annual European Meeting on Hypertension, June 2007. [Poster]
- Sellin L, Stegbauer J, Laeis P, et al. Adding hydrochlorothiazide to olmesartan dose dependently improves 24-h blood pressure and response rates in mild to-moderate hypertension. *Journal of Hypertension*. 2005;23(11):2083-92.
- Shuji M, Megumi S, Ryuji M, et al. Antihypertensive efficacy and cerebral hemodynamic advantage of olmesartan medoxomil compared with amlodipine in post-stroke patients. 21st Scientific Meeting of International Society of Hypertension (ISH), October 2006. [Poster]
- Simko F, Simko J. The potential role of nitric oxide in the hypertrophic growth of the left ventricle. *Physiologic Res*. 2000;49:37-46.
- Simons WR. Comparative Cost Effectiveness of Angiotensin II Receptor Blockers in a US Managed Care Setting: Olmesartan Medoxomil Compared with Losartan, Valsartan, and Irbesartan. *Pharmacoeconomics*. 2003; 21(1): 61-74.
- Smith D, Bailey W, Jones M, et al. Efficacy of an olmesartan medoxomil-based treatment algorithm in african american patients with hypertension. 65th Annual American Society of Hypertension Conference, May 2006. [Abstract]

AZOR™ (amlodipine and olmesartan medoxomil)  
Formulary Dossier

**OLMESARTAN MEDOXOMIL CLINICAL BIBLIOGRAPHY**  
(continued)

- Smith D, Bailey W, Jones M, et al. Treatment of hypertension in blacks utilizing an olmesartan medoxomil-based algorithm. 65th annual american society of hypertension conference, May 2006. [Poster]
- Smith D, Neutel J, Silfani T, et al. Quality of life improvements in patients achieving blood pressure goal with an olmesartan medoxomil-Based treatment algorithm. 65th Annual American Society of Hypertension Conference, May 2006. [Abstract]
- Smith D, Neutel J, silfani t, et al. Quality of life improvements in patients achieving blood pressure goal with an olmesartan medoxomil-based treatment algorithm. 17th Annual European Meeting on Hypertension, June 2007 [Poster]
- Smith D, Neutel J, Silfani T, et al. Quality of life in patients achieving blood pressure goal with an olmesartan medoxomil-based treatment algorithm. 65th Annual American Society of Hypertension Conference, May 2006. [Poster]
- Smith D, Neutel J, Silfani T, et al. Time to achieve blood pressure goal with an olmesartan medoxomil-Based treatment algorithm. 65th Annual American Society of Hypertension Conference, May 2006. [Poster]
- Smith D, Neutel J, Silfani T, et al. Time to achieve blood pressure goal with an olmesartan medoxomil-based treatment algorithm. 65th Annual American Society of Hypertension Conference, May 2006. [Abstract]
- Smith D, Neutel J, Silfani T, et al. Time to achieve blood pressure goal with an olmesartan medoxomil-based treatment algorithm. 17th Annual European Meeting on Hypertension, June 2007. [Poster]
- Smith David H.G. Strategies to Meet Lower Blood Pressure Goals With a New Standard in Angiotensin II Receptor Blockade. *American Journal of Hypertension*. 2002; 15: 108S-114S.
- Smith DH, Dubiel R, Jones M. Use of 24-hour ambulatory blood pressure monitoring to assess antihypertensive efficacy a comparison of olmesartan medoxomil, losartan potassium, valsartan, and irbesartan. *Am J Cardiovasc Drugs*. 2005; 5(1):41-50.
- Smith DHG, Dubiel R, Jones, M. Use of 24-hour ambulatory blood pressure monitoring to assess antihypertensive efficacy. A comparison of olmesartan medoxomil, losartan potassium, valsartan, and irbesartan. *Am J Cardiovasc Drugs* 2005; 5: 41-50.
- Smith R, Averill D, Brosnihan KB, et al. Effects of antihypertensive therapy on insulin sensitivity leptin values. 22nd Annual American Society of Hypertension Scientific Meeting, May 2007. [Poster]
- Smith R, Brosnihan KB, Owen J. Cross talk of the angiotensin at1 receptor and the thromboxane receptor. 22nd Annual American Society of Hypertension Scientific Meeting, May 2007. [Abstract]
- Smith R, Brosnihan KB, Owen J. Cross talk of the angiotensin at1 receptor and the thromboxane receptor. 22nd Annual American Society of Hypertension Scientific Meeting, May 2007. [Poster]
- Smith R, Ferrario C, Walker F. Number of antihypertensive agents required to achieve goal blood pressure with an olmesartan medoxomil vs atenolol based regimen. 17th Annual European Meeting on Hypertension, June 2007. [Poster]
- Smith R, Walker F, Ferrario C. Number of antihypertensive agents required to achieve goal blood pressure with an olmesartan medoxomil vs atenolol based regimen. 22nd Annual American Society of Hypertension Scientific Meeting, May 2007. [Poster]
- Smith R, Walker F, Ferrario C. Number of antihypertensive agents required to achieve goal blood pressure with an olmesartan medoxomil vs atenolol based regimen. 22nd Annual American Society of Hypertension Scientific Meeting, May 2007. [Poster]

AZOR™ (amlodipine and olmesartan medoxomil)  
Formulary Dossier

**OLMESARTAN MEDOXOMIL CLINICAL BIBLIOGRAPHY**  
(continued)

Smith R, Yokoyama H, Ferrario C. Noninvasive hemodynamic data reflect treatment effects on vascular remodeling. 22nd Annual American Society of Hypertension Scientific Meeting, May 2007. [Abstract]

Smith R, Yokoyama H, Ferrario C. Noninvasive hemodynamic data reflect treatment effects on vascular remodeling. 22nd Annual American Society of Hypertension Scientific Meeting, May 2007. [Abstract]

Smith R, Yokoyama H. Patient satisfaction allows better blood pressure control. 22nd Annual American Society of Hypertension Scientific Meeting, May 2007. [Abstract]

Smith R, Yokoyama H. Patient satisfaction allows better blood pressure control. 22nd Annual American Society of Hypertension Scientific Meeting, May 2007. [Poster]

Smith R, Yokoyama D, Averill E, et al. The relative effect of renin angiotensin system and the adrenergic system on vascular remodeling. 15th European Meeting on Hypertension, June 2005.

Smith R, Yokoyama H, Averill E, et al. The protective effects of angiotensin ii blockade with olmesartan medoxomil on resistance vessel remodeling (the VIOS study): Rationale and baseline characteristics. *American Journal of Cardiovascular Drugs*. 2006;6(5):335-342.

Smith RD, Averill D, Brosnihan KB, et al. Effects of antihypertensive therapy on insulin sensitivity and leptin values. 21st Scientific Meeting of International Society of Hypertension (ISH), October 2006. [Abstract]

Smith RD, Brosnihan KB, Owen J. Cross talk of the angiotensin at1 receptor and the thromboxane receptor. 21st Scientific Meeting of International Society of Hypertension (ISH), October 2006. [Abstract]

Smith RD, Yokoyama H, Brosnihan KB, et al. Inflammatory markers (im) in uncomplicated hypertension. 21st Scientific Meeting of International Society of Hypertension (ISH), October 2006. [Abstract]

Smith RD, Yokoyama H, Brosnihan KB, et al. Inflammatory markers in uncomplicated hypertension. 22nd Annual American Society of Hypertension Scientific Meeting, May 2007. [Poster]

Smith RD, Yokoyama H, Ferrario CM. Noninvasive hemodynamic data reflect treatment effects on vascular remodeling. 16th Scientific Meeting, European Society of Hypertension (ESH), June 2006. [Poster]

Smith RD, Yokoyama H, Ferrario CM. Patient satisfaction allows better blood pressure control. 21st Scientific Meeting of International Society of Hypertension (ISH), October 2006. [Abstract]

Smolensky MH, Hermida RC, Portaluppi F. Comparison of the efficacy of morning versus evening administration of olmesartan in uncomplicated essential hypertension. *Chronobiol Int*. 2007;24(1):171-81.

Song JC, White CM. Olmesartan medoxomil (CS-866) - An angiotensin II receptor blocker for treatment of hypertension. *Formulary*. 2001; 36:487-499.

Stefenelli T, Ewald S, Boehm M. OLMEBEST - study: Reduction of blood pressure in the treatment of patients with mild-to-moderate essential hypertension. - Interim analysis of the Austrian part. 65th Annual American Society of Hypertension Conference, May 2006. [Poster]

Steinberg JS., RPh, BS Pharm. Angiotensin II receptor antagonists: Mechanisms and therapeutic utility. *Hospital Pharmacy*. 1998; 33: 53-65.

Stone AA. An efficacy evaluation of olmesartan medoxomil/hydrochlorothiazide (om/hct) and amlodipine besylate/benazepril hydrochloride (am/bn). *AJH* (antihypertensive drugs and pharmacology). 2005;18(5): part 2. [Abstract]

AZOR™ (amlodipine and olmesartan medoxomil)  
Formulary Dossier

**OLMESARTAN MEDOXOMIL CLINICAL BIBLIOGRAPHY**  
(continued)

Stump CS, Sowers JR. Prevention of type 2 diabetes: Role of the renin-angiotensin-aldosterone system and antihypertensive therapy. *Adv Stud Med*. 2006;6(5):231-239.

Stumpe K, Agabiti-Rosei E, Scholze J, et al. Effects of olmesartan and atenolol on carotid atherosclerosis. The multicenter olmesartan atherosclerosis regression evaluation (MORE) trial. 17th annual european meeting on hypertension, June 2007. [Poster]

Stumpe KO, Agabiti-Rosei E, Scholze J, et al. Carotid atherosclerosis regression by angiotensin receptor blockade: Results of the multicentre olmesartan atherosclerosis regression evaluation (MORE) study. 22nd annual american society of hypertension scientific meeting, May 2007. [Poster]

Stumpe KO, M Ludwig. Antihypertensive efficacy of olmesartan compared with other antihypertensive drugs. *J Hum HTN*. 2002; 16 (Supl 2): S24-S28.

Stumpe KO. Olmesartan Medoxomil: A Viewpoint by KO Stumpe. *Drugs*. 2002; 62(9):1354-6.

Stumpe KO. Olmesartan compared with other angiotensin ii receptor antagonists: Head-to-head trials. *Clinical Therapeutics*. 2004;26(suppl a):a33-a37.

Sugie T, Kagaya Y, Takeda M, Yahagi H, et al. Should increasing the dose or adding an AT1 receptor blocker follow a relatively low dose of ACE inhibitor initiated in acute myocardial infarction? *Cardiovascular Research*. 2003; 58(3):611-20.

Takai S, Miyazaki M. Effect of olmesartan medoxomil on atherosclerosis: Clinical implications of the emerging evidence. *Am J Cardiovasc Drugs*. 2006;6(6):363-6.

Terra SG. Cardiology patient page. Angiotensin receptor blockers. *Circulation*. 2003; 107(24):E215-6.

Trenkwalder P, Samer H, Hempel RD, et al. Fixed dose combination of valsartan 160/hctz 25 in hypertensive patients not controlled by monotherapy with olmesartan 40 or olmesartan 20 plus hctz. 21st Scientific Meeting of International Society of Hypertension (ISH), October 2006. [Abstract]

Turnbull F. Effects of different blood-pressure-lowering regimens on major cardiovascular events: results of prospectively-Designed overviews of randomised trials. *Lancet*. 2003;362:1527-35.

Uen S, Ewald S, Kaufmann S, et.al. Telemonitoring of blood pressure self-measurement (bpsm) in the OLMETEL study. *Journal of Hypertension*. 2004;22(s2):s26 [Abstract p1.20]

Unger T, Mcinnes GT, Neutel JM et al. The role of olmesartan medoxomil in the management of hypertension. *Drugs*. 2004;64(24):2731-2739.

Ura N, Yamagushi K, Togashi N, et al. The mechanism of insulin resistance improvement by angiotensin 2 receptor blocker. 15th European Meeting on Hypertension, June 2005.

Urae, A, Amamoto T, Urae R, Hisaoka M, et al. Tolerability, Pharmacokinetics and Pharmacodynamics of a New Angiotensin II Receptor Antagonist CS-866 in Healthy Male Volunteers. Aug. 5, 1996. The 6th International Meeting of Clinical Pharmacology.

Van Der Hoorn JWA, Jukema JW, Maas MC, et al. Olmesartan and pravastatin additively reduce development of atherosclerosis. Dept of Cardiology, Leiden University Medical Centre and Biomedical Research, Tno Gaubius Lab, the Netherlands. [Abstract]

Van Liefde I, Le MT, Pugsley M, et al. A characterization of the molecular interactions of olmesartan and telmisartan with human angiotensin at1 receptors. 65th annual american society of hypertension conference, May 2006. [Abstract]

AZOR™ (amlodipine and olmesartan medoxomil)  
Formulary Dossier

**OLMESARTAN MEDOXOMIL CLINICAL BIBLIOGRAPHY**  
(continued)

- Van W Miegheem. A Multi-Centre, Double-Blind, Efficacy, Tolerability and Safety Study of the Oral Angiotensin II-Antagonist Olmesartan Medoxomil versus Atenolol in Patients with Mild to Moderate Essential Hypertension. *Journal of HTN*. 2001; 19 (suppl 2): P2.174.
- Venkata C., Ram S. Angiotensin receptor blockers and diuretics as combination therapy: clinical implications. *American Journal of Hypertension*. 2004; 17(3):277-280.
- Venketa C, Ram S. Antihypertensive efficacy of angiotensinreceptor blockers in combination with hydrochlorothiazide: A review of the factorial-design studies. *J Clin Hypertens*. 2004;6:569–577.
- Volpe M, Trimarco B. A novel molecule of the angiotensin ii receptor blocker class: Focus on olmesartan medoxomil. *High Blood Pressure Cardiovascu Prev*. 2005;12(2):79-89.
- Voors AA. Angiotensin II AT1 receptor antagonisten anno. *Heart Bulletin*. 1997; 28: 157-159.
- Ward K, Humme A. Olmesartan, benicar, for hypertension. *American Family Physician*. 2005;72(4).
- Warner GT, B Jarvis. Olmesartan Medoxomil. *Drugs*. 2002: 62(9):1345-53.
- Weber M, Basile J, Izzo J, et al. Blood pressure goal attainment: meeting the challenge of the JNC 7's blood pressure goals and the role of renin-angiotensin-aldosterone system blockade. *The Journal of Clinical Hypertension*. 2004;6(12):699-705.
- Weber M. Concluding Comment. *American Journal of Cardiology*. 2001;87(8A):44C.
- Wehling M. Can the pharmacokinetic characteristics of olmesartan medoxomil contribute to the improvement of blood pressure control? *Clinical Therapeutics*. 2004;26(suppl a):a21-a27.
- Williams PA. A Multi-Centre, Double-Blind, Efficacy, Tolerability and Safety Study of the Oral Angiotensin II-Antagonist Olmesartan Medoxomil versus Captopril in patients with Mild to Moderate Essential Hypertension. Abstract: 207.
- Woittiez AJ, Ewald S, Boehm M. OLMEBEST - study: Reduction of blood pressure in the treatment of patients with mild-to-moderate essential hypertension. - Interims analysis of the Dutch part. 65th Annual American Society of Hypertension Conference, May 2006. [Poster]
- Yokoyama H, Smith RD, Averill DB, et al. Vascular improvement with olmesartan medoxomil. *AJH (Arterial structure and compliance)*. 2005;18(5):part 2 [Abstract]
- Yokoyoma H, Averill D, Brosnihan KB, et al. Role of blood pressure reduction in prevention of cardiac and vascular hypertrophy. *AJH*. 2005;18:922-929.
- Yoshida K, Kohzuki M. Clinical and experimental aspects of olmesartan medoxomil, A new angiotensin ii receptor antagonist. *Cardiovascular Drug Reviews*. 2004;22(4):285-308.
- Yoshihara K, Gao Y, Shiga H, et al. Population pharmacokinetics of olmesartan following oral administration of its prodrug, olmesartan medoxomil - In healthy volunteers and hypertensives patients. *Clin Pharmacokinet*. 2005;44(12):1329-1342.
- Zannad F, Fay R. Blood pressure-lowering efficacy of olmesartan relative to other angiotensin ii receptor antagonists: An overview of randomized controlled studies. *Fundam Clin Pharmacol*. 2007;21(2):181-90.

AZOR™ (amlodipine and olmesartan medoxomil)  
Formulary Dossier

**d. OLMESARTAN MEDOXOMIL PRE-CLINICAL BIBLIOGRAPHY**

69th Scientific Sessions of the American Heart Association; New Orleans, Circulation 94, No. 8, Sup., I657, 1996 (Abstract: 3839).

Agata J, Ura N, Shinshi Y, et al. Olmesartan is an Angiotensin II Receptor Blocker with an Inhibitory Effect of Angiotensin-Converting Enzyme. 21st Scientific Meeting of International Society of Hypertension (ISH), October 2006. [Abstract]

Akishita M, Nagai K, Xi H, et al. Renin-Angiotensin System Modulates Oxidative Stress-Induced Endothelial Cell Apoptosis in Rats. *Hypertension*. 2005;45:1-6.

Anderson BE, Zeiger E, Shelby MD, et al. Chromosome aberration and sister chromatid exchange test results with 42 chemicals. *Environ. Molec. Mutagen*. 16(Suppl.18):55-137.

Arishiro K, Hoshiga M, Negoro N, et al. Angiotensin receptor-1 blocker inhibits atherosclerotic changes and endothelial disruption of the aortic valve in hypercholesterolemic rabbits. *J Am Coll Cardiol*. 2007;49(13):1482-9.

Asano Y, et al. Reproductive and developmental toxicity studies of imidapril hydrochloride in rats and rabbits. *The Clinical Report*. 1992; 26:4669-4676.

Bagdon WJ, Bokelman DL, Stone CA. Toxicity study of MK-421 (Enalapril maleate)(II); Subacute, chronic toxicity studies, saline supplementation, sodium-depletion studies in rats. *Yakuri to Chiryō* 1985;13:425-466.

Barr M, Jr. Teratogen update: angiotensin-converting enzyme inhibitors. *Teratology* 1994; 50:399-409.

Bo Zhang, Ping Fan, Hiroaki Tanaka and Keijiro Saku. Angiotensin II type-1 receptor antagonist, CS-866, reduces blood pressure without affecting glucose/insulin metabolism in cholesterol-fed rabbits. *British Journal of Biomedical Science*. 2001; 58: 217-225.

Brunner H. Olmesartan medoxomil: a new AT1 receptor antagonist for the treatment of hypertension. *Journal of Hypertension*. 2001; 19:S1-S14.

Bucher JR. Toxicology and carcinogenesis studies of hydrochlorothiazide (CAS No. 58-93-5) in F344/N rats and B6C3F1 mice (feed studies). U.S. Department of Health and Human Services, National Toxicology Program. Report No. NTP TR 357. NIH Publication No. 89-2812.

Buchwalder-Csajka C., Buclin T., Brunner HR., Biollaz J. Evaluation of the angiotensin challenge methodology for assessing the pharmacodynamic profile of antihypertensive drugs acting on the renin-angiotensin system. *British Journal of Clinical Pharmacology*. 1999; 48(4): 594-604.

Buclin T. Buchwalder-Csajka C. Brunner HR. Biollaz J. Evaluation of noninvasive blood pressure recording by photoplethysmography in clinical studies using angiotensin challenges. *British Journal of Clinical Pharmacology*. 1999; 48(4): 586-593.

Carey RM. Angiotensin type-1 receptor blockade increases ACE 2 expression in the heart. *Hypertension*. 2004;43(5):943-4.

Celebier M, Altinoz S. Determination of Olmesartan Medoxomil in Tablets by UV-Vis Spectrophotometry. *Pharmazie*. 2007;62:419-422.

Chappell MC, Gallagher PE, Averill DB, et al. Estrogen or the AT1 Antagonist Olmesartan Reverses the Development of Profound Hypertension in the Cogenic mRen2.Lewis Rat. *Hypertension*. 2003;42[part 2]:781-786.

AZOR™ (amlodipine and olmesartan medoxomil)  
Formulary Dossier

**OLMESARTAN MEDOXOMIL PRE-CLINICAL BIBLIOGRAPHY**  
(continued)

- Chappell MC, Gallagher PE, Averill DB, Ferrario CM, Brosnihan KB. Estrogen or the AT1 Antagonist Olmesartan Reverses the Development of Profound Hypertension in the Cogenic mRen2.Lewis Rat. *Hypertension*. 2003;42[part 2]:781-786.
- Chen R, Iwai M, Tomono Y, et al. Effect of Combination of Pravastatin and Olmesartan on Glucose Intolerance in Type 2 Diabetic kk-Ay MICE. 21st Scientific Meeting of International Society of Hypertension (ISH), October 2006. [Abstract]
- Chiba S, Yang XP. The preferential inhibitory effect of olmesartan, a new angiotensin II type 1 antagonist, on sympathetic nerve terminals in isolated canine splenic artery. *Journal of Pharmacological Sciences*. 2003 August;92(4):381-386.
- Coyle J, Gardener S. The Renal Protective Effects of Angiotensin II Receptor Blockers in Type II Diabetes Mellitus. *The Annals of Pharmacotherapy*. 2004;38:1731-8.
- de Faire U. Anti-atherosclerotic effects by AT1 receptor blockade. *Journal of Hypertension*. 2003;21:259-260.
- De Faire U. Anti-atherosclerotic effects by AT1 receptor blockade. *Journal of Hypertension*. 2003;21:259-260.
- Eguchi S, Dempsey PJ, Frank GD, et al. Activation of MAP kinases by angiotensin II in vascular smooth muscle cells: metalloprotease-dependent EGF receptor activation is required for activation of ERK and p38 MAP kinase, but not for JNK. *J Biol Chem*. 2001; 276:7957-7962.
- Erbe DV, Gartrell K, Zhang YL, et al. Molecular Activation of PPAR- $\gamma$  by Angiotensin II type 1-receptor antagonists. *Vascular Pharmacology*. 2006;(45)154-162.
- Fan YY, Baba R, Nagai Y, et al. Augmentation of Intrarenal Angiotensin II Levels in Uninephrectomized Aldosterone/Salt-Treated Hypertensive Rats; Renoprotective Effects of an Ultrahigh Dose of Olmesartan. *Hypertens Res*. 2006;29(3):169-78.
- Faure S, Oudart N, Javellaud J, et al. Synergistic protective effects of erythropoietin and olmesartan on ischemic stroke survival and post-stroke memory dysfunctions in the gerbil. *J Hypertens*. 2006;24(11):2255-61.
- Ferrario CM. Use of angiotensin II receptor blockers in animal models of atherosclerosis, *Am Journal of HTN*. 2002; 15(1 Pt 2):9S-13S.
- Fleming I, Kohlstedt K, Busse R. The tissue renin-angiotensin system and intracellular signaling. *Curr Opin Nephrol Hypertens*. 2006;15:8-13.
- Fujita M, Okuda H, Tsukamoto O, et al. Blockade of Angiotensin II Receptors Reduces the Expression of Receptors for Advanced Glycation End Products in Human Endothelial Cells. *Arteriosclerosis, Thrombosis & Vascular Biology*. 2006;26:e138.
- Fukuda N, Satoh C, Hu WY, Nakayama M, Kishioka H, et al. Endogenous angiotensin II suppresses insulin signaling in vascular smooth muscle cells from spontaneously hypertensive rats. *J Hypertens*. 2001;19(9):1651-8.
- Furuhashi M, Ura N. Blockade of the Renin Angiotensin System Decreases Adipocyte Size with Improvement in Insulin Sensitivity. *Journal of Hypertension*. 2004;22(10):1977-1982.
- Galloway SM, Armstrong MJ, Reuben C, et al. Chromosome aberrations and sister chromatid exchanges in Chinese hamster ovary cells: evaluations of 108 chemicals. *Environ. Molec. Mutagen*. 1987; 10(Suppl. 10):1-175.



AZOR™ (amlodipine and olmesartan medoxomil)  
Formulary Dossier

**OLMESARTAN MEDOXOMIL PRE-CLINICAL BIBLIOGRAPHY**  
(continued)

Galloway SM. Cytotoxicity and chromosome aberrations in vitro: Experience in industry and the case for an upper limit on toxicity in the aberration assay. *Environ. Molec. Mutagen.* 2000, 35:191-201.

George JD, Price CJ, Wolkowshki-Tyl RW, et al. The evaluation of the developmental toxicity of hydrochlorothiazide in mice and rats. *Fund. Appl. Toxicol.* 1995, 26:174-180.

Ghanim H, Viswanathan P, Sia CL, et al. Olmesartan, an Angiotensin II Receptor Blocker, Suppresses Oxidative Stress and Inflammation in Patients with Coronary Artery Disease or Risk Factors for Atherosclerosis. Endocrine Society 89th Annual Meeting. [Abstract]

Gonzalez-Villalobos RA, Klassen RB, Allen PL, et al. Megalin binds and internalizes Angiotensin-(1-7). *American Journal of Physiology. Renal physiology.* 2006;290(5):F1270-5.

Graul A., Lesson P., Castaner J. CS-866 and etanercept. *Drugs of the Future.* Vol 22(11) (pp 1205-1209), 1997.

Harada N, Ohnaka M, Sakamoto S et al. Cilnidipine improves insulin sensitivity in the Otsuka Long-Evans Tokushima Fatty rat, a model of spontaneous NIDDM. *Cardiovascular Drugs and Therapy.* 1999, 13: 519-523.

Hideharu Tomita, Kensuke Egashira, Yuichi Ohara et al. Early Induction of Transforming Growth Factor-beta via Angiotensin II Type 1 Receptors Contributes to Cardiac Fibrosis Induced by Long-term Blockade of Nitric Oxide Synthesis in Rats. *Hypertension* 1998;32:273-279.

Hidetomo Nakamoto, Hiroe Imai, Rie Fukushima, et al. Angiotensin Receptor Blocker (ARB) Protects the Development of Encapsulating Peritoneal Sclerosis on Peritoneal Dialysis in Spontaneously Hypertensive Rats. 5th international Meeting on Animal Models in Peritoneal Dialysis(ANIMOD V) Brussels, 3-4 May 2002. *Perit Dial Int* 2002; 22: 157.

Higashiura K, Ura N, Takada T, Agata J, Yoshida H et al The effect of angiotensin converting enzyme inhibitor and angiotensin receptor antagonist on insulin sensitivity in fructose-fed rats. International Society of Hypertension 17th Scientific Meeting; Amsterdam, *Journal of Hypertension* 16, Sup. 2, S154, 1998.

Higashiura K, Ura N, Takada T, Li Y, Torii T, et al. The effects of an angiotensin-converting enzyme inhibitor and an angiotensin II receptor antagonist on insulin resistance in fructose-fed rats. *Am J Hypertens.* 2000 Mar;13(3):290-7.

Hillard CA, Armstrong MJ, Bradt RB, et al. Chromosome aberrations in vitro related to cytotoxicity of nonmutagenic chemicals and metabolic poisons. *Environ. Molec. Mutagen.* 1998, 31:316-326.

Hiramatsu N, Hiromura K, Shigehara T, et al. Angiotensin II Type 1 Receptor Blockade Inhibits the Development and Progression of HIV-Associated Nephropathy in a Mouse Model. *J Am Soc Nephrol.* Jan 2007.

Hirano T, Ran J, Adachi M. Opposing actions of angiotensin II type 1 and 2 receptors on plasma cholesterol levels in rats. *J Hypertens.* 2006;24(1):103-8.

Hirano T, Ran J, Okada K, et al. Angiotensin II receptor blocker, olmesartan medoxomil, ameliorates triglyceride overproduction in insulin-resistant rats independent of hypotensive action. American Diabetes Association Scientific Meeting [938-P], June 2004.

Hirano T. Angiotensin II Infusion Decreases Plasma Adiponectin Level via Type 1 Receptor in Rats: Implication For Insulin Resistance Associated with Hypertension. [ABSTRACT]

AZOR™ (amlodipine and olmesartan medoxomil)  
Formulary Dossier

**OLMESARTAN MEDOXOMIL PRE-CLINICAL BIBLIOGRAPHY**  
(continued)

Hiroaki Yanagisawa, Yoshiya Amemiya, Takuro Kanazaki, et al. Nonpeptide Angiotensin II Receptor Antagonist: Synthesis, Biological Activities, and Structure-Activity Relationships of Imidazole-5-carboxylic Acids Bearing Alkyl, Alkenyl, and Hydroxyalkyl substituents at the 4-Position and Their Related Compounds. *Journal of Medicinal Chemistry*, 1996, Vol. 39, No.1: 323-338.

Hiroaki Yanagisawa. Antihypertensive Agents Suppressing the Renin-Angiotensin system. Aug. 24, 1996. The 14th Summer Meeting of the Synthetic Organic Chemistry.

Hiromi Rakugi, Atsunori Okamura, et al. Recognition of Tissue- and Subtype-Specific Modulation of Angiotensin II Receptors Using Antibodies antagonist AT1 and AT2 Receptors. *Hypertension Research*, 1997, Vol. 20, No. 1, 51-55.

Hirose A, Ono M, Saibara T, et al. Angiotensin II type 1 receptor blocker inhibits fibrosis in rat nonalcoholic steatohepatitis. *Hepatology*. 2007;45(6):1375-81.

Hong HJ, Chan P, Liu JC, et al. Angiotensin II induces endothelin-1 gene expression via extracellular signal-regulated kinase pathway in rat aortic smooth muscle cells *Cardiovascular Research*. 2004; 61(1):159-168.

Hong HJ, Chan P, Liu JC, Juan SH, Huang MT, Lin JG, Cheng TH. Angiotensin II induces endothelin-1 gene expression via extracellular signal-regulated kinase pathway in rat aortic smooth muscle cells. *Cardiovascular Research*. 2004 January; (61)1:159-168.

Hosomi N, Nishiyama A, Ban CR, et al. Angiotensin type 1 receptor blockage improves ischemic injury following transient focal cerebral ischemia. *Neuroscience*. 2005;134(1):225-31.

Hussain W, Patel P, Dhillon P. Olmesartan Reverses Changes In Conduction Velocity, Connexin43 Expression, And Distribution Associated With Localized In Situ Ventricular Stretch Univ, Sapporo, Japan Salutory, November 2004. [Abstract]

Ideura H, Hiromura K, Hiramatsu N, et al. Angiotensin II provokes podocyte injury in murine model of HIV-associated nephropathy. *Am J Physiol Renal Physiol*. July 2007.

Ido A, Hasebe N, Takeuchi T, Kikuchi K. Effects of temocapril and olmesartan on myocardial sympathetic nervous activity and fatty acid metabolism in rats with chronic beta-adrenergic stimulation. *Journal of Cardiovascular Pharmacology*. 2003 Jan;41 Suppl 1:S133-7.

Igase M, Strawn WB, Gallagher PE, et al. Angiotensin II AT1 receptors regulate ACE2 and angiotensin-(1-7) expression in the aorta of spontaneously hypertensive rats. *American Journal of Physiology (Heart & Circulatory Physiology)*. 2005;289(3):H1013-9.

Imai H, Nakamoto H, Fukushima R, et al. Role of adhesion molecules in the progression of peritoneal sclerosis. *Advances in Peritoneal Dialysis*. 2003;19:180-185.

Imai K, Yoshimura S., Hashimoto K. One month studies on the subacute toxicity of captopril in the rat. *Toxicol. Sci.* 1981; 6(Suppl.II): 189-214.

Inoue T, Nakamura H, Arakawa N, et al. Olmesartan medoxomil, an angiotensin II receptor antagonist, may have beneficial effects on diabetic retinopathy American Diabetes Association Scientific Meeting [887-P], June 2004.

Inoue T, Nakamura H, Arakawa N, et al. Olmesartan Medoxomil, an Angiotensin Receptor Antagonist, May Have Beneficial Effects on Diabetic Retinopathy. Univ, Sapporo, Japan Salutory, November 2004. [Abstract]

AZOR™ (amlodipine and olmesartan medoxomil)  
Formulary Dossier

**OLMESARTAN MEDOXOMIL PRE-CLINICAL BIBLIOGRAPHY**  
(continued)

- Inukai T, Yoshida N, Wakabayashi S, et al. Angiotensin-converting enzyme inhibitors and angiotensin II receptor blockers effectively and directly potentiate superoxide scavenging by polymorphonuclear leukocytes from patients with type 2 diabetes mellitus. *Am J Med Sci.* 2005;329(5):222-7.
- Isfort RJ, Kerckaert GA, LeBoeuf RA. Comparison of the standard and reduced pH Syrian hamster embryo (SHE) cell in vitro transformation assays in predicting the carcinogenic potential of chemicals. *Mutat. Res.* 1996, 356:11-63.
- Ishibashi M, Hiasa K, Zhao Q, et al. Critical role of monocyte chemoattractant protein-1 receptor CCR2 on monocytes in hypertension-induced vascular inflammation and remodeling. *Circ Res.* 2004;94(9):1203-10.
- Ishida Y, Tomori K, Nakamoto H, et al. Effects of antihypertensive drugs on peritoneal vessels in hypertensive dogs with mild renal insufficiency. *Advances in Peritoneal Dialysis.* 2003;19:10-14.
- Ishiyama Y., Gallagher DB, et al. Upregulation of Angiotensin Converting Enzyme 2 After Myocardial Infarction by Blockade of Angiotensin II Receptors. *Hypertension.* 2004;43:970-976.
- Ito N, Ohishi M, Yamamoto K, et al. Renin-Angiotensin Inhibition Reverses Advanced Cardiac Remodeling in Aging Spontaneously Hypertensive Rats. *AJH.* 2007;20:792-799.
- Ito N, Ohishi M, Yamamoto K, et al. Renin-angiotensin system blockade effects on improvement of advance renal damages in high-aged spontaneously hypertensive rats via suppression of oxidized stress. 15th European Meeting on Hypertension, June 2005.
- Itoh S, Ding B, Shishido T, et al. Role of p90 ribosomal S6 kinase-mediated prorenin-converting enzyme in ischemic and diabetic myocardium. *Circulation.* 2006 Apr 11;113(14):1787-98.
- Itoh T, Kajikuri J, Tada T, Suzuki Y, Mabuchi Y. Angiotensin II-induced modulation of endothelium-dependent relaxation in rabbit mesenteric resistance arteries. *Journal of Physiology.* 2003 May 1;548(Pt 3):893-906.
- Iwai M, Chen R, Ide A, et al. The calcium-channel blocker, azelnidipine, enhances the inhibitory action of AT1 receptor blockade on ischemic brain damage. *Journal of Hypertension.* 2006;24(10):2023-31.
- Iwai M, Li HS, Chen R, et al. Calcium Channel Blocker, Azelnidipine, Reduces Glucose Intolerance in Diabetic Mice via Different Mechanism to Angiotensin Receptor Blocker, Olmesartan. *Journal of Pharmacology and Experimental Therapeutics.* 2006 Sep 21.
- Iwasaki Y, Ichikawa Y, Igarasi O. Effect of Olmesartan, A novel ATR antagonist, on spinal motor neurons in vitro and in vivo. *Amyotroph Lateral Scler Other Motor Neuron Disord* 2001; 2(Suppl.2): 154.
- Iwasaki Y, Ichikawa I, Igarashi O, et al. Trophic effect of olmesartan, a novel AT 1 receptor antagonist, on spinal motor neurons in vitro and in vivo. *Neurological Research.* 2002; 24(5): 468-72.
- Izuhara Y, Nangaku M, Inagi R, et al. Renoprotective Properties of Angiotensin Receptor Blockers beyond Blood Pressure Lowering. *Journal of the American Society of Nephrology.* 2005;16(12):3631-41.
- Jackson EK. Commentary on Liu et al: Effect of Estrogen and AT1 Receptor Blocker on Neointima Formation. *Hypertension.* 2002; 40(4): 448-50.

AZOR™ (amlodipine and olmesartan medoxomil)  
Formulary Dossier

**OLMESARTAN MEDOXOMIL PRE-CLINICAL BIBLIOGRAPHY**  
(continued)

- Jagadeesh G. Angiotensin II receptor-antagonists, molecular biology, and signal transduction. *Indian Journal of Experimental Biology*. 1998; 36(12): 1171-1194.
- Jia N, Okamoto H, Shimizu T, et al. A newly developed angiotensin II type 1 receptor antagonist, CS866, promotes regression of cardiac hypertrophy by reducing integrin beta1 expression. *Hypertens Res*. 2003;26(9):737-42.
- Jinno T, Iwai M, Li Z, et al. Calcium Channel Blocker Azelnidipine Enhances Vascular Protective Effects of AT1 Receptor Blocker Olmesartan. *Hypertension*. 2004;43:1-7.
- Jinno T, Iwai M, Li Z, Li JM, Liu HW, Cui TX, Rakugi H, Ogihara T, Horiuchi M. Calcium Channel Blocker Azelnidipine Enhances Vascular Protective Effects of AT1 Receptor Blocker Olmesartan. *Hypertension*. 2004 February;43:1-7.
- Kamada Y, Tamura S, Kiso S, Fukui K, Doi Y, et al. Angiotensin II stimulates the nuclear translocation of Smad2 and induces PAI-1 mRNA in rat hepatic stellate cells. *Hepatology Research*. 2003 Mar;25(3):296-305.
- Kamiyama E, Yoshigae Y, Kasuya A, et al. Inhibitory Effects of Angiotension Receptor Blockers on CYP2C9 Activity in Human Liver Microsomes. *Drug Metab Pharmacokinet*. 2007;22(4):267-275.
- Kanamori H, Takemura G, Li Y, et al. Inhibition of Fas-associated apoptosis in granulation tissue cells accompanies attenuation of postinfarction left ventricular remodeling by olmesartan. *Am J Physiol Heart Circ Physiol*. Jan 2007.
- Kanazawa K, Inoue I, Ikeda M, et al. Angiotensin Type 1 Receptor Blocker, Olmesartan, Induces Transcription of Peroxisome Proliferator-Activated Receptor  $\gamma$  mRNA through the Promoter of PPAR  $\gamma$ . 21st Scientific Meeting of International Society of Hypertension (ISH), October 2006. [Abstract]
- Kang JJ, Toma I, Sipos A, et al. Imaging the renin-angiotensin system: An important target of anti-hypertensive therapy. *Advanced Drug Delivery Reviews*. 2006;58(7):824-33.
- Kato M, Sada T, Mizuno M, et al. Effect of Combined Treatment With an Angiotensin II Receptor Antagonist and an HMG-CoA Reductase Inhibitor on Atherosclerosis in Genetically Hyperlipidemic Rabbits. *Journal of Cardiovascular Pharmacology*. 2005;46(4):556-562.
- Katoh M, Egashira K, Kataoka C, Usui M, Koyanagi M, et al. Regression by ACE inhibition of arteriosclerotic changes induced by chronic blockade of NO synthesis in rats. *American Journal of Physiology-Heart and Circulatory Physiology*. 2001;280(5):H2306-2312.
- Kawasaki H, Inaizumi K, Nakamura A, Hobara N, et al. Chronic angiotensin II inhibition increases levels of calcitonin gene-related peptide mRNA of the dorsal root ganglia in spontaneously hypertensive rats. *Hypertension Research*. 2003 Mar;26(3):257-63.
- Kei Kamide, Hiromi Rakugi, Jitsuo Higaki, et al. The Renin-Angiotensin and Adrenergic Nervous System in Cardiac Hypertrophy in Fructose-Fed Rats. *American Journal of Hypertension*. 2002; 15: 66-71.
- Kikuchi T, Ito N, Masahiko Suzuki M, et al. Self-association properties of 4-[1-hydroxy-1-methylethyl]-2-propyl-1-[4-[2-[tetrazole-5-yl]phenyl]phenyl] methylimidazole-5-carboxylic acid monohydrate (CS-088), an antiglaucoma ophthalmic agent International. *Journal of Pharmaceutics*. 2005;299:100–106.
- Kikuchi T, Suzuki M, Kusai A, et al. Mechanism of permeability-enhancing effect of EDTA and boric acid on the corneal penetration of 4-[1-hydroxy-1-methylethyl]-2-propyl-1-[4-[2-[tetrazole-5-yl]phenyl]phenyl] methylimidazole-5- carboxylic acid monohydrate (CS-088). *Int J Pharm*. 2005;299(1-2):107-14.

AZOR™ (amlodipine and olmesartan medoxomil)  
Formulary Dossier

**OLMESARTAN MEDOXOMIL PRE-CLINICAL BIBLIOGRAPHY**  
(continued)

Kim S, Izumi Y, Izumiya Y, Zhan Y, Taniguchi M, Iwao H. Beneficial effects of combined blockade of ACE and AT1 receptor on intimal hyperplasia in balloon-injured rat artery. *Arteriosclerosis, Thrombosis and Vascular Biology*. 2002; 22(8): 1299-1304.

Kim-Mitsuyama S, Izumi Y, Izumiya Y, et al. Additive beneficial effects of the combination of a calcium channel blocker and an angiotensin blocker on a hypertensive rat-heart failure model. *Hypertens Res*. 2004;27(10):771-9.

Kitamoto S, Egashira K, Kataoka C, Usui M, et al. Chronic inhibition of nitric oxide synthesis in rats increases aortic superoxide anion production via the action of angiotensin II. *J Hypertens*. 2000;18(12):1795-800.

Kobayashi N, Nakano N, Fukushima H, et al. Cardioprotective Mechanism of Combination Therapy with Angiotensin II Receptor Blocker and Statin in Failing Heart of Dahl Salt-Sensitive Rats. 17th Annual European Meeting on Hypertension, June 2007. [Poster]

Kobori H, Prieto-Carrasquero MC, Ozawa Y, et al. AT1 Receptor Mediated Augmentation of Intrarenal Angiotensinogen in Angiotensin II-Dependent Hypertension. *Hypertension*. 2004;43:1126-1132.

Kobori H. Ozawa Y. Suzuki Y. et al. Enhanced intrarenal angiotensinogen contributes to early renal injury in spontaneously hypertensive rats. *Journal of the American Society of Nephrology*. 2005;16(7):2073-80.

Koga K, Yamagishi S, Takeuchi M, Inagaki Y, et al. CS-886, A New Angiotensin II Type 1 Receptor Antagonist, Ameliorates Glomerular Anionic Site Loss and Prevents Progression of Diabetic Nephropathy in Otsuka Long-Evans Tokushima Fatty Rats. *Mol Med*. 2002 Oct;8(10):591-9.

Koike H. New Pharmacologic aspects of CS-866, the newest angiotensin II receptor antagonist. *American Journal of Cardiology*. 2001;87(8A):33-6.

Kurikawa N, Suga M, Kuroda S, et al. An angiotensin II type 1 receptor antagonist, olmesartan medoxomil, improves experimental liver fibrosis by suppression of proliferation and collagen synthesis in activated hepatic stellate cells. *British Journal of Pharmacology*. 2003;139:1085-1094.

Kurikawa N, Suga M, Kuroda S, Yamada K, Ishikawa H. An angiotensin II type 1 receptor antagonist, olmesartan medoxomil, improves experimental liver fibrosis by suppression of proliferation and collagen synthesis in activated hepatic stellate cells. *British Journal of Pharmacology*. 2003(139):1085-1094.

Le MT, Pugsley MK, Vauquelin G, et al. Molecular characterisation of the interactions between olmesartan and telmisartan and the human angiotensin II AT1 receptor. *British Journal of Pharmacology*. 2007;151:952-962.

Lee TM, Lin MS, Chou TF, et al. Effects of Combined Blockade of AT1 Receptor and HMG-CoA Reductase on Left Ventricular Remodeling in Infarcted Rats. 2004 [POSTER]

Lee TM, Lin MS, Chou TF, et al. Additive Effects of Combined Blockade of AT1 Receptor and HMG-CoA Reductase on Left Ventricular Remodeling in Infarcted Rats. *Am J Physiol Heart Circ Physiol*. March 2006.

Lee TM, Lin MS, Chou TF, et al. Additive effects of combined blockade of AT1 receptor and HMG-CoA reductase on left ventricular remodeling in infarcted rats. *American Journal of Physiology (Heart & Circulatory Physiology)*. 2006;291(3):H1281-9.

Li Y, Kishimoto I, Saito Y, Harada M, et al. Guanylyl Cyclase-A Inhibits Angiotensin II Type 1A Receptor-Mediated Cardiac Remodeling, an Endogenous Protective Mechanism in the Heart. *Circulation*. 2002; 106(13): 1722-8.

AZOR™ (amlodipine and olmesartan medoxomil)  
Formulary Dossier

**OLMESARTAN MEDOXOMIL PRE-CLINICAL BIBLIOGRAPHY**  
(continued)

- Li Z, Chen XD, Ni SK, et al. Olmesartan inhibits the expression of monocyte chemoattractant protein-1 and tumor necrosis factor-alpha and improves vascular remodeling after vascular injury in mouse. *Chinese Journal of Traumatology*. 2004;7(1):56-61.
- Lijinsky W, Reuber MD. Pathologic effects of chronic administration of hydrochlorothiazide, with and without sodium nitrite, to F344 rats. *Toxicol. Ind. Health*. 1987; 3:413-422.
- Lin Y, Matsumura K, Kagiya S, et al. Chronic administration of olmesartan attenuates the exaggerated pressor response to glutamate in the rostral ventrolateral medulla of SHR. *Brain Res.* 2005;1058(1-2):161-6.
- Liu HW, Iwai M, Takeda-Matsubara Y, et al. Effect of estrogen and AT1 receptor blocker on neointima formation. *Hypertension*. 2002; 40(4): 451-7, and 448-50.
- Liu N, Shimizu S, Ito-Ihara T, et al. Angiotensin II Receptor Blockade Ameliorates Mesangioproliferative Glomerulonephritis in Rats through Suppression of CTGF and PAI-1, Independently of the Coagulation System. *Nephron Exp Nephrol*. 2007;105(3):e65-74.
- Ma SF, Anraku M, Iwao Y, et al. Hydrolysis of Angiotensin II Receptor Blocker Prodrug Olmesartan Medoxomil by Human Serum Albumin and Identification of Its Catalytic Active Sites. *Drug metabolism and disposition*. 2005;33(12):1911-9.
- Makoto Mizuno, Mikio Kato, Chizuru Miyake, Toshio Sada, et al. Effects of CS-866 on nephropathy in diabetic rats. The 72nd Annual Meeting of the Japanese Pharmacological Society, Sapporo, Japan, 1999.
- Makoto Mizuno, Toshio Sada, Masahiro Ikeda, et al. Effect of CS-866 on Hypertensive Nephropathy in SHR and DOCA Rats. The 70th Annual Meeting of the Japanese Pharmacological Society, Chiba, Japan - *Japanese Journal of Pharmacology* 1997; 73(suppl 1).
- Makoto Mizuno, Toshio Sada, Masahiro Ikeda, et al. Pharmacology of CS-866, a novel nonpeptide angiotensin II receptor antagonist. *European Journal of Pharmacology* 1995; 285: 181-188.
- Makoto Usui, Kensuke Egashira, Hideharu Tomita et al. Important Role of Local Angiotensin II Activity Mediated via Type 1 Receptor in the Pathogenesis of Cardiovascular Inflammatory Changes Induced by Chronic Blockade of Nitric Oxide Synthesis in Rats. *Circulation*. 2000;101:305-310.
- Manotham K, Tanaka T, Matsumoto M, et al. Evidence of Tubular Hypoxia in the Early Phase in the Remnant Kidney Model. *Journal of the American Society of Nephrology*. 2004;15:1277-1288.
- Marshall TG, Lee RE, Marshall FE. Common Angiotensin Receptor Blockers may directly modulate the immune system via VDR, PPAR and CCR2b. *Theor Biol Med Model*. 2006;3(1):1.
- Masao Takemoto, Kensuke Egashira, Hideharu Tomita, et al. Chronic Angiotensin-Converting Enzyme Inhibition and Angiotensin II Type 1 Receptor Blockade. *Hypertension* 1997; 30:1621-1627.
- Matsubara H, Shibasaki Y, Okigaki M, Mori Y, Masaki H, et al. Effect of angiotensin II type 2 receptor on tyrosine kinase Pyk2 and c-Jun NH2-terminal kinase via SHP-1 tyrosine phosphatase activity: evidence from vascular-targeted transgenic mice of AT2 receptor. *Biochem Biophys Res Commun*. 2001;282(5):1085-91.
- Matsuda M, Kanda Y, Kawasaki F, et al. Combination Use of Azelnidipine and Olmesartan Increases Insulin Sensitivity in Spontaneous Hypertensive Rats. ADA, June 2005. [Poster]
- Matsumoto H. Toxicity studies of losartan potassium. Angiotensin II receptor antagonist(5th report), oral development toxicity study in rabbits. *The Clinical Report* .1994; 28:4043-4051.
- Mazzolai L, Duchosal MA, Korber M, et al. Endogenous Angiotensin II Induces Atherosclerotic Plaque Vulnerability and Elicits a Th1 Response in ApoE / Mice. *Hypertension*. 2004;44:277-282.

AZOR™ (amlodipine and olmesartan medoxomil)  
Formulary Dossier

**OLMESARTAN MEDOXOMIL PRE-CLINICAL BIBLIOGRAPHY**  
(continued)

Merlos M., Rabasseda X., Silvestre JS. Discovery of new angiotensin II receptor antagonists. A review of pharmacological studies. *Methods & Findings in Experimental & Clinical Pharmacology*. Vol 20(9) (pp 805-815), 1998.

Mima A, Matsubara T, Abe H, et al. Angiotensin II receptro blocker inhibits the expression of Smad1 in Diabetic Nephropathy. ADA, June 2005. [Poster]

Mima A, Matsubara T, Arai H, et al. Angiotensin II-dependent Src and Smad1 signaling pathway is crucial for the development of diabetic nephropathy. ADA 66th Scientific Sessions (Laboratory Investigation/ABSTRACT). June 2006.

Min LJ, Mogi M, Li JM, et al. Aldosterone and Angiotensin II Synergistically Induce Mitogenic Response in Vascular Smooth Muscle Cells. *Circ. Res.* 2005;97:434 – 442.

Miura S, Fujino M, Hanzawa H, et al. Molecular Mechanism Underlying Inverse Agonist of Angiotensin ii Type 1 Receptor. The American Society for Biochemistry and Molecular Biology, May 2006.

Miura S, Fujino M, Saku K. Angiotensin II Receptor Blocker as an Inverse Agonist: A Current Perspective. *Current Hypertension Reviews*. 2005;1:115-121.

Miura SI, Kiya Y, Kanazawa T, et al. Differential Bonding Interactions of Inverse Agonists of Angiotensin II Type 1 Receptor in Stabilizing the Inactive-state. *Mol Endocrinol*. 2007 Sep 27.

Miyazaki M, Kim S, Sakonjo H, et al. A new angiotensin II type I receptor antagonist, CS-866, inhibited development of atherosclerotic lesion in monkeys fed a high-cholesterol diet. *J Hypertension*. 2000; 18(suppl 4):S86.

Miyazaki M, Takai S. Anti-atherosclerotic efficacy of olmesartan. *J of Hum Hypertens*. 2002; 16(Suppl. 2), SS7-S12.

Miyazawa N, Abe M, Nakayama K, et al. Chronic or acute interaction of angiotensin II on oxidative stress production by glucose pretreated mesangial cells. 21st Scientific Meeting of International Society of Hypertension (ISH), October 2006. [Abstract]

Mizuno M, Sada T, Kato M, et al. Renoprotective effects of blockade of angiotensin II AT 1 receptors in an animal model of type 2 diabetes. *Hypertension Research - Clinical Experimental*. 2002;25(2): 271-8.

Mortelmans K, Haworth S, Lawlor T, et al. Salmonella mutagenicity tests: II. Results from testing of 270 chemicals. *Environ. Mutagen*. 1986; 8(Suppl 7):1-119.

Mukai Y, Shimokawa H, Higashi M, Morikawa K, et al. Inhibition of renin-angiotensin system ameliorates endothelial dysfunction associated with aging in rats. *Arteriosclerosis, Thrombosis and Vascular Biology*. 2002 Sep 1;22(9): 1445-50.

Mura, Shinichiro. Angiotensin II receptor blocker as an agonist. A current perspective. *Curren Hypertension Reviews*. 2005;1(2):115-121.

Nagai Yukiko et al. Temporary Angiotensin II Blockade at the Prediabetic Stage Attenuates the Development of Renal Injury in Type 2 Diabetic Rats. *Journal of the American Society of Nephrology*. 2005;16:1-9.

Nagashima H, Uto K, Sakomura Y, Aoka Y, et al. An angiotensin-converting enzyme inhibitor, not an angiotensin II type-1 receptor blocker, prevents beta-aminopropionitrile monofumarate-induced aortic dissection in rats. *Journal of Vascular Surgery*. 2002; 36(4): 818-823.

AZOR™ (amlodipine and olmesartan medoxomil)  
Formulary Dossier

**OLMESARTAN MEDOXOMIL PRE-CLINICAL BIBLIOGRAPHY**  
(continued)

- Nakagomi-Hagihara R, Nakai D, Kawai K, et al. OATP1B1, OATP1B3 AND MRP2 ARE Involved in Hepatobiliary Transport of Olmesartan, A Novel Angiotensin ii Blocker. *Drug Metab Dispos.* 2006;34(5):862-9.
- Nakamoto T, Harasawa H, Akimoto K, et al. Effects of olmesartan medoxomil as an angiotensin II-receptor blocker in chronic hypoxic rats. *Eur J Pharmacol.* 2005;528(1-3):43-51.
- Nakamura H, Inoue T, Arakawa N, et al. Pharmacological and pharmacokinetic study of olmesartan medoxomil in animal diabetic retinopathy models. *Eur J Pharmacol.* 2005;512(2-3):239-46.
- Nakamura Y, Yoshiyama M, Omura T, Yoshida K, et al. Beneficial effects of combination of ACE inhibitor and angiotensin II type 1 receptor blocker on cardiac remodeling in rat myocardial infarction. *Cardiovascular Research.* 2003 Jan;57(1):48-54.
- Nakanishi K, Akimoto T, Kato H, et al. Olmesartan medoxomil, and angiotensin-II receptor blocker, ameliorates the pressure-natriuresis relationship in obese Zucker rats. 22nd Annual American Society of Hypertension Scientific Meeting, May 2007. [Abstract]
- Nakano N, Kobayashi N, Fukushima H, et al. Cardioprotective Mechanism of Combination Therapy with Selective Aldosterone Blocker and Angiotensin II Receptor Blocker Associated with JAK-STAT Pathway in Failing Rat Hearts. 17th Annual European Meeting on Hypertension, June 2007. [Poster]
- Nakayama S, Uehara Y, Urata H, Kinoshita A et al. Differential regulation of angiotensin II (All) forming enzymes by tissues in 2-kidney 1-clip (2K1C) hypertensive hamster. International Society of Hypertension 17th Scientific Meeting; Amsterdam, *Journal of Hypertension* 16, Sup. 2, S128, 1998.
- Namara I, Nakajima H, Fujishima H. A 12-month oral toxicity study of Trandolapril (RU44570) with 3-month recovery test in rats. *J. Toxicol. Sci.* 18(Suppl):37-92.
- Namikoshi T, Tomita N, Satoh M, et al. Olmesartan Ameliorates Renovascular Injury and Oxidative Stress in Zucker Obese Rats Enhanced by Dietary Protein. *Am J Hypertens.* 2007;20(10):1085-91.
- Nangaku M, Miyata T, Sada T, Mizuno M, Inagi R, et al. Anti-hypertensive agents inhibit in vivo the formation of advanced glycation end products and improve renal damage in a type 2 diabetic nephropathy rat model. *Journal of the American Society of Nephrology.* 2003 May;14(5):1212-22.
- Nimata M, Kishimoto C, Yuan Z, et al. Beneficial effects of olmesartan, a novel angiotensin II receptor type 1 antagonist, upon acute autoimmune myocarditis. *Mol Cell Biochem.* 2004;259(1-2):217-222.
- Nishio M, Sakata Y, Mano T, et al. Therapeutic effects of angiotensin II type 1 receptor blocker at an advanced stage of hypertensive diastolic heart failure. *Journal of Hypertension.* 2007;25(2):455-461.
- Nomura M, Nishii H, Ozaki Y, et al. An angiotensin II receptor blocker increases sexual behavior in type 2 diabetic mice. *Physiol Behav.* March 2007.
- Oberly TJ, Garriott ML. Influence of cytotoxicity on test results in the L5178Y TK+/- mouse lymphoma assay. *Environ. Molec. Mutagen.* 1996, 27:75-78.
- Ogata S, Sharyo S, Hinman DJ, et al. Renal effects of 26-week administration of olmesartan medoxomil/hydrochlorothiazide in rats. *The Journal of Toxicological Sciences.* 2004;29(1):37-46.
- Ohki R, Yamamoto K, Ueno S, et al. Effects of olmesartan, an angiotensin II receptor blocker, on mechanically-modulated genes in cardiac myocytes. *Cardiovascular Drugs & Therapy.* 2003;17(3):231-236.
- Ohshima Y, et al. Reproductive and developmental toxicity of TCV-116 in mice, rats and rabbits. *Jpn. Pharmacol.* 1996, 24:99-111.



AZOR™ (amlodipine and olmesartan medoxomil)  
Formulary Dossier

**OLMESARTAN MEDOXOMIL PRE-CLINICAL BIBLIOGRAPHY**  
(continued)

- Ohta T, Hasebe N, Tsuji S, et al. Unequal Effects of Renin Angiotensin System Inhibitors in Acute Cardiac Dysfunction Induced by Isoproterenol. *Am J Physiol Heart Circ Physiol*. 2004; 5: 1-42 (part 1 and 2).
- Ohtani K, Egashira K, Zhao G, et al. Angiotensin II AT1 Receptor Blockade Attenuates the Stent-Associated Neointimal Formation by Inhibiting Inflammation and by Reducing Recruitment/ Differentiation of Smooth Muscle Progenitor Cells in Rabbits and Monkeys. Univ, Sapporo, Japan Salutory, November 2004. [Abstract]
- Okada K, Hirano T, Ran J, et al. Olmesartan medoxomil, an angiotensin II receptor blocker ameliorates insulin resistance and decreases triglyceride production in fructose-fed rats *Hypertens Res*. 2004;27(4):293-299.
- Okada K, Minamino T, Tsukamoto Y, et al. Prolonged endoplasmic reticulum stress in hypertrophic and failing heart after aortic constriction: possible contribution of endoplasmic reticulum stress to cardiac myocyte apoptosis. *Circulation*. 2004;110(6):705-12.
- Oparil S. Are there meaningful differences in blood pressure control with current antihypertensive agents? *Am Journal of HTN* 2002. 15(1 Pt 2):14S-21S.
- Olmesartan Medoxomil/CS-866/Benevas. *Drugs Fut*. 2000, 25(11):1217-1218.
- Owen RA, Molon-Noblot S, Hubert MF, et al. The morphology of juxtaglomerular cell hyperplasia and hypertrophy in normotensive rats and monkeys given an angiotensin II receptor antagonist. *Toxicol. Pathol*. 1995, 22:606-619.
- Porteri E, Rodella L, Rizzoni D, et al. Effects of olmesartan and enalapril at low or high doses on cardiac, renal and vascular interstitial matrix in spontaneously hypertensive rats. *Blood Pressure*. 2005;14(3):184-92.
- Prieto-Carrasquero MC, Kobori H, Ozawa Y, et al. AT1 receptor-mediated enhancement of collecting duct renin in angiotensin II-dependent hypertensive rats. *American Journal of Physiology (Renal Fluid & Electrolyte Physiology)*. 2005;289(3):F632-7.
- Quadri L., Gobbin M., Monti L. Recent advances in antihypertensive therapy. *Current Pharmaceutical Design*, Vol 4(6) (pp 489-512), 1998.
- Ran J, Hirano T, Adachi M. Angiotensin II infusion increases hepatic triglyceride production via its type 2 receptor in rats. *J Hypertens*. 2005;23(8):1525-30.
- Ran J, Hirano T, Adachi M. Angiotensin II type 1 receptor blocker ameliorates overproduction and accumulation of triglyceride in the liver of Zucker fatty rats. *Am J Physiol Endocrinol Metab*. 2004;287(2):E227-32.
- Ran J, Hirano T, Fukui T, et al. Angiotensin II infusion decreases plasma adiponectin level via its type 1 receptor in rats: an implication for hypertension-related insulin resistance. *Metabolism*. 2006;55(4):478-88.
- Rodriguez-Iturbe B., Sato T. et al. AT1 Receptor Blockade Prevents Proteinuria, Renal Failure, Hyperlipidemia, and Glomerulosclerosis in the Imai Rat. *Kidney International*. 2004;66:668-675.
- Sada T, Mizuno M. Pharmacological Profiles and Clinical Effects of Olmesartan Medoxomil, a novel Angiotensin II Receptor Antagonist. *Pharmacology and Molecular Research Laboratories*. 2004;124(4):257-69.
- Sagawa K, Nagatani K, Komagata Y, et al. Angiotensin Receptor Blockers Suppress Antigen-Specific T Cell Responses and Ameliorate Collagen-Induced Arthritis in Mice. *Arthritis & Rheumatism*. 2005;52(6):1920-1928.

AZOR™ (amlodipine and olmesartan medoxomil)  
Formulary Dossier

**OLMESARTAN MEDOXOMIL PRE-CLINICAL BIBLIOGRAPHY**

(continued)

Sanada S, Kitakaze M, Node K, Takashima S, Ogai A, et al. Differential subcellular actions of ACE inhibitors and AT(1) receptor antagonists on cardiac remodeling induced by chronic inhibition of NO synthesis in rats. *Hypertension*. 2001;38(3):404-11.

Sandmann S, Li J, Fritzenkotter C, et al. Differential effects of olmesartan and ramipril on inflammatory response after myocardial infarction in rats. *Blood Pressure*. 2006;15(2):116-28.

Sasaki Y, Kanazawa M, Tufescu A, et al. Renoprotective and antihypertensive effects of combination therapy with a low protein diet (LPD), long-term exercise (EX) and olmesartan (OLS) in rats with renal ablation. 21st Scientific Meeting of International Society of Hypertension (ISH), October 2006. [Abstract]

Sato M, Iwanaga T, Mamada H, et al. Involvement of Uric Acid Transporters in Alteration of Serum Uric Acid Level by Angiotensin II Receptor Blockers. *Pharmaceutical Research*. June 2008.

Satoh K, Ichihara K, Landon EJ, et al. 3-Hydroxy-3-methylglutaryl-CoA reductase inhibitors block calcium-dependent tyrosine kinase pyk2 activation by angiotensin II in vascular endothelial cells. *J Biol Chem*. 2001;276(19):15761-7.

Satoru Sharyo, Shinya Sehata, Hideki Sagisaka, et al. Effects of Sodium Supplementation on CS-866 Repeated-Dose Toxicity Study on Rats. July 25-26, 1996. The 23rd Japanese Society of Toxicology - *Journal of Toxicological Sciences* 1996; 21(5): 364.

Schildcrout JS, Margolin BH, Zeiger E. Predicting rodent carcinogenicity using potency measures of the in vitro sister chromatid exchange and chromosome aberration assays. *Environ. Molec. Mutagen*. 1999, 33:59-64.

Seko Y. Effect of an angiotensin II receptor blocker olmesartan on the development of murine acute myocarditis caused by coxsackievirus B3. *Clinical Science*. 2006;110(3):379-86.

Shao J, Nangaku M, Miyata T, Inagi R, et al. Imbalance of T-cell subsets in angiotensin II-infused hypertensive rats with kidney injury. *Hypertension*. 2003 Jul;42(1):31-8. Epub 2003 May 27.

Shibasaki Y, Matsubara H, Nozawa Y, Mori Y, et al. Angiotensin II type 2 receptor inhibits epidermal growth factor receptor transactivation by increasing association of SHP-1 tyrosine phosphatase. *Hypertension*. 2001 Sep;38(3):367-72.

Shokei Kim, Toshio Sada, Makoto Mizuno, et al. Effects of Angiotensin AT<sub>1</sub> Receptor Antagonist on Volume Overload-Induced Cardiac Gene Expression in Rats. *Hypertension Research*, 1997; 20:133-142.

Soichiro Fujiyama, Hiroaki Matsubara, Yoshihisa Nozawa, et al. Angiotensin AT<sub>1</sub> and AT<sub>2</sub> Receptors Differentially Regulate Angiopoietin-2 and Vascular Endothelial Growth Factor Expression and Angiogenesis by Modulating Heparin Binding-Epidermal Growth Factor (EGF)-Mediated EGF Receptor Transactivation. *Circulation Research*. 2001;88:22-29.

Sonta T, Inoguchi T, Matsumoto S, et al. In vivo imaging of oxidative stress in the kidney of diabetic mice and its normalization by angiotensin II type 1 receptor blocker. *Biochem Biophys Res Commun*. 2005;330(2):415-22.

Sonta T, Inoguchi T, Matsumoto S, et al. Noninvasive imaging of increased oxidative stress in kidney of diabetic mice by a novel technique of in vivo ESR-CT measurement. American Diabetes Association Scientific Meeting [842-P], June 2004.

Spalding JW, et al. Recent retrospective and prospective studies in the TG.AC and p53<sup>±</sup> transgenic mouse lines on chemicals completed in the two-year NTP bioassay. *The Toxicologist* 1999, 48:253.

AZOR™ (amlodipine and olmesartan medoxomil)  
Formulary Dossier

**OLMESARTAN MEDOXOMIL PRE-CLINICAL BIBLIOGRAPHY**  
(continued)

Suda O, Tsutsui M, Morishita T, Tanimoto A, et al. Long-term treatment with N(omega)-nitro-L-arginine methyl ester causes arteriosclerotic coronary lesions in endothelial nitric oxide synthase-deficient mice. *Circulation*. 2002; 106(13):1729-35.

Suda O, Tsutsui M. et al. Asymmetric Dimethylarginine Causes Arteriosclerotic Lesions in Endothelial Nitric Oxide Synthase Deficient Mice: Involvement in the Renin-Angiotensin System and Oxidative Stress. *Arteriosclerosis, Thrombosis, and Vascular Biology*. 2004; 1-7.

Sugawara A, Kudo M, Salto A, et al. Differential Effects of Angiotensin-Converting Enzyme Inhibitor and Angiotensin II Receptor Blocker on Gene Expression in Vascular Endothelial Cells Revealed by DNA Microarray analyses. 21st Scientific Meeting of International Society of Hypertension (ISH), October 2006. [Abstract]

Sugie T, Kagaya Y, Morihiko T, Yahagi H, Takahashi C, et al. Should increasing the dose or adding an AT1 receptor blocker follow a relatively low dose of ACE inhibitor initiated in acute myocardial infarction? *Cardiovascular Research*. 2003 Jun 1;58(3):611-20.

Suzuki J, Iwai M, Li Z, et al. Effect of combination of calcium antagonist, azelnidipine, and AT1 receptor blocker, olmesartan, on atherosclerosis in apolipoprotein E-deficient mice. *J Hypertens*. 2005;23(7):1383-9.

Tada T, Nawata J, Wang H, et al. Pulsatile Pressure accelerates Vascular Smooth Muscle Migration through IP3-Receptor. 21st Scientific Meeting of International Society of Hypertension (ISH), October 2006. [Abstract]

Takahashi H, Takeishi Y, Miyamoto T, et al. Protein kinase C and extracellular signal regulated kinase are involved in cardiac hypertrophy of rats with progressive renal injury. *European Journal of Clinical Investigations*. 2004;34(2):85-93.

Takahashi T, Konta T, Takasaki S, et al. An Angiotensin II Type-1 Receptor Blocker, Olmesartan Medoxomil, Attenuates Lipid Peroxidation in Renal Injury Induced by Subtotal Nephrectomy. *Clin Exp Nephrol*. 2007;11(3):202-8.

Takai S, Jin D, Sakaguchi M, et al. The regressive effect of an angiotensin II receptor blocker on formed fatty streaks in monkeys fed a high-cholesterol diet. *Journal of Hypertension*. 2005;23(10):1879-86.

Takai S, Kim S, Sakonjo H, et al. Mechanisms of angiotensin II type 1 receptor blocker for anti-atherosclerotic effect in monkeys fed a high-cholesterol diet. *Journal of Hypertension*. 2003;21(2):361-9.

Takai S, Kim S, Sakonjo H, Miyazaki M. Mechanisms of angiotensin II type 1 receptor blocker for anti-atherosclerotic effect in monkeys fed a high-cholesterol diet. *Journal of Hypertension*. 2003 Feb;21(2):361-9.

Takao T, Horino T, Kagawa T, et al. Angiotensin Type 1 Receptor Blockers Protect Human Renal Proximal Tubular Epithelial Cells from Albumin-Induced Cell Damage. 21st Scientific Meeting of International Society of Hypertension (ISH), October 2006. [Abstract]

Takaori Kasuo., Ohasi Yasuko., Kitagawa Taizo. et al. Effect of an angiotensin II Type 1 Receptor Antagonist on Carotid Injury in Rats. *Memoirs of Osaka Kyoiku University Series III a Natural Science & Applied Science*. 46(1). Aug., 1997. 95-100.

Takayanagi M, Sano N, Takikawa H. Biliary excretion of olmesartan, an angiotensin II receptor antagonist, in the rat. *Journal of Gastroenterology & Hepatology*. 2005;20(5):784-8.

AZOR™ (amlodipine and olmesartan medoxomil)  
Formulary Dossier

**OLMESARTAN MEDOXOMIL PRE-CLINICAL BIBLIOGRAPHY**  
(continued)

- Takemoto M, Egashira K, Usui M, Tsutsui H. Angiotensin AT1 receptor blockade versus inhibition of bradykinin breakdown: effects on cardiac remodeling in rats induced by long-term blockade of nitric oxide synthesis.
- Tanabe A, Naruse M, Shimamoto Y, et al. Effects of AT1 Receptor Antagonist (ARB) on Insulin Resistance in Experimental Model Rats for Metabolic Syndrome. Endocrine Society 89th Annual Meeting. [Abstract]
- Tanabe Y, Morikawa Y, Kato T, et al. Effects of olmesartan, an AT1 receptor antagonist, on hypoxia-induced activation of ERK1/2 and pro-inflammatory signals in the mouse lung. *Naunyn Schmiedeberg Arch Pharmacol.* 2006;374(3):235-48.
- Tanaka T, Matsumoto M, Inagi R, et al. Induction of protective genes by cobalt ameliorates tubulointerstitial injury in the progressive Thy1 nephritis. *Kidney Int.* 2005;68(6):2714-25.
- Tanifuji C, Suzuki Y, Geot WM, et al. Reactive Oxygen Species-Mediated Signaling Pathways in Angiotensin II-Induced MCP- Expression of Proximal Tubular Cells. *Antioxid Redox Signa.* 2005;7(9-10):1261-8.
- Taniyama Y, Morishita R, Matsumoto K, Ogihara T. Prevention of myocardial injury and fibrosis by angiotensin II blockade in cardiomyopathic hamster: potential contribution of a novel anti-fibrotic factor, hepatocyte growth factor (HGF). 71st Scientific Sessions of the American Heart Association; Dallas, *Circulation* 98, No. 17, Sup., I179, 1998 (Abstract: 921).
- Togashi N, Ura N, Higashiura K, Murakami H, et al. The contribution of skeletal muscle tumor necrosis factor- $\alpha$  to insulin resistance and hypertension in fructose-fed rats. *J Hypertens.* 2000;18(11):1605-10.
- Tokuyama H, Hayashi K, Matsuda H, et al. Differential Regulation of Elevated Renal Angiotensin II in Chronic Renal Ischemia. *Hypertension.* 2002; 40:34-40.
- Tokuyama H, Hayashi K, Matsuda H, et al. Distinct Role of intrarenal cyclo-oxygenase-  $\frac{1}{2}$  in chronic unilateral renal ischemia. *Nephron* Sept 2002; 92(1): 183-91.
- Tomita Hideharu, Egashira Kensuke et al. Angiotensin type 1 receptor antagonist prevents vascular monocyte infiltration and monocyte chemoattractant protein-1 expression in the rat heart induced by chronic inhibition of nitric oxide synthesis. 70th Scientific Sessions of the American Heart Association; Orlando, *Circulation* Vol. 96, Sup 8, I176;1997 (Abstract: 975).
- Toshio Sada, Makoto Mizuno, Chizuru Miyata, et al. Long-term blockade of angiotensin II (AT1) receptors suppresses Ca<sup>2+</sup>-dependent tone of aorta in spontaneously hypertensive rats. The 72nd Annual Meeting of the Japanese Pharmacological Society, Sapporo, Japan - *Japanese Journal of Pharmacology* 1999; 79(1).
- Tsuda M, Iwai M, Li JM, et al. Inhibitory Effects of AT1 Receptor Blocker, Olmesartan, and Estrogen on Atherosclerosis Via Anti-Oxidative Stress. *Hypertension.* 2005;45:1-7.
- Uemura H, Ishiguro H, Nakaigawa N, et al. Angiotensin II receptor blocker shows antiproliferative activity in prostate cancer cells: A possibility of tyrosine kinase inhibitor of growth factor. *Mol Cancer Ther.* 2003;2:1139-1147.
- Urakaze M, Kobashi C, Asamizu S, et al. Olmesartan Inhibits IL-8-Induced ERK Phosphorylation in Human Adipocytes In Vitro. 21st Scientific Meeting of International Society of Hypertension (ISH), October 2006. [Abstract]
- Valencia R, Mason JM, Woodruff RC, et al. Chemical mutagenesis testing in Drosophila. III. Results of 48 coded compounds tested for the National Toxicology Program. *Environ. Mutagen.* 1985; 7:325-348.

AZOR™ (amlodipine and olmesartan medoxomil)  
Formulary Dossier

**OLMESARTAN MEDOXOMIL PRE-CLINICAL BIBLIOGRAPHY**  
(continued)

- Wang L-X, Ideishi M, Yahiro E, Urata H, et al. Mechanism of the cardioprotective effect of inhibition of the renin-angiotensin system on ischemia/reperfusion-induced myocardial injury. *Hypertens Res*. 2001;24(2):179-87.
- Wennemuth G, Aumu"ller G. Angiotensin II-mediated calcium signals and mitogenesis in human prostate stromal cell line hPCPs. *British Journal of Pharmacology*. 2005;144:3–10.
- Xu HL, Yoshida K, Wu XM, M Kohzuki. Effects of CS-866, an angiotensin II receptor antagonist, in 5/6 nephrectomized spontaneously hypertensive rats. Nippon Jinzo Gakkai Shi. *Japanese Journal of Nephrology*. 2001;43(7):580-8.
- Yamada A, Maeda K, Kamiyama E, et al. Multiple Human Isoforms of Drug Transporters Contribute to the Hepatic and Renal Transport of Olmesartan, A Selective Antagonist of the Angiotensin II AT1-receptor. *Drug Metab Dispos*. September 2007.
- Yamagishi S, Takeuchi M, Inoue H. Olmesartan medoxomil, a newly developed angiotensin II type 1 receptor antagonist, protects against renal damage in advanced glycation end product (age)-injected rats. *Drugs Exp Clin Res*. 2005;31(2):45-51.
- Yamagishi SI, Matsui T, Nakamura K, et al. Pigment-epithelium-derived factor (PEDF) inhibits angiotensin-II-induced vascular endothelial growth factor (VEGF) expression in MOLT-3 T cells through anti-oxidative properties. *Microvasc Res*. April 2006.
- Yamaguchi K, Ura N, Murakami H, et al. Olmesartan Ameliorates Insulin Sensitivity by Modulating Tumor Necrosis Factor- $\alpha$  and Cyclic AMP in Skeletal Muscle. *Hypertens Res*. 2005; 28:773-778.
- Yamamoto E, Yamashita T, Tanaka T, et al. Pravastatin Enhances Beneficial Effects of Olmesartan on Vascular Injury of Salt-Sensitive Hypertensive Rats, via Pleiotropic Effects . *Arteriosclerosis, Thrombosis & Vascular Biology*. 2006.
- Yamamoto T, Kajikuri J, Watanabe Y, et al. Chronic nitroglycerine administration reduces endothelial nitric oxide production in rabbit mesenteric resistance artery. *Br J Pharmacol*. 2005;146(4):534-42.
- Yamashita C, Hiyashi T, Mori T, et al. Nifedipine Added to the Combination with Pravastatin and Olmesartan Shows Preventive Effect on Hypoxia-Induced Left Ventricular Remodeling in Diabetes Mellitus 56th Annual American College of Cardiology Scientific Session, March 2007. [Poster]
- Yamashita T, Sekiguchi A, Kato T, et al. Angiotensin type 1 receptor blockade prevents endocardial dysfunction of rapidly paced atria in rats. *J Renin Angiotensin Aldosterone Syst*. 2007;8(3):127-32.
- Yao L, Kobori H, Rahman M, et al. Olmesartan improves endothelin-induced hypertension and oxidative stress in rats. *Hypertens Res*. 2004;27(7):493-500.
- Yasuda N, Akazawa H, Qin Y, et al. Structural and functional analysis of Olmesartan as an inverse agonist for Angiotensin type 1 receptor. 21st Scientific Meeting of International Society of Hypertension (ISH), October 2006. [Abstract]
- Yasuko Ito, Makoto Mizuno, Mikio Kato, Toshio Sada, et al. Effects of CS-866 on anti-thy 1.1 antibody induced glomerulonephritis in rats. The 72nd Annual Meeting of the Japanese Pharmacological Society, Sapporo, Japan - *Japanese Journal of Pharmacology* 1999; 79(1).
- Yatabe J, Sanada H, Yatabe MS, et al. Olmesartan Attenuates the Activation of ERK by cyclic Mechanical Strain in Rat Mesangial Cells in the absence of Angiotensin II. 21st Scientific Meeting of International Society of Hypertension (ISH), October 2006. [Abstract]

AZOR™ (amlodipine and olmesartan medoxomil)  
Formulary Dossier

**OLMESARTAN MEDOXOMIL PRE-CLINICAL BIBLIOGRAPHY**  
(continued)

Yokota T, Utsunomniya K, Taniguchi K, et al. Olmesartan Ameliorates expression of platelet-derived growth factor (PDGF) and Vascular Endothelial Growth Factor (VEGF) Induced by Glycated Albumin in Porcine Retinal Pericytes as an Inverse Agonist. ADA, June 2005. [Poster]

Yoshiaki Taniyama, Ryuichi Morishita, Hironori Nakagami et al. Potential Contribution of a Novel Anti-Fibrotic Factor, Hepatocyte Growth Factor, to Prevention of Myocardial Fibrosis by Angiotensin II Blockade in Cardiomyopathic Hamsters. *Circulation*. 2000;102(2):246-52.

Yoshida J, Yamamoto K, Mano T, et al. AT1 Receptor Blocker Added to ACE Inhibitor Provides Benefits at Advanced Stage of Hypertensive Diastolic Heart Failure. *Hypertension*. 2004;43:1-6.

Yoshida J, Yamamoto K, Mano T, Sakata Y, Nishikawa N, Nishio M, Ohtani T, Miwa T, Hori M, Masuyama T. AT1 Receptor Blocker Added to ACE Inhibitor Provides Benefits at Advanced Stage of Hypertensive Diastolic Heart Failure. *Hypertension*. 2004 March;43:1-6.

Yoshida K, Xu HL, Kawamura T, Ji L, Kohzuki M. Chronic Angiotensin-Converting Enzyme Inhibition and Angiotensin II Antagonism in Rats with Chronic Renal Failure. *Journal of Cardiovascular Pharmacology*. 2002; 40(4): 533-42.

Yu X, Murao K, Imachi H, et al. Regulation of Scavenger Receptor Class BI Gene Expression by Angiotensin II in Vascular Endothelial Cells. *Hypertension*. April 2007.

Yuan Z, Nimata M, Okabe TA, et al. A Novel Angiotensin II Receptor Type 1 Antagonist, Suppressed Cytotoxic Myocardial Injury in Autoimmune Heart Failure. *Am J Physiol Heart Circ Physiol*. 2005 May 6.

Zaman T, Fujii S, Goto D, et al. Effects of Attenuation of Angiotensin II Mediated JNK Signaling on Coronary Perivascular Fibrosis Associated with Insulin Resistance and Obesity. Univ, Sapporo, Japan Salutory, November 2004. [Abstract]

Zaman T., Fujii S. et al. Salutory Effects of Attenuation of Angiotensin II on Coronary Perivascular Fibrosis Associated with Insulin Resistance and Obesity. *Journal of Molecular and Cellular Cardiology*. 2004; 37:529-535.

Zeiger E, Haseman JK, Shelby MD, et al. Evaluation of four in vitro genetic toxicity tests for predicting rodent carcinogenicity: confirmation of earlier results with 41 additional chemicals. *Environ. Molec. Mutagen*. 1990; 16(Suppl 18):1-14.

AZOR™ (amlodipine and olmesartan medoxomil)  
Formulary Dossier

**OLMESARTAN MEDOXOMIL PRE-CLINICAL BIBLIOGRAPHY**  
(continued)

Zhang GX, Ohmori K, Nagai Y, et al. Role of AT1 receptor in isoproterenol-induced cardiac hypertrophy and oxidative stress in mice. *J Mol Cell Cardiol.* Feb 2007.

Zhao Q, Ishibashi M, Hiasa K, et al. Essential role of vascular endothelial growth factor in angiotensin II-induced vascular inflammation and remodeling. *Hypertension.* 2004;44(3):264-70.